

Page numbers in italics denote figures; those followed by a "t" denote tables; and those followed by a "c" denote capsules

| A  |
|--|
| Abbott Sanitary Counting Tray, 257<br>Abbreviated New Drug Application<br>bioavailability data, 177<br>bioequivalence data, 185<br>for generic drugs, 20 |
| Abelia Abertifacient, 753 Abradant, 753  |
| Absorbent, 753   |
| Absorption<br>bases, for ointments, 317–318<br>drug  |
| active transport, 121, 171–173, 409<br>administration route and, 43, 47, 50,<br>61, 66, 167–169, 187–197, 334, 510,<br>625, 745, 768                     |
| amorphous, 112<br>area under curve representing,<br>183–184  |
| barriers, 66, 121, 166, 168–172, 192, 193, 307, 343, 345, 368, 369 bioequivalence factors, 177–187   |
| controlled, 51, 175, 345, 349, 504,<br>514   |
| crystalline, 173, 175, 176, 276, 424, 550, 554   |
| dissolution and, 50, 51, 112, 117, 121, 172–177  |
| dosage form and, 43, 172–177<br>in elderly persons, 197  |
| epicutaneous, 187t<br>from extended-release products,  |
| 302, 309, 309<br>facilitating diffusion, 171   |
| fate after, 172<br>gastric emptying time, 172, 173,<br>183, 192, 274   |
| hydration state and, 177 from injections, 194  |
| from liposomes, 740, 741<br>membrane permeability and, 121   |
| nasal, 196<br>ocular, 196  |
| otic, 196<br>parenteral, 188t, 193   |
| particle size and, 112<br>passive diffusion, 168–171   |
| percutaneous. See Transdermal  |
| drug delivery systems<br>principles, 167   |
| protein binding and, 197, 204<br>pulmonary, 197, 642, 710  |
| rate, 135, 135c–137c<br>rectal, 188t, 192–193, 369–371   |
| in salt form, 177 serum concentration and, 61,   |
| 183–184<br>small intestinal, 172–177,  |
| 192, 368 specialized transport mechanisms, 171–172   |
| surface area and, 174–175<br>time course, 180  |
| transdermal. See Transdermal drug  |

delivery systems

```
urethral, 386-387
    vaginal, 735
  in packaging, 12
Absorption, distribution, metabolism/
      biotransformation and excretion
      (ADME) studies, 47
Abuse, drug regulations, 17
Acacia, as emulsifying agent, 469
Accelerated stability testing, 254
Acesulfame potassium, 158
Acetaminophen
  with codeine, 247
  rapidly dissolving tablet, 267
  in syrup, 416
Acetazolamide, in nuclear medicine, 659
Acid(s)
  diluted, 434-436
  dissociation constant, 124c
  pK<sub>a</sub> values, 170t
  weak, solubility, 399t
Acidification, of urine, 460t
Acidifier, 753
Acidifying agent, 151t
Acquired immunity, 579
Action letters, in new drug approval, 70
Activated protein C, recombinant, 692
Activation energy of, 454c
Active immunity, 579–580
Active ingredient, 763
Active pharmaceutical ingredients
  definition, 83
  regulations, 129
Active transport, 171
Acute myeloid leukemia, gemtuzumab
      ozogamicin for, 683t
Acute toxicity studies, 48
Adalimumab (Humira), 700
Additive properties, drugs, 517c
ADD-Vantage System, for intravenous
      administration, 534, 534
Adenocarcinoma, diagnostic imaging of,
      satumomab pendetide in, 705
Adenosine deaminase (ADA)
      deficiency, gene therapy, 679
Adenosine, in nuclear medicine,
      659-660
Adhesive layer, in transdermal drug
      delivery systems, 349
Administration routes
  dosage forms and
    parenteral, 188t, 193
    rectal, 188t, 192-193
  inhalation. See Inhalation
      administration route)
  nasal. See Nasal administration route
  ocular. See Ocular administration
      route
  oral. See Oral administration route
  otic. See Otic administration route
  for powders, 195
  rectal. See Rectal administration route
  sublingual, 188, 188t, 191
  for systemic effects, 187
  urethra, 188t, 197
  vaginal. See Vaginal administration
      route)
Administration timing, dosage and, 66
```

Adrenergic drug, 753 Adrenocorticosteroids anti-inflammatory, 753 salt-regulating, 753 Adrenocorticotropic hormone, 753 Adsorbent, 151t, 753 Adults, immunization schedules for, 603 Adult-senior use packaging, 93, 93-94 Advair Diskus, 229 Adverse reactions product recall and, 24 reporting, 35, 36, 56 Advisory Committee on Immunization Practices, 578 Aerosols, 763 administration methods, 498-500 advantages, 490-491 compressed gas, 494 containers for, 494-495 definition, 488 examples, 498, 499t filling of, 496-498 foam, 763 rectal administration, 500-501 vaginal administration, 500-501 inhalation examples, 498, 499t metered dose, 496 particle size of, 175 labeling, 498 packaging, 498 partial pressure and, 488c–489c particle sizes, 490 powders, 227, 228, 500 principles, 491–492 product concentrate, 492 propellants for, 151t blends, 493 compressed gas, 494 filling of, 496-498 physical properties, 491t in three-phase systems, 493-494 in two-phase systems, 493 rectal, 500–501, 501 space, 493 storage, 498 surface, 493 systems, 493 testing, 498 three-phase, 493-494 topical, 500 two-phase, 493 vaginal, 500-501 valve assembly of, 494 Agar, as emulsifying agent, 469 Age. See also Elderly persons; Pediatric patients in clinical studies, 54-55 dosage form design, 104 drug dosage, 62-63 drug metabolism and, 200 flavor preferences and, 152t gastrointestinal absorption and, 173 Aggregation, of globules, 475–476 Agitation, in syrup preparation, 418 Agonist, 753

| Air  | American Society of Health System                | lyophilization, 533   |
|--|--|---|
| displacement of, 151t                          | Pharmacists (ASHP)                               | manufacturing, 533  |
| humidifiers for, 627, 627–628                  | pharmaceutical care statement, 28                | in ophthalmic preparations, 607                                 |
| suspension tablet coating, 288–289,            | quality assurance guidelines, 543–544            | in suspension, 464  |
| 288–289  | Amino acids, in parenteral nutrition,            | Antibody(ies)   |
| Airway, 228. See also Inhalation               | 563  | chimeric, 704   |
| administration route                           | Ammonia, radiolabeled, in positron               | human antimouse, 677  |
| Albumin, drug binding to, 197                  | emission tomography, 656–657                     | man-made, 677   |
| Alcohol  | Amorphous compounds, in dosage                   | monoclonal. See Monoclonal                                      |
| dehydrated, 401                                | form design, 112                                 | antibodies  |
| diluted, 402                                   | Amorphous forms, dissolution of,                 | Anticholesterol agent, 754                                      |
| in pediatric patients, 406                     | 175–177  | Anticholinergic agents, 754                                     |
| as preservative                                | Amorphous polyethylene terephthalate             | gastrointestinal absorption and, 173                            |
| in syrups, 402                                 | glycol (APET), 91                                | Anticholinesterase antidote, 754                                |
| as solvents                                    | Amoxicillin, USP, 13, 14                         | Anticoagulants  |
| in tinctures, 424                              | Amphiphilic colloid, 479                         | antagonist, 754   |
| Alcohol-abuse deterrent, 753                   | Amphotec, 741                                    | from biotechnology, 685   |
| Aldehyde dehydrogenase inhibitor, 753          | Amphotericin B liposome formulation,             | Anticonvulsant, 754   |
| Aldesleukin (Proleukin), 682, 683t,            | 741  | Antidepressant, 754   |
| 697–698  | Ampicillin, anhydrous form, 177                  | Antidiabetic drug, 754  |
| Alendronate sodium tablets, 257                | Ampuls, 763                                      | Antidiarrheal drug, 754   |
| Alginic acid, as gelling agent, 481            | definition, 523                                  | Antidiuretic drug, 754  |
| Alkalinization                                 | filling and sealing of, 537, 538                 | Antidote, 754   |
| systemic, 411                                  | for parenterals, 523                             | for radiation exposure, 658, 658–659                            |
| of urine, 201                                  | sizes of, 535                                    | Antidyskinetic drug, 755  |
| Alkalinizer, 753                               | Amyl nitrite                                     | Antieczematic drug, 754   |
| Alkalinizing agents, 151t                      | inhalation, 628                                  | Antiemetic drug, 755  |
| Alkylating agents, 753                         | volatility, 106                                  | Antiepileptic drug, 755   |
| Allergic reactions                             | Anabolic steroid, 753                            | Antiestrogen, 755   |
| sulfites, 140                                  | Anakinra (Kineret), 698, 698                     | Antifibrinolytics, 755  |
| to vaccines, 591, 595                          | Analeptic, 753                                   | Antifilarial drug, 755  |
| All-in-one granulation methods, for            | Analgesics, 753                                  | Antiflatulent, 755  |
| tablet manufacture, 279, 279–280               | in ear preparations, 630                         | Antifungal agents, for ophthalmic use,                          |
| Allogeneic tumor vaccines, 586                 | patient-controlled analgesia, 511                | 532, 607  |
| Alloying substance, for tablet coatings,       | Anaphylaxis, in vaccination, 591                 | Antifungal preservative, 151t                                   |
| 287  | Andreasen pipette, in particle size              | Antigens  |
| Alpha glucosidase inhibitor, 753               | measurement, 218c                                |   |
| Alpha particle, 640                            |  | lymphocyte recognition of, 579 skin, diagnostic, 601–602        |
| Alpha receptor agonist, 753                    | Androderm patch, 348t, 355<br>Androgen, 753      | tumor associated, in vaccine                                    |
| Alpha receptor antagonist, 753                 | Anemia, erythropoietin in, 690                   |   |
| Alpha reductase inhibitor, 753                 | Anesthetics, for ophthalmic use, 607             | preparation, 585  |
| Alpha tocopherol as antioxidants, 140          | Angiotensin-converting enzyme                    | Antiglaucoma agent, 755<br>Antigonadotropin, 755                |
| Alpha value, of drugs, 197                     | inhibitor (ACE) inhibitor, 43, 754               | Antigout agent, 755   |
| Alprostadil, in urethral                       | Angle of repose, of powders, 218c–219c           | Antihemophilic drug, 755  |
| microsuppository, 386–387                      | Anhydrous forms, dissolution, 177                | Antihemophilic factors (Kogenate,                               |
| Alteplase (Activase), recombinant, 706,        | Aniline derivatives, as colorants, 158           | Recombinate), 686–687   |
| 706–707  | Animal powders, definition, 216                  | Antiherpes agent, 755   |
| Aluminum Acetate Topical Solution, 428         | Animals  | Antihistamines  |
| Aluminum, for aerosol containers, 486          | drug sources, 39                                 | 4   |
|  | drug testing                                     | elixirs, 421–423, 422t  |
| Aluminum hydroxide                             | metabolism studies, 47                           | syrups, 413t<br>Antihyperlipidemic drug, 755                    |
| in antacids, 459, 462, 463<br>as gels, 485–486 | pharmacology studies, 46, 47                     |   |
| in suspensions, 463                            | *  | Antihypertensive drug, 755                                      |
| Aluminum subacetate topical solution,          | toxicology studies, 47–48, 49<br>Anorexiant, 754 | Antihypocalcemic drug, 755<br>Antihypoglycemic drug, 755        |
| 428  | Anorexic, 754                                    | Anti-infective drug, 755  |
| Aluminum tubes, 325                            | Antacid, 754                                     | Anti-inflammatory drug, 755                                     |
| Alzet osmotic pump, 734–735                    | chewable tablet, 282                             | Antileishmanial drug, 755                                       |
| AmBisome, 741                                  | side effects, 459                                | Antileprotic drug, 755  |
| Ambulatory care setting, pharmacy              | suspensions, 459, 462–463                        | Antilipemic drug, 755   |
| practice in, 26                                | Antagonist, 754                                  | Antimpenne drug, 755 Antimalarial drug, 755                     |
| Amebicide, 753                                 | Anthelmintic drug, 754                           | Antimanic drug, 755   |
| American Association of Colleges of            | Anthrax  | Antimetabolite, 755   |
| Pharmacy Janus Commission,                     | adsorbed vaccine, 588                            | Antimicrobial preservative, 151t                                |
| pharmaceutical care statement, 29              | in bioterrorism, 600–601                         | Antimigraine agent, 755   |
| American Association of Pharmaceutical         | Antiacne agent, 754                              | Anti-motion sickness agent, 755                                 |
| Scientists, code of ethics, 31                 | Antiadrenergic, 754                              | Antimuscarinic drug, 755  |
| American National Standards Institute,         | Antiamebic, 754                                  | Antinauseant drug, 755  |
| 15   | Antiandrogen, 754                                | Antineoplastic drug, 755  |
| American Pharmacists Association               | Antianemic, 754                                  | Antineoplastic drug, 755  |
| code of ethics for pharmacists, 30–31          | Antianginal agent, 754                           | Antipeuritic drug, 755 Antiparasitic drug, 755                  |
| drug standards, 8–15                           | Antiarrhythmic, 754                              | Antiparasitic drug, 755 Antiparkinsonian (antidyskinetic) drug, |
| mission statement, 28                          | Antiarthytitilic, 754 Antiarthritic, 754         | 755   |
| National Formulary of, 11                      | Antibacterial, 754                               | Antiperistaltic drug, 755                                       |
| Nuclear Pharmacy Practice                      | Antibiotics, 754                                 | Antiplatelet agent, 755   |
| Guidelines, 661                                | in ear preparations, 632                         | Antiprogestin, 756  |
| pharmacoutical care statement 28               | in frozon form 535                               | Antiprotezzal 756   |

| Antipruritic drug, 756  | Pastonial and atoxing test LICD E21   | Diagraphy along dura muchusta definition  |
|---|---|---|
| Antipsoriatic drug, 756   | Bacterial endotoxins test, USP, 531 Bacterial filtration  | Bioequivalent drug products, definition,<br>184   |
| Antipsoriatic drug, 756   | of ophthalmic preparations, 606   | Biologic characterization, of drugs,  |
| Antipyretic drug, 756   | of parenterals, 524   | 44–50, 46, 47   |
| Antirachitic drug, 756  | Bacterial vaccine, 583, 589t–591t   | Biologic indicators, of sterilization   |
| Antiretroviral drug, 756  | Bacteriostatic sodium chloride injection,   | effectiveness, 527  |
| Antirheumatic drug, 756   | USP, 516  | Biologic membrane, drug penetration   |
| Antirickettsial drug, 756   | Bacteriostatic water for injection, USP, 516  | of, 168   |
| Antischistosomal drug, 756  | Balance   | Biologicals, 578-604  |
| Antiscorbutic drug, 756   | electronic, 776–778   | for active immunity, 583-586  |
| Antiseborrheic drug, 756  | prescription, 776–778   | administration, toxicity associated   |
| Antisecretory agent, 756  | Barbiturate, 423, 756   | with, 587–600   |
| Antisense technology, 679   | Barium hydroxide, in pyrogen removal,   | animal immune sera, 587, 588t-591t  |
| Antispasmodic drug, 756   | 531   | antitoxins, 587, 590t   |
| Antithyroid agent, 756  | Barium sulfate suspension, 463–464  | antivenins, 587, 590t   |
| Antitreponemal, 756   | Base(s)   | bioterrorism and, 600-601   |
| Antitrichomonal drug, 756   | for ointments, 317–318  | case studies, 602–603   |
| Antitubercular drug, 756  | pKa values, 170t, 192   | definition, 578   |
| Antitussive drug, 756   | suppositories   | diagnostic skin antigens, 601–602   |
| Antivenins, 587, 590  | bioavailability and, 371  | examples, 588t–591t   |
| Antiviral agents  | classification, 372   | expiration date, 581  |
| for ophthalmic use, 607   | determination of amount required,   | handling, 581–583   |
| prophylactic, 632   | 378–382   | human immune sera and globulins,  |
| Antiviral drug, 756   | Basiliximab (Simulect), 700–701   | 587   |
| Antixerophthalmic drug, 756   | Batch, definition, 78   | immunity and, 578–581   |
| Anxiolytic drug, 756  | Batchwise control, definition, 78   | labeling, 581   |
| Apothecary, first, 5  | Beads, coated, for extended-release   | mixed, 583, 586   |
| Apothecary system, 770, 772–773, 777  | products, 303, 303  | for passive immunity, 587, 588t–591t  |
| Apparent density, of powders, 219c  | Becaplermin (platelet-derived growth  | production, 581   |
| Appearance, of pharmaceutical   | factor) (Regranex), 694   | shipping, 581–583   |
| ingredients and excipients, 150,<br>151t  | Becquerel, 640 Beeswax, as hardening agent, 373   | side effects, 587, 591  |
| Aqueous film-coating, 287   | Belladonna alkaloid, 756  | storage, 581–583<br>toxoids, 580, 586, 589  |
| Area under the concentration-time curve,  | Bentonite, as emulsifying agent, 470  | vaccines. See Vaccines and  |
| bioavailability assessment, 178   | Bentonite magma, NF, 485  | vaccination   |
| Aromas, for tablets, 287  | Benzalkonium chloride   | Biologics license applications, 72  |
| Aromatic water, 763   | as emulsifying agents, 470  | Biopharmaceutics, 166   |
| Arthritis, rheumatoid   | as preservative, 608  | Biotechnology products  |
| adalimumab for, 700, 700  | Benzodiazepine, 756   | anticoagulants, 685   |
| anakinra for, 698, 698  | Benzoic acid, 322, 399t, 484  | antisense drugs, 679  |
| Artificial colloids, 479  | Beta particles, 641   | case studies, 721–723   |
| Artificial sweeteners, 158  | Beta receptor agonist, 756  | classification, 682   |
| Artificial tears, 607–608   | Beta receptor antagonist, 756   | clotting factors, 682   |
| Ascorbic acid, antioxidant, 140   | Beta-blockers, 510  | and any atimately time for those 607 600  |
| Ascorbyl palmitate, antioxidant, 140  |   | colony stimulating factors, 687–689,  |
|   | for ophthalmic use, 607   | 688   |
| Aseptic, definition, 763  | Betaseron (interferon beta-1b), 695–696   | 688<br>definition, 672  |
| Aseptic, definition, 763<br>Aseptic processing, 763   | Betaseron (interferon beta-1b), 695–696<br>Bile, drug excretion, 200  | 688<br>definition, 672<br>erythropoietins, 690–693  |
| Aseptic, definition, 763<br>Aseptic processing, 763<br>Aspartame, 158   | Betaseron (interferon beta-1b), 695–696<br>Bile, drug excretion, 200<br>Binders, for tablets, 264   | 688<br>definition, 672<br>erythropoietins, 690–693<br>Food and Drug Administration office   |
| Aseptic, definition, 763<br>Aseptic processing, 763<br>Aspartame, 158<br>Aspirin  | Betaseron (interferon beta-1b), 695–696<br>Bile, drug excretion, 200<br>Binders, for tablets, 264<br>Binding sites, for drugs, 207  | 688 definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714  |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173   | Betaseron (interferon beta-1b), 695–696<br>Bile, drug excretion, 200<br>Binders, for tablets, 264<br>Binding sites, for drugs, 207<br>Binding, to plasma proteins, 197  | 688 definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694   |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366  | Betaseron (interferon beta-1b), 695–696<br>Bile, drug excretion, 200<br>Binders, for tablets, 264<br>Binding sites, for drugs, 207<br>Binding, to plasma proteins, 197<br>Bingham bodies, 454c  | definition, 672<br>erythropoietins, 690–693<br>Food and Drug Administration office<br>on, 713–714<br>fusion inhibitors, 693–694<br>future, 711–713  |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479   | Betaseron (interferon beta-1b), 695–696<br>Bile, drug excretion, 200<br>Binders, for tablets, 264<br>Binding sites, for drugs, 207<br>Binding, to plasma proteins, 197<br>Bingham bodies, 454c<br>Bioanalytical research laboratories, 210  | definition, 672<br>erythropoietins, 690–693<br>Food and Drug Administration office<br>on, 713–714<br>fusion inhibitors, 693–694<br>future, 711–713<br>growth factors, 695   |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703   | Betaseron (interferon beta-1b), 695–696<br>Bile, drug excretion, 200<br>Binders, for tablets, 264<br>Binding sites, for drugs, 207<br>Binding, to plasma proteins, 197<br>Bingham bodies, 454c<br>Bioanalytical research laboratories, 210<br>Bioavailability   | 688 definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711   |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time  | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694   |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184   | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696  |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426  | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181   | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699  |
| Aseptic, definition, 763 Aseptic processing, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245  | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694  |
| Aseptic, definition, 763 Aseptic processing, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184  | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715,   |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580  | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185  | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t  |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641  | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179   | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production   |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629   | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679  |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641  | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179   | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production   |
| Aseptic, definition, 763 Aseptic processing, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616   | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717–720t production gene therapy in, 678–679 monoclonal antibodies in,   |
| Aseptic, definition, 763 Aseptic processing, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t  | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706  |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586  | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182   | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade/antisense  |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586 Automation, for drug manufacture,  | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182 time of peak, 182–183   | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade/antisense technique, 679   |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586 Automation, for drug manufacture, 82–83, 83, 84 Autoxidations, in drug degradation, 134 Avoirdupois system, 770, 773   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182 time of peak, 182–183 Bioavailable fraction first-pass effect, 187 formula for, 187   | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade/antisense technique, 679 peptide technology, 679 polymerase chain reaction in, 678 recombinant DNA techniques,   |
| Aseptic, definition, 763 Aseptic processing, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586 Automation, for drug manufacture, 82–83, 83, 84 Autoxidations, in drug degradation, 134  | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182 time of peak, 182–183 Bioavailable fraction first-pass effect, 187 formula for, 187 Bioequivalence  | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade/antisense technique, 679 peptide technology, 679 polymerase chain reaction in, 678 recombinant DNA techniques, 674–676   |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586 Automation, for drug manufacture, 82–83, 83, 84 Autoxidations, in drug degradation, 134 Avoirdupois system, 770, 773   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182 time of peak, 182–183 Bioavailable fraction first-pass effect, 187 formula for, 187 Bioequivalence absolute, 183  | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade/antisense technique, 679 peptide technology, 679 polymerase chain reaction in, 678 recombinant DNA techniques, 674–676 protein-based, advantages, 711  |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586 Automation, for drug manufacture, 82–83, 83, 84 Autoxidations, in drug degradation, 134 Avoirdupois system, 770, 773   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182 time of peak, 182–183 Bioavailable fraction first-pass effect, 187 formula for, 187 Bioequivalence absolute, 183 assessment, 180–184  | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade / antisense technique, 679 peptide technology, 679 polymerase chain reaction in, 678 recombinant DNA techniques, 674–676 protein-based, advantages, 711 rasburicase, 710   |
| Aseptic, definition, 763 Aseptic processing, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586 Automation, for drug manufacture, 82–83, 83, 84 Autoxidations, in drug degradation, 134 Avoirdupois system, 770, 773 Avonex (interferon beta-1a), 696  | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182 time of peak, 182–183 Bioavailable fraction first-pass effect, 187 formula for, 187 Bioequivalence absolute, 183 assessment, 180–184 definition, 177  | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade/antisense technique, 679 peptide technology, 679 polymerase chain reaction in, 678 recombinant DNA techniques, 674–676 protein-based, advantages, 711 rasburicase, 710 reimbursement issues, 716                                       |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586 Automation, for drug manufacture, 82–83, 83, 84 Autoxidations, in drug degradation, 134 Avoirdupois system, 770, 773 Avonex (interferon beta-1a), 696  B B lymphocytes, in immunity, 579   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182 time of peak, 182–183 Bioavailable fraction first-pass effect, 187 formula for, 187 Bioequivalence absolute, 183 assessment, 180–184 definition, 177 differences, 178                         | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade/antisense technique, 679 peptide technology, 679 polymerase chain reaction in, 678 recombinant DNA techniques, 674–676 protein-based, advantages, 711 rasburicase, 710 reimbursement issues, 716 tissue plasminogen activators,        |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586 Automation, for drug manufacture, 82–83, 83, 84 Autoxidations, in drug degradation, 134 Avoirdupois system, 770, 773 Avonex (interferon beta-1a), 696  B B lymphocytes, in immunity, 579 Bacille Calmette-Guérin (BCG), USP, 586 | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182 time of peak, 182–183 Bioavailable fraction first-pass effect, 187 formula for, 187 Bioequivalence absolute, 183 assessment, 180–184 definition, 177 differences, 178 factors affecting, 186t | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade/antisense technique, 679 peptide technology, 679 polymerase chain reaction in, 678 recombinant DNA techniques, 674–676 protein-based, advantages, 711 rasburicase, 710 reimbursement issues, 716 tissue plasminogen activators, 706–708 |
| Aseptic, definition, 763 Aseptic processing, 763 Aspartame, 158 Aspirin absorption, 173 bioavailability, in suppositories, 366 Association colloid, 479 Asthma, omalizumab, 703 Astringent, 756 ophthalmic use, 607 Atomizers, 426 Atropine, sodium chloride equivalent of, 610–611, 611t Attenuated vaccines, 580 Auger electron, 641 Aural, 763 Autoclave, in parenteral sterilization, 522 Autoinjection systems, 749, 749–750 Autologous tumor vaccines, 585–586 Automation, for drug manufacture, 82–83, 83, 84 Autoxidations, in drug degradation, 134 Avoirdupois system, 770, 773 Avonex (interferon beta-1a), 696  B B lymphocytes, in immunity, 579   | Betaseron (interferon beta-1b), 695–696 Bile, drug excretion, 200 Binders, for tablets, 264 Binding sites, for drugs, 207 Binding, to plasma proteins, 197 Bingham bodies, 454c Bioanalytical research laboratories, 210 Bioavailability area under serum concentration time curve, 183–184 assessment, 180–181 in capsule products, 245 concentration time curve, 183–184 definition, 185 FDA submission requirements, 179 intranasal, 629 in ophthalmic preparations, 616 of oral drugs, factors affecting, 186t peak height, 181–182 time of peak, 182–183 Bioavailable fraction first-pass effect, 187 formula for, 187 Bioequivalence absolute, 183 assessment, 180–184 definition, 177 differences, 178                         | definition, 672 erythropoietins, 690–693 Food and Drug Administration office on, 713–714 fusion inhibitors, 693–694 future, 711–713 growth factors, 695 human DNase 1, 710–711, 711 human growth hormone, 694 interferons, 695–696 interleukins, 696–699 patient education on, 693–694 pharmacist information on, 714, 715, 716, 717t–720t production gene therapy in, 678–679 monoclonal antibodies in, 699–706 nucleotide blockade/antisense technique, 679 peptide technology, 679 polymerase chain reaction in, 678 recombinant DNA techniques, 674–676 protein-based, advantages, 711 rasburicase, 710 reimbursement issues, 716 tissue plasminogen activators,        |

| Bioterrorism, 600-601                      | Caloric requirements                      | vaginal, 388                            |
|--|---|---|
| Biotransformation. See Metabolism,         | in enteral nutrition, 565                 | weight variation, 253–254               |
| drug                                       | in parenteral therapy, 562                | wetted, 241                             |
| Bisacodyl, in suppositories, 367t          | Cancer                                    | Captopril, in nuclear medicine, 659     |
|  |   |   |
| Bisphosphonate, 756                        | detection, satumomab pendetide in,        | Carbamide peroxide                      |
| Bitter taste, 157                          | 705                                       | cerumen-removing solutions, 630         |
| Black box warnings, 18                     | interleukins for, 696–699                 | dental solutions, 433                   |
| Blending, of powders, 223–225, 224–225     | pain in, fentanyl lollipop for, 293       | Carbohydrates                           |
| Blinded studies, 51–52, 58                 | positron emission tomography, 654         | as emulsifying agents, 469              |
| Blister packaging, 91                      | rituximab for, 704                        | requirements, in enteral nutrition, 565 |
| Blood, isotonicity, 609                    | trastuzumab for, 705-706                  | Carbomer resins, 482                    |
| Blowers, for powders, 228, 228             | vaccines for, 585–586                     | Carbon dioxide, radiolabeled, in        |
| Board of Pharmaceutical Specialties,       |   |   |
|  | Candida albicans skin test antigen, 591t, | positron emission tomography,           |
| 26–27                                      | 602                                       | 657                                     |
| Body fluids, isotonicity, 609              | Caplets, 190, 763                         | Carbon-11, in positron emission         |
| Body membranes, classification, 168        | Capping defect, in tablets, 281, 282      | tomography, 656                         |
| Body surface area, dosage                  | Capromab pendetide (ProstaScint), in      | Carbonic anhydrase inhibitor, 756       |
| determination and, 64, 65                  | prostate cancer, 649–650                  | Carbowaxes, 481                         |
| Body weight, dosage determination          | Capsules, 763                             | Carcinoembryonic antigen detection,     |
| and, 64                                    | case studies, 259–261                     | radiopharmaceuticals in, 643            |
| Boiling point, of mixtures, 517            | colorants for, 240, 247                   | Carcinogenicity studies, 49             |
| Bolus, 763                                 | commercially available, 255, 255t         |   |
|  | a. a                                      | Cardiac depressant, antiarrhythmic, 757 |
| Bone disease, radiopharmaceuticals         | compendial requirements, 251–254          | Cardiac glycoside, 757                  |
| in, 647                                    | content uniformity, 254                   | Cardiotonic, 422t                       |
| Bone marrow, granulocyte macrophage        | counting, 255–256, 257                    | Carmustine implant, after brain tumor   |
| colony-stimulating factor for,             | definition, 240                           | resection, 745, 745-746, 746            |
| 687–688                                    | diluents for, 244, 245                    | Cascade impaction, particle size        |
| Bone metabolism regulator, 756             | disintegrants for, 244                    | measurement, 221                        |
| Boric acid                                 | disintegration test for, 251–252          | Castor oil emulsion, 477                |
| in ear preparations, 631                   | dispensers for, 251                       | Catapres TTS, 347t, 353                 |
|  |   |   |
| in ophthalmic preparations, in             | dissolution test for, 252–253             | Catecholamine synthesis inhibitor, 757  |
| tonicity considerations, 610               | extended-release                          | Catheter                                |
| Bottle (Forbes) method, 474–475            | coated particles in, 240                  | Broviac/Hickman, 513                    |
| Bottom spray method, for tablet coating,   | glidants for, 245, 247                    | central venous, 513                     |
| 289  | hard gelatin                              | indwelling plastic, 513                 |
| Botulism antitoxins, 590                   | cleaning, 249, 252                        | Cauterizing agent, 757                  |
| Botulism, in bioterrorism, 600, 601        | contents of, examples, 247                | Caventou, Joseph, 7                     |
| Brain imaging                              | design, 242–243, 243                      | Cavitation, in phonophoresis, 731       |
| . 0 9                                      | dissolution, 241                          |   |
| drug interference with, 667t               |   | Cell cultures, in preclinical testing,  |
| positron emission tomography, 658          | drug absorption from, 241                 | 44–45                                   |
| Brain, tumor resection in, carmustine      | filling, 247–249, 248, 249                | Center for Biologics Evaluation and     |
| implant for, 745, 745–746, 746             | formulations for, 244–247, 245            | Research, 56, 581                       |
| Breast cancer, trastuzumab for, 705        | gelatin properties, 243                   | Center for Drug Evaluation and          |
| Breast-feeding, radiopharmaceuticals       | liquids in, 246                           | Research, 23, 56, 541                   |
| during, 668                                | manufacturing, 241–243, 242, 243          | Centipoise, 615                         |
| British Pharmacopeia (BP), 8               | passage of, tracking methods, 241         | Centistokes, 615                        |
| Bronchodilator, 756                        | polishing, 249, 252                       | Central compartment, in                 |
| Brookfield Viscometer, 450, 450            | preparation, 243–244                      | pharmacokinetics, 204                   |
|  |   |   |
| Broviac catheter, for repeated injection,  | sealing, 249, 252                         | Central vein parenteral nutrition       |
| 513  | sizes, 244, 244, 245                      | (central vein TPN), 565                 |
| Buccal system, mucoadhesive                | small capsules or tablets inserted        | Central venous catheters, 513           |
| testosterone, 733–734                      | into, 245                                 | Centrally acting drug, 757              |
| Buccal tablets, 265–266                    | swallowing, 241                           | Cephalosporin, 757                      |
| Buffer capacity, 138, 138c-140c            | transit time, 241                         | Certification                           |
| Buffering agent, 151t                      | uses, 240                                 | colorants, 158                          |
| drug stability, 138                        | weight variation, 254                     | finished pharmaceuticals, 78            |
| ophthalmic preparation, 612                | identification, 239                       | pharmacist, 27                          |
| Buildings, for drug manufacture,           | inspection, 255–256, 257                  | Cerumen-removing solutions, 630         |
| regulations on, 79                         |   |   |
|  | labeling, 254                             | Cesium, radioactive, exposure to,       |
| Bulk powders, 229–230                      | lubricants for, 245, 247                  | antidote for, 658, 658–659              |
| Bulk volume, of powders, 219c–220c         | moisture permeation test for, 254–255     | Chelating agent, 152t, 757              |
| Bulkiness, of powders, 219c                | official, examples, 255–256, 255t         | Chemical abstracts service (CAS)        |
| Butyl hydroxy anisole, as antioxidant, 140 | opaquants for, 247                        | registry number, 12                     |
|  | opening, 242, 243, 256                    | Chemical enhancers, transdermal drug    |
| ^  | for oral use, 256                         | delivery systems, 344                   |
| C  | overview, 240                             | Chemotherapy                            |
| Calamina Linimant 475                      | packaging, tamper-evident, 249            | bone marrow, granulocyte                |
| Calairm archanata 451 450 462              |   |   |
| Calcium carbonate, 451, 459, 462           | rectal, 388                               | macrophage colony-stimulating           |
| Calcium channel blocker, 756               | soft gelatin                              | factor for, 687–688                     |
| Calcium hydroxide                          | preparation, 250–251, 253, 254            | drugs for, safe handling, 543           |
| solubility, 397                            | utilization, 251                          | Gliadel Wafer, 745, 745–746, 746        |
| topical solution, 397                      | weight variation, 253–254                 | neutropenia in, granulocyte colony-     |
| Calcium, in parenteral nutrition, 563      | stability testing, 254                    | stimulating factor, 687-688             |
| Calcium soaps, emulsion preparation,       | storage, 255–256, 257                     | thrombocytopenia in, oprelvekin         |
| 475  | substances added to, 251                  | for, 698                                |
| Calibration, of suppository molds, 378     | swallowing, 239, 240                      | Chewable tablets, 266                   |
| Tr   |   |   |

| Children. See Pediatric patients          | soft, 621–622   | Cytomegalovirus (CMV)                     |
|---|---|---|
| Child-resistant packaging, 93–94, 94      | storage solutions for, 622  | immunoglobulin, 590t                      |
| Chimeric antibodies, 704                  | types, 620  | retinitis, fomivirsen, 685                |
| Chloramphenicol, crystalline vs.          | wetting solutions for, 623  | ,   |
| * 4 .5_                                   | Containers, 79–80   | <b>B</b>                                  |
| amorphous, 175                            |   | D   |
| Chloride, in oral rehydration solutions,  | for aerosols, 494–495   | D 1: 1 (7 ) 701                           |
| 410                                       | definition, 80  | Daclizumab (Zenapax), 701                 |
| Chlorobutanol, as preservative, 162, 609  | for packaging, 88–93, 89, 90, 91t, 543  | Darbepoetin alfa (Aranesp), 691–692,      |
| Chlorofluorocarbons, in aerosols, 491     | regulations, 83   | 692                                       |
| Cholelitholytic, 757                      | Continental (dry gum) method, 473   | Decongestant, 757                         |
| Cimetidine                                | Contraceptives, 757   | Delayed-release products                  |
| discovery, 41, 42                         | oral, 745   | definition, 301                           |
|   |   |   |
| in nuclear medicine, 659                  | transdermal, 355  | drug release rate from, 309, 309          |
| Citric acid                               | vaginal preparations, 735, 735  | enteric coatings, 307                     |
| in effervescent granulated salts, 232,    | Corticosteroid, 757   | examples, 307t                            |
| 233c-234c                                 | Coupling agents, for phonophoresis, 730   | Demulcent, 757                            |
| sodium citrate and, 411                   | Creams, 764   | Dental caries prophylactic drug, 757      |
| Clarifying agent, 152t                    | definition, 323   | Dentin desensitizer, 757                  |
| Cleaning solutions, for contact lenses,   | dermatologic, 327–329, 330t, 331t, 332t   | Depigmenting agent, 757                   |
|   |   |   |
| 621–623                                   | packaging, 325–326  | Detergent, 757                            |
| Clearance, drug, 207                      | rectal, 334–335, 336t   | Diagnostic aid, 757                       |
| Clinical studies                          | vaginal, 336, 336–337, 336t   | Diagnostic imaging                        |
| controls, 58–59                           | Cresol, as preservative, 162  | radiopharmaceuticals for, 642,            |
| designs, 58–60, 60                        | Crinone gel, 737  | 642-643                                   |
| drug dosage and terminology, 59-66,       | Crohn disease, infliximab for, 702  | satumomab pendetide in, 704               |
| 60, 61, 62t, 63t, 65                      | Crossover designs   | Dietary Supplement Health and             |
| phases, 56–58, 57t, 58                    | for bioequivalence studies, 178   | Education Act (DSHEA) of 1994,            |
|   |   |   |
| Clinical trial materials, regulations, 84 | clinical studies, 58–59, 60   | 21–22, 96                                 |
| Clonidine, transdermal drug delivery      | Cross-tolerance, 66   | Diffusion                                 |
| systems, 347t, 353–354                    | Crystalline structure, 106  | coefficient, 118–119c, 169, 484, 738      |
| Clotting factors, recombinant, 686–687    | dissolution of, 173   | facilitated, 171                          |
| Code of Federal Regulations (CFR), 23     | Crystals, polymorphism of, 176  | layer, 172                                |
| Collodion, 764                            | Cups, for measuring doses, 776  | passive, 168–171                          |
| Colloidal dispersion, 764                 | Curie, 640  | Digestive aid, 757                        |
|   |   | ~ .                                       |
| Colloidal silicon dioxide, 483            | Current Good Compounding Practices,   | Digoxin                                   |
| Colony-stimulating factors,               | 85–88   | discovery, 8                              |
| recombinant, 687–690                      | Current Good Manufacturing Practice   | dosage regimen for, 201                   |
| Color Additive Petition, for contact lens | for active pharmaceutical ingredients,  | in elixirs, 424                           |
| manufacture, 621                          | 83  | half-life, 201                            |
| Colorants, 152t                           | for biologics, 84   | Diluents, for capsules, 244, 245          |
| capsules, 160                             | for clinical trial materials, 84  | Dinoprostone, in vaginal insert, 736, 737 |
|   |   |   |
| certification status, 158                 | for finished pharmaceuticals  | Dipyridamole, in nuclear medicine,        |
| for contact lenses, 621                   | automation, 82–83, 83, 84   | 659–660                                   |
| formulation, 160                          | control of components, containers,  | Disinfectant, 757                         |
| lake pigments, 160                        | and closures, 79–80   | Disinfection, of contact lenses, 622      |
| powdered drug, 160                        | distribution procedures, 81   | Diskhaler inhalation device, 229          |
| for tablets, 160                          | equipment, 79   | Disposable contact lenses, 620            |
| Comminution, of drugs, 222–223, 223       | facilities, 77t, 79   | Dissociation constants                    |
| Compliance                                | general provisions, 76–78   | in dosage form design, 123                |
|   | holding procedures, 81  | 0   |
| finished pharmaceuticals, 78              |   | passive diffusion, 169                    |
| packaging for, 94                         | information technology, 82–83,  | Dissolution                               |
| Components, drug product                  | 83, 84  | of amorphous forms, 175–177               |
| definition, 78                            | labeling control, 80–81   | hydration state and, 177                  |
| regulation, 83                            | laboratory controls, 81–82  | of salt form, 177                         |
| Comprehensive Drug Abuse Prevention       | organization, 78–79   | Diuretic drug, 757                        |
| and Control Act of 1970, 16, 17           | packaging control, 80-81, 82t   | Diuretics                                 |
| Compressed tablets. See Tablets,          | personnel requirements, 78–79   | loop, 561, 660, 759                       |
| compressed                                | process controls, 80  | potassium-sparing, 561, 760               |
| Concomitant drug therapy, 66              |   | thiazide, 201, 761                        |
|   | production controls, 80   |   |
| Contact lenses                            | records management and reports,   | Diverticulum, Meckel's, nuclear           |
| advantages, 620                           | 82  | imaging of, 659                           |
| care, 621                                 | returned drug products, 82  | Divided powders, 230                      |
| cleaners for, 621–622                     | salvaged drug products, 82  | DNA technology, recombinant, in           |
| clinical considerations, 623-624          | for labeling, 94–97   | biotechnology products,                   |
| color additives to, 621                   | for manufacturing in pharmacies, 85   | 674–676                                   |
| combination solutions for, 623            | for medical devices, 84–85  | Dopamine receptor agonist, 757            |
| disadvantages, 620                        | noncompliance with, 85  | Dosage form(s), 764                       |
|   |   |   |
| disinfecting solutions for, 622           | for packaging, 88–93, 89, 90, 91t   | administration routes, 66                 |
| disposable, 620                           | for pharmaceutical excipients, 83   | nasal, 196                                |
| drug interactions, 623                    | standards, 76, 76t  | ocular, 196                               |
| extended wear, 620                        | for storage, 97   | oral, 196                                 |
| hard, 620                                 | for transportation, 97  | otic, 196                                 |
| neutralizing solutions for, 622           | Cycloplegic drug, 757   | parenteral, 188t, 193                     |
| rigid gas permeable, 620                  | Cyclotron, in radiopharmaceutical   | rectal, 188t, 192–193                     |
| rinsing solutions for, 622                | production, 657, 662  | novel. See Novel dosage forms and         |
| soaking solutions for, 545, 623           | Cystic fibrosis, human DNase I for, 710   | drug delivery systems                     |
|   | _ , , 100, _ 100 |   |

| Dosage form design                                  | loading of, in rapidly dissolving                            | definition, 420                                     |
|---|--|---|
| biopharmaceutical considerations                    | tablets, 267   | digoxin, 424  |
| passive diffusion, 168–171                          | new. See New drug(s)   | medicated, 421                                      |
| specialized transport mechanisms,                   | pharmacy, heritage of, 4–8                                   | nonmedicated, 421                                   |
| 171–172<br>general considerations                   | physical properties, 104 regulation. See Drug regulation and | phenobarbital, 423–424                              |
| microscopic examination, 107                        | control  | preparation, 421 proper administration and use, 425 |
| oral dosage devices, 104, 105                       | semisynthetic, 38  | vs. syrups, 420                                     |
| various forms, 104, 104                             | uses, 2  | theophylline, 420                                   |
| importance, 160                                     | Drug discovery and drug design, 3                            | Emetic drug, 758                                    |
| low usual doses, 189t                               | FDA's definition, 43-44                                      | Empirical approach, to dosage regimen,              |
| pharmaceutical ingredients and                      | goal drug in, 40   | 208   |
| excipients  | lead compound (prototype) in, 41–43                          | Emulsifying agent, 152t                             |
| appearance, 156                                     | methods, 40–41   | Emulsions, 764                                      |
| palatability, 156–161<br>types, 150, 151t–155t, 155 | nomenclature and, 44<br>prodrugs in, 43                      | aggregation, 475–476<br>coalescence, 475–476        |
| preformulation studies                              | sources of new drugs, 38–40                                  | creaming, 476                                       |
| complex organic molecules, 131,                     | Drug Enforcement Administration, 17                          | definition, 465                                     |
| 131c-133c   | Drug Listing Act of 1972, 18, 20                             | examples, 477–478                                   |
| dissolution, 117-118, 118c-120c                     | Drug Price Competition and Patent                            | fat, intravenous, 196                               |
| heat of vaporization, 107, 107c–108c                | Term Restoration Act of 1984,                                | injectable, 560                                     |
| hydrates and solvates, 124–125, 125c                | 20–21  | inversion, 470                                      |
| melting point depression, 108, 109c                 | Drug product, definition, 78                                 | oil-in-water, 465                                   |
| membrane permeability, 121                          | Drug product recall, 24                                      | oral, 477   |
| microscopic examination, 107                        | Drug regulation and control                                  | packaging, 476                                      |
| organic ester considerations, 128,<br>128c–129c     | Code of Federal Regulation. See Code of                      | phases, 476   |
| organic salt considerations,                        | Federal Regulations Comprehensive Drug Abuse                 | preparation<br>bottle (Forbes) method, 474–475      |
| 126–127, 127c                                       | Prevention and Control Act of                                | continental (dry gum) method, 473                   |
| particle size, 112                                  | 1970, 16, 17   | emulsifying agents for, 468–470                     |
| partition coefficient, 121, 121c–123c               | Dietary Supplement Health and                                | English (wet gum) method, 473                       |
| pH, 161   | Education Act (DSHEA) of 1994,                               | foam, 501   |
| phase rule, 108, 110c–111c                          | 21–22  | HLB system in, 470, 471c-472c                       |
| physical description, 106–107                       | Drug Listing Act of, 18, 20                                  | homogenizer in, 473                                 |
| pKa/dissociation constants, 123,                    | Drug Price Competition and Patent                            | mechanical blender in, 470                          |
| 123c-124c   | Term Restoration Act of 1984,                                | microemulsion, 475                                  |
| polymorphism, 112                                   | 20–21  | mortar and pestle in, 451                           |
| potency-designated active                           | drug product recall, 24                                      | in situ soap method, 475                            |
| pharmaceutical ingredients,                         | Durham-Humphrey Amendment of                                 | preservatives for, 476                              |
| 129–130, 130c–131c<br>solubility, 112               | 1951, 16<br>Federal Food, Drug, and Cosmetic                 | purpose, 465–466<br>stability, 475–476              |
| stability considerations                            | Act of 1938, 15–25   | temperature and, 476                                |
| accelerated stability studies,                      | Food and Drug Administration                                 | theories of emulsification, 466–468,                |
| 147–150, 148c–149c                                  | Modernization Act of 1997, 22–23                             | 467c-468c   |
| degradation mechanism and, 134                      | Kefauver-Harris Amendments of                                | topical, 477–478                                    |
| drug product, 134–135                               | 1962, 16–17  | water-in-oil, 465                                   |
| drug substance, 147                                 | Prescription Drug Marketing Act of                           | Enalapril   |
| enhancement, 137–142, 138c–140c                     | 1987, 21   | metabolites of, 199                                 |
| excipient, 147                                      | Dry powder inhaler, 764                                      | as prodrug, 43                                      |
| expiration date, 147                                | Durham-Humphrey Amendment of                                 | Encapsulating agent, 152t                           |
| kinetics, 134–135                                   | 1951, 16   | Endothelin receptor antagonist, 758                 |
| $Q_{10}$ method of shelf life estimation, 137       | _  | Enema, 764<br>Enfuvirtide (Fuzeon), 693, 693–694    |
| rate reactions, 135, 135c–137c                      | E  | English (wet gum) method, 473                       |
| shelf life, 137                                     | Ear, drug administration to. See Otic                        | Enzyme, 758   |
| stress testing, 147                                 | administration route   | Epicutaneous administration route, 187t             |
| testing, 147–148                                    | Early formulation studies, 50-52                             | Epidermal growth factor, 758                        |
| Dressing, 764                                       | Effervescent dosage form, 764                                | Epinephrine, autoinjection systems, 749,            |
| Drops   | Effervescent granulated salts, 232,                          | 749–750   |
| eye. See Ophthalmic preparations                    | 233c-234c  | Ergot alkaloid, 758                                 |
| nasal, 628–629                                      | Elderly persons  | Erythropoietins, recombinant, 690–693               |
| Orotrecogin alfa (activated) (Xigris),              | absorption in, 197   | Estradiol, transdermal drug delivery                |
| 692–693, 693<br>Drug(s)                             | in clinical studies, 54–55                                   | systems, 347t, 354–355                              |
| administration routes. See                          | dosage form design for, 105                                  | Estring vaginal drug delivery system, 735, 736      |
| Administration routes                               | drug dosage for, 459<br>gastrointestinal absorption in, 169  | Estrogen, 758                                       |
| chemical properties, 104                            | packaging for, 94  | Eutectic mixtures                                   |
| definition, 2                                       | protein binding in, 197                                      | capsules, 246                                       |
| delivery system, 764                                | Electrical current, in iontophoresis, 728                    | of powders, 223                                     |
| discovery. See Drug discovery and                   | Electroendosmotic flow, in                                   | Expectorant, 758                                    |
| drug design   | iontophoresis, 729   | Expiration date                                     |
| early, 5–6, 6                                       | Elimination, drug, 200, 201                                  | for biologics, 581                                  |
| formulations, 4                                     | Elixirs, 764   | for finished pharmaceuticals, 81                    |
| homeopathic, 12                                     | antihistamine, 421–423, 422t                                 | Extended-release products                           |
| interactions with food, 177                         | barbiturate, 423   | advantages, 300, 301t                               |
| legend, 16  | components, 420  | chemical complexes, 305                             |

| control bonds, granules, or                          | dermatalogie 224t                                 | UMC CoA reductors inhibitor 758  |
|--|---|--|
| coated beads, granules, or<br>microspheres, 303, 303 | dermatologic, 334t                                | HMG-CoA reductase inhibitor, 758<br>Hormone, 758   |
|  | examples, 484–487                                 | The state of the s |
| definition, 299, 301                                 | formulation considerations, 484–485               | Human DNase I, recombinant   |
| disadvantages, 300                                   | gelling agents for, 481–484                       | (Pulmozyme), 710–711, 711  |
| drug candidates, 302                                 | lubricating jelly, 487                            | Human growth hormone, recombinant,   |
| drug release rate from, 309, 309                     | nasal, 333–334                                    | (Protropin, Humatrope), 694–695,   |
| embedded in matrix system, 304–305                   | ophthalmic, 329, 331–333, 334t                    | 695  |
| examples, 307t                                       | packaging, 322                                    | Human immunodeficiency virus (HIV)   |
| ion exchange resins, 305–306                         | preparation, 481                                  | infection  |
| microencapsulated drug, 304, 304                     | principles, 478                                   | efavirenz for, 686   |
| multitablet system, 303                              | single-phase, 478                                 | enfuvirtide for, 694   |
| oral, 302–307, 303–304, 306                          | terminology, 479–480                              | Humectant, 152t  |
| osmotic pump, 306, 306–307                           | two-phase. See Magmas                             | Hybridoma technology, for monoclonal   |
| rationale for, 300                                   | vaginal, 336, 737                                 | antibodies, 676  |
|  | Gelatin, 483                                      |  |
| repeat-action tablets, 307                           | Gender  | Hydantoin, 758   |
| transdermal. See Transdermal drug                    |   | Hydration state, dissolution and, 177  |
| delivery systems                                     | clinical studies, 54                              | Hydrocortisone   |
| in vitro/in vivo correlations, 309–310               | dosage determination, 64, 66                      | in ear preparations, 630   |
|  | Gene therapy, 40                                  | phonophoresis of, 731  |
| F  | in biotechnology production                       | in suppositories, 367t   |
| •  | preparation, 678–679                              | Hydrolytic drug, 759   |
| Facilitated diffusion in, 171                        | General anesthetic, 196                           | Hydrophilic ointment, 317  |
| Factor VIII (ReFacto), recombinant,                  | Gliadel Wafer, 745, 745-746, 746                  | Hydrophilic petrolatum, 317  |
| 687, 687   | Globule, 765                                      | Hydrophilic-lipophilic balance (HLB)   |
| Feathering method, in                                | Glucocorticoid, 758                               | system, 740  |
| pharmacokinetics, 205                                | Glutamate inhibitor, 758                          | Hyperglycemic drug, 759  |
|  | Glycerogelatins, 325–326                          | rijpergrj cenne drug, ros  |
| Fecal softener, 758                                  | Gonadotropin, 758                                 |  |
| Federal Food, Drug, and Cosmetic Act                 | Goserelin (Zoladex), 746–747                      |  |
| of 1938  |   | T  |
| Comprehensive Drug Abuse                             | Granules, 765                                     | Ibritumomab tiuxetan (Zevalin), in   |
| Prevention and Control Act of                        | clinical case study, 236–237                      | non-Hodgkin lymphoma, 650,   |
| 1970, 17   | definition, 215                                   | <i>650,</i> 702  |
| Durham-Humphrey Amendment of                         | effervescent, 233c–234c                           | Imbibition, of gel, 479–480  |
| 1951, 16   | dry or fusion method, 234                         | Immunizing agent, 759  |
| Kefauver-Harris Amendments of                        | wet method, 234–235                               | Immunoglobulin, 759  |
| 1962, 16–17  | flow properties, 232                              | Immunoglobulin G   |
| Fentanyl   | particle size, 216                                | monoclonal antibody preparation  |
| in lollipops, 293                                    | pharmaceutics case study, 235–236                 | from, 676  |
| transdermal drug delivery systems,                   | preparation, 231, 231–232                         | placental transfer of, 580   |
| 347t-348t  | Granulocyte colony-stimulating factor             | Immunosuppressant, 759   |
| Fertility agent, 758                                 | monomethoxypolyethylene glycol                    | Implants, 744–748, 745, 746, 765   |
|  | (pegfilgrastim), 689, 689                         |  |
| First and a biration 100                             | recombinant (Filgrastim), 687–689,                | in brain tumor resection cavity, 745,  |
| First-order kinetics, 169                            | . 0   | 745–746, 746   |
| Flavorant, 152t                                      | 689   | goserelin in, 746–747  |
| Flavoring pharmaceuticals, 156–158                   | Granulocyte macrophage colony-                    | Indium-111 chloride, 649–651, 650  |
| Flexible collodion, 433, 437                         | stimulating factor, recombinant,                  | Information technology, in drug  |
| Foam, 765  | 689–690   | manufacture, 82–83, 83, 84   |
| Food   | Gray, as radiation unit, 640                      | Infusion, 765  |
| drug interactions with metabolism                    | Growth factor, recombinant, 694                   | Inhalant, 765  |
| and, 177   | Growth hormone, 758                               | Inhalation, 766  |
| oral administration timing, 256                      | human, 39, 675, 683t, 694–695                     | Inhalation administration route  |
| Food and Drug Administration                         | recombinant, 694                                  | aerosols   |
| Current Good Manufacturing                           | Guide to the Inspection of Bulk                   | examples, 499t   |
| Practice, 52, 68, 70, 76, 88, 142                    | Pharmaceutical Chemicals, 83                      | metered dose, 496-498  |
| drug classification system, 56                       | Gum, 765  | inhalants, 550, 628  |
| Modernization Act of 1997, 22–23, 85                 |   | powders, 227, 228  |
| pregnancy categories, 18                             | 11  | solutions, 628   |
| Forbes bottle method, 474–475                        | Н   | Initial product formulations, 51–52  |
| Furosemide, in nuclear medicine, 660                 | Haemophilus influenzae vaccines                   | Injections, 766  |
|  |   | ,  |
| Fusion inhibitors, recombinant, 693–694              | recombinant (HibTITER,                            | intravenous injections   |
| Fusion method  | PedvaxHIB, ActHIB), 709–710                       | advantages, 510  |
| for effervescent granule preparation,                | Half-life   | automated, 511   |
| 232–235  | of drugs  | disadvantages, 517   |
| ointment preparation, 321–322                        | vs. total body clearance, 207                     | flow rates for, 510  |
| Fusion protein, special handling,                    | of radioisotopes, 639–640                         | history, 509–510   |
| 743–744  | Hard-shell capsule, 765                           | infusion devices for, 566-567  |
|  | Heavy metal antagonist, 758                       | procedure, 511   |
| G  | Hematinic drug, 758                               | transfer of additives to, 522  |
| G  | Hemorheologic agent, 758                          | volumes of, 547  |
| GABA agonist, 758                                    | Heparin, thrombocytopenia induced by,             | procedure, 511   |
| Gastric acid secretion inhibitor, 758                | lepirudin for, 685                                | safe systems for, 750  |
|  | Hepatitis B vaccine (Engerix-B,                   | veins for, 510   |
| Gastroprokinetic drug, 758                           | Recombivax HB), 709                               | Inotropic agent, 759   |
| Gel(s), 765  | Hirudin, recombinant, 685                         | Inserts  |
| classification, 480–481, 480t                        |   |  |
| colloidal, 478–479                                   | Histamine H <sub>1</sub> receptor antagonist, 758 | ophthalmic, 737–739, 738   |
| definition, 478                                      | Histamine H, receptor antagonist, 758             | vaginal, 736, 737  |

| Insulin  | of prescription container, 95  | Medical devices, good manufacturing                            |
|--|--|--|
| autoinjection systems, 749, 749–750  | storage conditions on, 97  | practice standards for, 95                                     |
| crystalline vs. amorphous, 175–176   | Lacrisert system, 737–738  | Melting point, depression of, 108, 109c                        |
| Interferons, recombinant, 695–696, 743<br>Interleukins, recombinant, 696–699   | Lactinov granulas 22, 23   | Metabolism, drug, 47<br>active metabolites from, 168           |
| International Conference on  | Lactinex granules, 22–23<br>Lanolin, 318                                   | examples, 190t   |
| Harmonization of Technical   | Laxative, 759  | factors affecting, 200   |
| Requirements for Registration of   | Leaching, from plastic packaging, 92                                       | first-pass effect, 187   |
| Pharmaceuticals for Human Use  | Lead compound (prototype), 41–43   | first-pass effect in, 47                                       |
| (ICH), 73  | Length, units  | pathways, 200  |
| International Organization for   | intersystem conversion of, 773   | time course, 180   |
| Standardization, 15<br>Intrauterine devices, 735–736, 736, 766   | metric system, 770–773<br>Lepirudin (Refludan), 685                        | Metal complexing agent, 759<br>Methylcellulose, 483            |
| Intravenous injections   | Leukemia, acute myeloid, gemtuzumab  | Methylphenidate, transdermal drug                              |
| advantages, 510  | ozogamicin for, 683t   | delivery systems, 356  |
| automated, 511   | Leukotriene receptor antagonist, 759                                       | Metric system, 770–773   |
| disadvantages, 517   | Levigating agent, 152t   | Microemulsions, 475 Microstreaming, in phonophoresis, 731      |
| flow rates for, 510<br>history, 509–510  | Levigation, for particle size reduction,<br>223                            | Milk of magnesia, 486  |
| infusion devices for, 566–567  | Liniment, 766  | Mineralocorticoids, 759  |
| procedure, 511   | Lipase inhibitor, 759  | Minimum fill test, 322   |
| transfer of additives to, 522  | Liposomes, 739–742, 766  | Modified-release products                                      |
| veins for, 510   | advantages, 740  | case studies, 311–313  |
| volumes of, 547 Investigational New Drug Application   | amphotericin formulation with, 741 configurations for, 740                 | clinical consideration, 310–311<br>definition, 299, 301        |
| clinical protocol, 54–56   | daunorubicin formulation with, 741   | driving forces, 299  |
| clinical studies   | description, 739   | examples, 307t   |
| controls, 58-60  | disadvantages, 740   | labeling, 310  |
| designs, 58–59, 60   | doxorubicin formulation with, 742  | oral, 302–307  |
| drug dosage and terminology,<br>59–66, 60, 61, 62t, 63t, 65  | examples, 741<br>preparation, 739–740                                      | packaging, 311<br>storage, 311                                 |
| phases, 56–58, 57t, 58   | stealth, 740, 742  | terminology, 301   |
| content, 53  | Lipstick, 766  | uniformity of dosage units, 309                                |
| drug classification, 56  | Liquid(s), 766   | USP requirements, 309, 309-310                                 |
| FDA review, 56   | measurement techniques, 776–780  | in vitro/in vivo correlations, 309–31                          |
| meetings before, 56  | quantitative product strength, 774–775                                     | Molded tablet, 766   |
| for orphan drug, 67<br>termination, 67   | volume measurement, 778–780 Liquid preparations. <i>See also</i> Solutions | Monoamine oxidase inhibitor, 759<br>Monoclonal antibodies, 759 |
| treatment, 67  | advantages, 160  | in clinical practice, 699                                      |
| withdrawal, 67   | in capsules  | preparation, 702   |
| Ion exchange resins, 305-306, 759  | hard gelatin, 246  | radiolabeled, 645  |
| Iontophoresis, 344   | soft gelatin, 371  | toxic protein fusion with, 708                                 |
| Irrigation, 766  | dosage forms for, 103  | Mouthwash, 766 Mucoadhesive testosterone buccal                |
| Irritant, 759  | flavorants for, 160 incorporation into ointments, 319,                     | system, 733–734  |
| A CONTRACTOR OF THE CONTRACTOR | 319–321, 320, 321  | Muromonab-CD3 (Orthoclone OKT3),                               |
| J  | for nuclear medicine. See  | 703  |
| Jars, ointment, 326  | Radiopharmaceuticals   | Muscle relaxant, 759   |
| Jejunostomy feeding tube, 566  | oral administration, 103, 103t   | Mydriatic drug, 759  |
| Jelene, 481  | packaging, 24<br>Lotion, 766   | M  |
| Jelly(ies), 766  | Lozenge, 766   | N  |
| Iubricating, 487 Jet milling, for suspensions, 451   | Lung   | Narcotic antagonist, 759                                       |
| jet mining, for outperioretty, for   | drug administration to, 187t   | Narcotic drug, 759   |
| V  | aerosolized powders for, 227–228,  | Nasal administration route, 624–630                            |
| K  | drug excretion, 200  | for decongestants, 625   |
| Kefauver-Harris Amendments of 1962,  | inhalation solutions for, 625–628, 627                                     | drops for, 628–629<br>inhalants for, 628                       |
| 16–17  | Lung surfactant, 759   | inhalation solutions for, 625–628, 627                         |
| Keratolytic drug, 759  | Lymphoma, non-Hodgkin's  | ointments for, 333-334   |
| drug excretion, 200  | radiopharmaceuticals in  | preparations for   |
| radiopharmaceuticals for, 642, 642   | rituximab for, 689<br>Lyophilization, 766                                  | examples, 624–625, 625, 626t                                   |
| Kinematic viscosity, 615   | Lyophinzation, 700   | procedures, 629–630<br>sprays for, 628–629                     |
| Kineret, 698, 698  | М  | for systemic effects, 629–630                                  |
| Kinetic approach, to dosage regimen,   | IAI  | National Association of Boards of                              |
| 208<br>King's Guide to Parenteral Admixtures, 559  | Magmas   | Pharmacy, compounding  |
| King 5 Guiac to 1 archierai 11amixtures, 557   | bentonite, 485   | practices of, 87–88  |
| · ·  | definition, 478  | National Formulary, monographs, 11–12<br>86                    |
| L  | preparation, 481<br>principles, 478  | Natural colloids, 479  |
| Labeling   | Magnesium aluminum silicate, 483   | Needles, safe systems for, 750                                 |
| dietary supplement label, 96-97, 97  | Manufacture, liposomes, 739–742  | Nernst-Planck equation, in                                     |
| manufacturer's, 94–95  | Mathematical models, for   | iontophoresis, 728   |
| over-the-counter (OTC) drugs,  | compartmental analysis, in   | Neuraminidase inhibitor, 760                                   |
| 95–96, 96  | pharmacokinetics, 202  | Neuromuscular blocking agent, 760                              |

| New drug(s)   | microbial content, 322                                    | Parasympathomimetic drug, 760            |
|---|---|--|
| biologic characterization, 44–50, 46, 47                          | minimum fill test, 322                                    | Parenterals                              |
| clinical trial materials, 51-52                                   | nasal, 333–334  | administration routes                    |
| development, overview, 34-37, 35, 36                              | ophthalmic, 329, 331-333, 334t                            | intradermal, 195, 513                    |
| FDA-approved, standards in, 22                                    | packaging, 322  | intramuscular, 194, 511-512              |
| formulations for, 50–52   | polyethylene glycol in, 318                               | intravenous, 194–195                     |
| metabolism, 47  | preparation, 319, 319–322, 320, 321                       | specialized access, 513                  |
| pharmacology, 45–47, 46   | preservatives for, 318, 322, 324, 335                     | subcutaneous, 194                        |
| preformulation studies, 50–51. See also                           | rectal, 334–335, 335, 336t                                | subcutaneous route, 512–513              |
| Preformulation  | requirements, 322–323                                     | case studies, 573–575                    |
| toxicology, 47–50, 49   | storage, 322  | chemotherapeutic, handling and           |
| New drug application  | vaginal, 336–337, 336t                                    | disposal, 571                            |
| abbreviated, 72   | white, 317  | dialysis solutions, 571–572              |
| action letters, 70  | yellow, 317   | drug adsorption, 526                     |
| for animals, 72–73  | Omnibus Budget Reconciliation Act of                      | frozen, 535                              |
| annual reports on, 71–72  | 1990 (OBRA 90), 30  | implants, 513                            |
| Biologics License Applications, 72                                | One-compartment model, for                                | irrigation solutions, 571                |
| drug product labeling, 69–70                                      | pharmacokinetics, 207                                     | labeling, 535–542                        |
| FDA review, 70  | Open-model system, 202                                    | large-volume                             |
| for medical devices, 73   | Ophthalmic preparations                                   | administration, 543                      |
| NDA submission, 68–69   | buffering, 612–615, 615t                                  | caloric requirements, 562–563            |
| phase 4 studies, 70–71<br>postmarketing activities, 71            | case studies, 633–634                                     | electrolyte requirements, 561–562        |
|   | clarity, 617  | examples, 572t<br>incompatibilities, 559 |
| supplemental, 72<br>Nicotine, transdermal drug delivery           | for contact lens, 623<br>with contact lens wear, 620, 623 | intravenous infusion devices for,        |
| systems, 347t, 354  | contact lens-related, 623                                 | 566–567                                  |
| Nitroglycerin   | drug delivery from, 606–607                               | for maintenance therapy, 560–561         |
| administration route, 187   | gels, 329, 331–333, 334t                                  | manufacturing, 532                       |
| sublingual, 187   | inserts, 737–739, 738                                     | for nutrition, 563–565                   |
| transdermal drug delivery systems,                                | isotonicity, 609–612, 612t                                | packaging, 535–542                       |
| 345, 347t, 352–353  | novel, 737–739, 738                                       | for replacement therapy, 561             |
| Noncompliance   | ointments, 329, 331–333, 334t                             | transfer of additives to, 522            |
| with manufacturing regulations, 85                                | packaging, 617, 617                                       | water requirements, 561                  |
| patient, factors associated with, 94                              | pH, solubility and, 613c–614c                             | long-acting, 514, 739, 739t              |
| Nonsteroidal anti-inflammatory drug,                              | retention time, 607                                       | look-alike products, 569                 |
| 760   | sterility, 608, 608–609                                   | manufacturing                            |
| Novel dosage forms and drug delivery                              | thickening agents for, 615-616                            | antibiotics, 532                         |
| systems, 726–751  | viscosity, 615–616  | clarity in, 537                          |
| autoinjection systems, 749, 749-750                               | Opioid, 760   | contaminant exclusion in, 537            |
| benefits, 727   | Opioid antagonist, 760                                    | drug colligative properties and,         |
| drug composition for, 727   | Oprelvekin (Neumega), 698-699, 699                        | 517c-521c                                |
| fusion protein, 743-744   | Oral administration route                                 | dry, 524                                 |
| implants, 744–748, 745, 746                                       | absorption  | environment for, 542–543                 |
| iontophoresis, 728-729, 731, 732, 732t,                           | capsules for, 188t  | for extemporaneous compounding,          |
| 733t  | solutions for, 188t                                       | 555                                      |
| liposomes, 739–742  | suspensions for, 188t                                     | liquid with powder, 532                  |
| mechanisms, 727   | tablets for, 188t   | packaging in, 535–542                    |
| microspheres, 748   | osmotic pump formulations for,                            | quality assurance, 543–544               |
| mucoadhesive system, 733–734                                      | 734–735   | suspensions, 532                         |
| ophthalmic, 737–739, 738  | Organogels, 481   | novel drug delivery system, 739–742,     |
| oral, 733–735   | Orphan drugs, Investigational New                         | 739t                                     |
| osmotic pump, 734, 734–735  | Drug Application withdrawal, 67                           | official types, 513–514                  |
| parenteral, 739–742, 739t   | Osmotic pump drug delivery system,                        | pellets, 566                             |
| pegylated dosage forms, 742–743                                   | 734–735   | pyrogen removal and testing, 531         |
| phonophoresis, 729–731  | Otic administration route, 630–632                        | small-volume                             |
| safe-needle systems, 750  | procedures, 632   | advantages, 547                          |
| subgingival, 748, 749<br>topical, 727–733, 730, 731, 732t–733t    | solutions for, 630–632, 631t<br>analgesic, 630–632        | disadvantages, 547                       |
| * .   | anti-infective, 630–632                                   | examples, 545t–546t<br>insulin, 548–558  |
| vaginal, 735–737<br>Novobiocin, crystalline <i>vs.</i> amorphous, | anti-inflammatory, 630–632                                | manufacturing, 550                       |
| 175   | cerumen-removing, 630                                     | precautions with, 547–548                |
| Noyes-Whitney equation, 173                                       | suspensions for, 632                                      | solvents for, 514–516                    |
| Nuclear medicine. See   | Otic suspension, 767                                      | special considerations, 567–571          |
| Radiopharmaceuticals  | Ovulation stimulator, 760                                 | sterilization, 522–526                   |
| Numby Stuff iontophoresis system, 729                             | Oxytocic drug, 760  | storage, 535–542                         |
| rtaine y etain terrespresses system, 725                          | ony toda arag, 700  | substances added to, 521–522             |
| •   | D.  | vehicles for                             |
| 0   | Р   | aqueous, 514                             |
| Ocular administration route, 196                                  | Packaging   | nonaqueous, 516–517, 517c–521c           |
| Ointments, 152t, 766  | child-resistant, 93–94, 94                                | volume overages, 541                     |
| bases for, 317–318  | compliance, 94  | Particle size, of powders, 216–217, 216t |
| definition, 316   | containers for, 88–93, 89, 90, 91t                        | Partition coefficient, in dosage form    |
| dermatologic, 316, 327-329, 328, 330t,                            | glass, 91, 91t  | design, 121, 121c-123c                   |
| 331t, 332t  | light-resistant, 90                                       | Passive diffusion, 168–171               |
| hydrophilic, 317  | strip, 90, 90   | Pastes, 324–325, 767                     |
| labeling, 322   | Parasympatholytic drug, 760                               | Pastille, 767                            |

| Patches  | Plastibase, 480   | biostability, 43                                     |
|--|---|--|
| for transdermal drug delivery,                         | Plasticizer, 152t   | prolonged release, 43                                |
| 357. See also Transdermal drug                         | Plastics, in packaging, 91  | solubility, 43                                       |
| delivery systems (TDDSs)                               | Plateau concentration, of drug, 203                                 | Progesterone   |
| Pathologic state, dosage and, 64                       | Platelet growth factor, 760   | in vaginal insert, 735–736, 736                      |
| Pediatric patients                                     | Platelet-derived growth factor,                                     | in vaginal suppositories, 374, 387                   |
| in clinical studies, 55                                | recombinant, 694  | Progestin, 760                                       |
| drug dosage for, 60, 62–63                             | Platelet-reducing agent, 760  | Prostaglandin synthetase inhibitor, 760              |
| packaging resistant to, 93–94, 94                      | Pledget, 767  | Protease inhibitor, 760                              |
| suppositories for, 386                                 | Polymorphism design form design 112                                 | Proteasome inhibitor, 760                            |
| suspensions for, 456<br>total nutrient admixtures, 559 | Polymorphism, dosage form design, 112                               | Protectant, 760                                      |
| Pediculicide, 760                                      | Posterior pituitary hormone, 760<br>Potassium-sparing diuretic, 760 | Prothrombogenic drug, 760 Proton pump inhibitor, 761 |
| Pegylated dosage forms, 742–743                        | Potentiator, 760  | Psychedelic drug, 761                                |
| Pellet, 767  | Poultice, 767   | Psychotherapeutic drug, 761                          |
| Penicillin adjuvant drug, 760                          | Povidone, 483   | Pump, osmotic, 734–735                               |
| Penicillin G, crystalline vs. amorphous,               | Powders, 767  | 1,   |
| 175  | angle of repose, 218c-219c  | Q  |
| Pennkinetic system, sustained-release                  | blending, 223–225, 224–225  | G.   |
| suspensions, 457                                       | blowers for, 228, 228   | $Q_{10}$ method of shelf life estimation, 137        |
| Percutaneous absorption. See                           | bulkiness, 219c   | Quality assurance                                    |
| Transdermal drug delivery                              | clinical case study, 236–237  | definition, 78                                       |
| systems (TDDSs)  | comminution of, 222, 223  | for parenterals, pharmacy-prepared,                  |
| Pessaries. See Suppositories, vaginal                  | definition, 215   | 543–544  |
| Petrolatum, 317  | density, 220c   | for radiopharmaceuticals, 663–664                    |
| pH, in passive diffusion, 170                          | flow properties, 218c   | Quality audit, definition, 78                        |
| Pharmaceutical alternatives, definition,               | granule preparation from, 230–232,                                  | Quality control, definition, 78                      |
| 184  | 231   | Quality control unit, definition, 78                 |
| Pharmaceutical care, definition, 28–29                 | medicated   | Quantitative product strength, 774–775               |
| Pharmaceutical equivalents, definition,<br>184         | aerosol, 227, 228<br>bulk, 229–230                                  | Quarantined area, 78                                 |
| Pharmaceutical measurement                             | divided, 230  | _  |
| errors in  | particle size and, 216–217, 216t                                    | R  |
| significance, 770                                      | pharmaceutics case study, 235–236                                   | Dahina ya asimoa E90t                                |
| volume, 779  | porosity, 219c  | Rabies vaccines, 589t                                |
| weight, 776  | volume, 219c  | Racial immunity, 579 Radiation                       |
| systems for  | Preformulation studies  | exposure, drug antidote for, 658,                    |
| apothecary, 770, 772, 773, 777, 778                    | complex organic molecules, 131,                                     | 658–659  |
| avoirdupois, 770, 773                                  | 131c–133c   | ionizing, in parenteral sterilization,               |
| conversion among, 770                                  | dissolution, 50, 117-118, 118c-120c                                 | 526  |
| dosage units, 774, 775                                 | heat of vaporization, 107, 107c-108c                                | Radiogardase, for radiation exposure,                |
| formula reduction and                                  | hydrates and solvates, 124-125, 125c                                | 658, 658   |
| enlargement, 775                                       | melting point depression, 108, 109c                                 | Radiographic agent. See X-ray contrast               |
| household, 775–776                                     | membrane permeability, 121  | medium   |
| metric, 770, 771, 773                                  | microscopic examination, 107  | Radioimmunotherapy, 643                              |
| quantitative product strength,                         | organic ester considerations, 128,                                  | Radionuclides, definition, 639                       |
| 774–775  | 128c-129c   | Radiopaque agent, 761                                |
| techniques   | organic salt considerations, 126–127,<br>127c                       | Radiopharmaceutical drug, 761                        |
| volume, 778–780  |   | Radiopharmaceuticals, 638–670, 767                   |
| weight, 777<br>Pharmacist, contemporary role, 26–30    | particle size, 112<br>partition coefficient, 50, 121,               | applications, 638–639                                |
| Pharmacogenetics, 63–64                                | 121c–123c   | background information, 639–642                      |
| Pharmacokinetics                                       | pH, 161   | case studies, 668–669                                |
| definition, 167  | phase rule, 108, 110c–111c  | consultation on, 666, 666t, 667t                     |
| principles, 167  | physical description, 106–107                                       | definition, 638<br>dispensing, 664–665               |
| Pharmacopeias, 8–15                                    | physical form, 50–51  | distribution, 665                                    |
| Pharmacy   | pKa/dissociation constants, 123,                                    | dosage units for, 640                                |
| early research, 7–8                                    | 123c-124c   | drug interference with, 666t, 667t                   |
| heritage of, 4–8                                       | polymorphism, 112   | education on, 666                                    |
| mission, 28  | potency-designated active   | examples, 645, 646t                                  |
| practice standards, 29–30                              | pharmaceutical ingredients,   | gallium-67 citrate, 649                              |
| school of  | 129–130, 130c–131c  | harmful exposure to, antidote for,                   |
| first, 8   | solubility, 50, 112   | 658–659  |
| scientific viewpoint introduced, 6–7                   | stability, 51   | holmium-166, 653                                     |
| Phase 4 clinical studies, 70–71                        | Pregnancy, 18   | indium-111 chloride, 649–651, 650                    |
| Phase rule, in dosage form design, 108,                | Prescription Drug Marketing Act of                                  | labeling, 662  |
| 110c-111c  | 1987, 21<br>Preservatives   | lutetium-177, 653                                    |
| Phenothiazine, 760<br>Phonophoresis, 729–731           | general considerations, 162   | monoclonal antibodies, 643                           |
| Photosensitizer, 760                                   | mode of action, 162   | nonradioactive drugs used with,                      |
| Pigmenting agent, 760                                  | selection, 161–162  | 659–660  |
| Pilocarpine, insert, 738, 738–739                      | with sterilization, 161   | outcome monitoring, 667–668                          |
| pK <sub>2</sub> (dissociation constants), in dosage    | Process validation, definition, 78                                  | physics, 639–642<br>in positron emission tomography, |
| form design, 123, 123c–124c                            | Procrit (epoetin alfa), 690–691                                     | 653–658, 657t  |
| Plasma volume expander, 760                            | Prodrugs  | practice guidelines, 660–668                         |
| Plasters, 325, 767                                     | absorption, 43  | preparation, 662–663, 663                            |
|  |   | A A  |

| procurement, 661–662                      | Reprocessing, definition, 78             | Solubility                               |
|---|--|--|
| production, 641                           | Reproduction studies, 49                 | particle size, 113                       |
| quality assurance, 663–664                | Research, in early times, 7–8            | pH, 113, 115c–117c                       |
| regulatory process, 668                   | Reserpine, sources, 38                   | Solutions, 767. See also Solubility;     |
| rhenium-186/188, 653                      | Residency programs, for pharmacists,     | Solvents                                 |
| rhenium-HEDP, 653                         | 27                                       | colligative properties, 610              |
| safety, 665–666                           | Resin, electrolyte removing. See Ion     | definitions, 396                         |
|   |  |  |
| samarium-153, 652–653                     | exchange resins                          | elixirs, 420–424, 422t, 424              |
| sodium iodide (I-123), 651                | Respiratory stimulant, 761               | inhalation, 628                          |
| sodium iodide (I-131), 651–652            | Respiratory syncytial virus infection,   | irrigation, 571, 572t                    |
| storage, 661–662                          | palivizumab for, 704                     | ophthalmic                               |
| strontium-89 chloride, 647                | Retention enemas, 432                    | administration, 617-618, 618t-619t       |
| targeting strategies, 639, 643            | Retention time, ophthalmic               | for contact lens use, 621–622            |
| technetium-99m, 645-647                   | preparations, 607                        | drug delivery from, 606-607              |
| thallous-201 chloride, 648, 648-649       | Reteplase (Retavase), recombinant, 707   | examples, 618t-619t                      |
| vs. traditional drugs, 639                | Reticuloendothelial system,              | packaging, 617, 617                      |
| uses                                      | radiopharmaceuticals in, 646–647         | pharmaceutical requirements, 608,        |
|   |  |  |
| diagnostic, 642, 642–643                  | Retinitis, CMV, fomivirsen for, 685      | 608–617, 611t, 612t, 613c–614c,          |
| therapeutic, 643–645                      | Retinoic acid derivative, 761            | 615t                                     |
| yttrium-90, 647–648, 648                  | Returned drug products, regulations, 82  | types, 607                               |
| Random screening, in drug                 | Reverse iontophoresis, 729               | oral                                     |
| discovery, 40                             | Reverse osmosis, 404                     | colonic lavage, 410                      |
| Ranitidine (Zantac), mechanism-based      | Review clock, for new drug approval, 70  | examples, 427t–428t                      |
| design, 41                                | Rhenium-186/188, 653                     | magnesium citrate, 410–411               |
| Rapidly dissolving tablets, 267–268, 268  | Rhenium-HEDP, 653                        | rehydration, 409–410                     |
| Rasburicase (Elitek), 710                 | Rheology, 452                            | sodium citrate with citric acid, 411     |
|   |  |  |
| Rate reactions, 135, 135c–137c            | Rheumatoid arthritis                     | syrups, 411–412                          |
| Rate-controlled delivery. See also        | adalimumab for, 700                      | otic, 630                                |
| Modified-release products                 | anakinra for, 698                        | parenterals, 514                         |
| definition, 301                           | Rigid gas permeable contact lenses, 620  | preparation, 438–441                     |
| Ratio strength, 774, 775                  | Ringer's injection, USP, 558t            | proper administration and use, 425       |
| Rauwolfia alkaloid, 761                   | Ringer's irrigation, USP, 572t           | rectal, 431–432                          |
| 82Rb, in positron emission tomography,    | Rings, vaginal, 735, 736                 | sorption tendency, 93                    |
| 656                                       | Rinse, 767                               | strength, 434                            |
| Reaction order, 135                       | Rinsing solutions, for contact           | syrups, 411–420                          |
| Reaction rate, 135                        | lenses, 622                              | topical, 425–431                         |
|   |  |  |
| Recall, of drug products, regulations, 24 | Rituximab, ibritumomab tiuxetan          | vaginal, 431–432                         |
| Receptor sites, for drugs, 45, 46, 198    | (Zevalin) in non-Hodgkin                 | Solvents, 153t                           |
| Recombinant DNA technology                | lymphoma, 689                            | for extraction, 401–404                  |
| for biotechnology products, 674–676       | RNA, antisense, 679                      | glycerin, 402                            |
| in drug preparation, 39                   | Robiquet, Pierre, 7                      | isopropyl rubbing alcohol, 402           |
| Reconstitution                            | Robotic devices, 83, 83                  | "like dissolves like," 399               |
| of dry forms, 138                         | Roller compaction, for granule           | for parenteral, 403                      |
| of powders, for suspensions, 464          | preparation, 279–280                     | propylene glycol, 402                    |
| Rectal administration route, 188t,        | Rotary die process, for capsule          | selection, 401                           |
| 192–193                                   |  | in tinctures, 424–425                    |
|   | preparation, 250, 253, 254               |  |
| aerosols for, 335, 500–501, 501           | Rotary tablet press, 281                 | water, 404                               |
| creams for, 334–335, 336t                 | Rotavirus vaccine, 594t                  | Somatostatin analog, 761                 |
| ointments for, 334–335, 335, 336t         | Rubbing alcohol, 402                     | Sonophoresis, 344                        |
| solutions for, 431                        | Rubefacient, 761                         | Sorption, in packaging, 93               |
| suppositories, 386                        | Rubella vaccine, 589t                    | Space sprays, 490                        |
| drug absorption from, 369–371             |  | Spatulation, in powder blending,         |
| examples, 367t, 386                       | C  | 223–225, 224–225                         |
| local action, 368-369                     | <b>S</b>                                 | Specialized transport mechanisms,        |
| systemic action, 369                      | Salicylic acid collodion, 437-438        | 171–172                                  |
| suspensions for, 464–465                  |  | Spirits, 767                             |
|   | Salts                                    |  |
| Rectal fluids, buffering capacity, 370    | dissolution, 177                         | Sponge, 768                              |
| Reduction, 199                            | substitute, 761                          | Steady state concentration, of drug, 209 |
| Refrigeration, of hydrolysis prone        | Salvaged drug products, regulations, 82  | Stealth liposomes, 740, 742              |
| drugs, 138                                | Scabicide, 761                           | Sterility, 768                           |
| Registered pharmacist (RPh), 26           | Sclerosing agent, 761                    | Sterilization, preservatives with, 161   |
| Regulation, of drugs. See Drug            | Scopolamine, transdermal drug delivery   | Stick, 768                               |
| regulation and control                    | systems, 348t                            | Stiffening agent, 153t                   |
| Rehydration solutions, oral, 410          | Sedative, 761                            | Stimulant, 761                           |
| Remicade, 702–703                         |  | Stool softener, 761                      |
| Renal cell carcinoma, aldesleukin for,    | Selective estrogen receptor modulator    | Storage, labeling information for, 97    |
|   | (SERM), 761                              |  |
| 683t                                      | Sepsis, drotrecogin alfa (activated) in, | Strength, definition, 78                 |
| Renal disease, erythropoietin in, 683t    | 692                                      | Striant mucoadhesive testosterone        |
| Renal transplantation                     | Serotonin agonist, 761                   | buccal system, 733–734                   |
| basiliximab in, 701                       | Serotonin antagonist, 761                | Strip packaging, 90, 90, 768             |
| daclizumab in, 701                        | Simethicone emulsion, 477                | Strontium-89 chloride, 647               |
| muromonab-CD3 in, 703                     | Smoking deterrent, 761                   | Sulfonylurea, 761                        |
| Repeat-action tablets, 307                | Sodium alginate, 484                     | Sunscreening agent, 761                  |
| Replacement therapy, parenterals, 561     |  | Suppositories                            |
|   | Sodium iodide (I-123), 651               |  |
| Repository injections, 193                | Sodium iodide (I-131), 651–652           | appearance, 365, 366                     |
| Representative sample, definition, 78     | Soft gel capsule, 767                    | base, 153t                               |

| Suppositories (continued)               | otic, 446                              | effervescent, 266                        |
|---|--|--|
| bases for                               | packaging, 459                         | enteric-coated, 288                      |
|   |  | extended-release, 269                    |
| bioavailability and, 371                | parenteral, 451                        |  |
| classification, 372                     | powders for, 464–465                   | fillers for, 276                         |
| determination of amount required,       | preparation, 456–458                   | film-coated, 265                         |
| 378–382                                 | rectal, 463–464                        | flavorants for, 287                      |
| case studies, 391–392                   | sedimentation rate in, 448–450,        | fluid-bed coating of, 288, 288–289       |
| cathartic, 367t, 386                    | 448c–449c                              | glidant, 155t                            |
| definition, 365                         | storage, 459                           | glidants for, 234                        |
| density (dose replacement)              | sustained-release, 457–458             | glossants for, 287                       |
| calculations for, 379c–382c             | viscosity, 457                         | hardness, 271–272, 272–273               |
| glycerin, 368, 374, 386                 | Sweeteners                             | hypodermic, 266                          |
| hemorrhoidal, 369, 386                  | artificial, 158                        | immediate-release, 267                   |
| insertion                               |  |  |
|   | comparison of, 153t                    | imprinting of, 286, 286                  |
| rectal, 365                             | Swelling, of gel, 479                  | inspection of, 271, 271                  |
| vaginal, 367t, 388                      | Syneresis, of gel, 479                 | lubricants for, 155t, 264                |
| local action, 368–369                   | Syrups, 768                            | manufacture                              |
| packaging, 384                          | agitation, 418                         | all-in-one granulation methods for,      |
| pediatric, 386                          | colorants for, 418                     | 279, 279–280                             |
| preparation                             | components of, 412-415                 | changes in, 290                          |
| dose calculation for, 379c-382c         | vs. elixirs, 411                       | coated, 284–290                          |
| by hand rolling and shaping, 383        | flavorants for, 418                    | direct compression method for,           |
| by molding, 377–379                     | heating of, 418–419                    | 281–282, 282                             |
|   |  |  |
| rectal, 386                             | medicated, 412t, 419                   | dry granulation method for, 279–280      |
| drug absorption from, 369–371           | nonmedicated, 412t                     | quality standards for, 270, 270          |
| examples, 367t, 385                     | percolation, 419–420                   | molded (triturated), 266, 283, 283–284   |
| local action, 368–369                   | preparation, 418–420                   | official, examples, 290, 291t            |
| systemic action, 369                    | preservatives for, 415                 | opaquants for, 287                       |
| requirements, 383                       | proper administration and use, 425     | oral administration, 293                 |
| storage, 384                            | sorbitol-base, 415                     | packaging, 292, 293                      |
| systemic action, 369                    | sucrose-based, 415                     | polishing agent, 155t, 286, 286          |
| urethral, 386–387                       | Systemically acting drug, 761          | rapidly-dissolving (instant-release),    |
|   | bysterifically acting arag, 701        | 267–268, 268                             |
| appearance, 365                         | _                                      |  |
| bases for, 374                          | I                                      | repeat action, 281, 285                  |
| local action, 368–369                   |  | scored, 263, 270, 270                    |
| vaginal                                 | Tablets, 768                           | sealing coats for, 285                   |
| appearance, 386                         | antiadherents, 153t                    | splitting defects, 282, 282              |
| bases for, 365, 374                     | aromas for, 287                        | storage, 292, 293                        |
| examples, 367t, 387                     | binders for, 154t, 264                 | subcoating, 285, 285                     |
| local action, 368–369                   | buccal (sublingual), 265–266           | sugar-coated, 284–285, 284–285           |
| progesterone, 374, 387                  | capping defect, 281, 282               | sweeteners for, 287                      |
| systemic action, 369                    | **                                     | thickness, 271, 272                      |
|   | capsule diluent, 154t                  | vaginal, 269–270                         |
| Suppressant, 761                        | capsule opaquant, 155t                 |  |
| Surface area, of drug, dissolution and, | case studies, 294–297                  | waterproofing, 285                       |
| 174–175                                 | chewable, 266                          | weight, 271, 271                         |
| Surface sprays, 490                     | coating agent, 154t                    | Tamper-evident packaging, 81             |
| Surfactant, 153t                        | colorants for, 287                     | Technetium-99m, 642, 645–647             |
| Suspending agent, 153t                  | commercially available, 290, 291t      | Teratogens, 19t                          |
| Suspensions, 768                        | compressed                             | Terminal sterilization, 768              |
| administration, 447                     | all-in-one granulation methods for,    | Testosterone                             |
| advantages, 446                         | 279, 279–280                           | mucoadhesive buccal system, 733-734      |
| antacid, 459, 462–463                   |  | transdermal drug delivery systems,       |
| antibacterial, 463                      | dedusting of, 282, 283                 | 355–356                                  |
|   | direct compression method for,         |  |
| case studies, 502–505                   | 281–282, 282                           | Thallium, radioactive, exposure to,      |
| definition, 446                         | dry granulation method for,            | antidote for, 658                        |
| desirable features, 455                 | 279–280, 280                           | Therapeutic dose, 62, 62t                |
| dispersed phase, physical features,     | effervescent, 266                      | Therapeutic equivalents, 184             |
| 450–451                                 | enteric-coated, 265                    | nicotine, 354                            |
| dispersion medium for, 451–452,         | film-coated, 265                       | nitroglycerin, 345, 349, 350, 352–353    |
| 452c-456c                               | gelatin-coated, 265                    | scopolamine, 352                         |
| drug interactions with, 495             | manufacture, 276                       | testosterone, 355–356                    |
| extemporaneous compounding,             | multiple, 264–265, 265                 | Therapeutic index, of drugs, 62, 62t     |
| 458–459                                 |  | Therapeutic window, of drugs, 62         |
|   | punch method for, 280–281, 281         |  |
| flocculated, 451                        | quality standards for, 270, 270        | TheraSphere, 647, 648                    |
| injectable, 458                         | rapidly-dissolving, 267–268, 268       | Thixotropy, 479                          |
| ophthalmic                              | shapes, 270, 270                       | Thrombocytopenia                         |
| administration, 617–618, 618t–619t      | sugar-coated, 265                      | chemotherapy-related, oprelvekin         |
| for contact lens use, 621–622           | Wet granulation method for,            | for, 698                                 |
| drug delivery from, 606-607             | 276–279, 277–278                       | heparin-induced, lepirudin for, 685      |
| examples, 618t–619t                     | compression-coated, 289–290, 290       | Thrombolytic enzyme, 761                 |
| packaging, 617, 617                     | content uniformity, 271                | Thyroid hormone, 761                     |
| pharmaceutical requirements, 608,       | dedusting of, 282, 283                 | Tinctures, 768                           |
| 608–617, 611t, 612t, 613c–614c, 615t    |  |  |
|   | direct compression excipient, 154t     | benzoin, 425, 432–433<br>definition, 424 |
| types, 607                              | disintegrant, 154t                     | definition, 424                          |
| oral                                    | disintegrants for, 264                 | iodine, 432                              |
| examples, 459, 462–463                  | dispensing, 266                        | thimerosal, 433                          |
| powders for, 464–465                    | dissolution test for, 274-275, 275-276 | topical, 432–433                         |

| Tolerance, drug, 66                                 | Uricosuric drug, 762   | and capsule counting and filling                              |
|---|--|---|
| Topical preparations                                | Urinary acidifier, 762   | apparatus, 258  |
| iontophoresis for, 728–729, 731, 732,               | Uterine contractant, 762   | Vesicant, 762   |
| 732t, 733t  | Uterine contraction inhibitor, 762   | Vinblastine, sources, 38                                      |
| phonophoresis for, 729–731 solutions                |  | Vinca alkaloid, 762   |
| aluminum acetate, 428                               | V  | Vincaleukoblastine, discovery, 8<br>Vincristine, sources, 38  |
| aluminum subacetate, 428                            | 77 . 1   | Viral vaccines, 583–585                                       |
| calcium hydroxide, 397                              | Vaccines and vaccination   | Viscosity   |
| chlorhexidine gluconate, 430                        | active, 579–580<br>administration, nasal, 598                                | definition, 615   |
| coal tar, 429                                       | attenuated, 580, 583   | measurement, in rheology, 452                                 |
| examples, 406                                       | bacterial, 583, 589t–591t  | of ophthalmic preparations,                                   |
| hydrogen peroxide, 429–430                          | benefits, 578  | 615–616   |
| povidone-iodine, 430                                | bioterrorism and, 600-601  | in rheology, 452  |
| sprays, 426, 428                                    | cancer, 585–586  | of suspensions, 449   |
| thimerosal, 430–431<br>Topically acting drug, 761   | definition, 578  | temperature and, 615<br>Vitamin, 762                          |
| Toxicology  | handling, 581–583  | Volume  |
| acute or short-term toxicity studies, 48            | immunity and, 578–581  | measurement techniques, 778–780                               |
| carcinogenicity studies, 49                         | inactivated, 580, 583  | in quantitative product strength,                             |
| genotoxicity or mutagenicity studies,               | labeling, 581<br>mixed, 583, 586   | 774–775   |
| 49–50   | in pregnancy, 580  | units   |
| reproduction studies, 49                            | production, 581  | in apothecary system, 770, 772                                |
| subacute or subchronic studies, 48–49               | shipping, 581–583  | intersystem conversion of, 773                                |
| Tragacanth gum, 484                                 | side effects, 583, 585, 587  | metric system, 770–773  |
| Tranquilizer, 762 Transdermal drug delivery systems | storage, 581–583   | Volume of distribution, drug, 202                             |
| (TDDSs), 768–769                                    | strength, 583  |   |
| absorption in, 343–344                              | subunit, 583   | W   |
| advantages, 351–352                                 | suppositories for, appearance, 386   | Wafer, 769  |
| case studies, 358–360                               | toxoids, 580, 586, 589<br>viral, 583–585, 589t–591t                          | Warfarin  |
| clinical considerations, 356-357                    | Vaginal administration route, 192  | drug interactions, 488  |
| clonidine, 353–354                                  | aerosols for, 337, 500–501   | protein binding of, 204                                       |
| contraceptives, 355                                 | gels for, 737  | volume of distribution, 202                                   |
| definition, 342                                     | inserts for, 736, 737  | Water   |
| design features, 346–351, 347t–348t, 350, 351, 353  | novel preparations for, 735–737  | bacteriostatic, for injection, 515                            |
| disadvantages, 351–352                              | preparations for, 334t, 336, 336–337,  | distillation, 403   |
| enhancement, 344                                    | 336t   | hydrogen bonds in, 399  |
| estradiol, 347t, 354-355                            | rings for, 735, 736  | for injection, 515  |
| examples, 351, 352-356, 353                         | suppositories for, 367, 387, 387–388<br>bases for, 373                       | ion exchange purification, 403 purified, 402–403              |
| factors affecting, 343–344                          | local action, 368–369  | radiolabeled, in positron emission                            |
| fentanyl, 347t                                      | progesterone, 374  | tomography, 654   |
| historical aspects, 342–343                         | systemic action, 369   | requirements, in parenteral nutrition,                        |
| layers in, 346                                      | tablets for, 269-270   | 515   |
| membrane-controlled, 349, 351                       | Vaginal tablet/insert, 769   | reverse osmosis, 404  |
| methylphenidate, 356<br>monolithic, 346, 349        | Validation, 769  | sterile   |
| in vitro studies, 345, 345, 346                     | definition, 78   | for injection, 533  |
| in vivo studies, 344–345                            | Validation protocol, definition, 78  | in irrigation solutions, 571 viscosity, 615–616               |
| Transdermal matrix patch, 769                       | Valve assembly, of aerosols, 494<br>Van der Waals forces, in solubility, 397 | Water channels, 169   |
| Transdermal membrane patch, 769                     | Vapor pressure, of mixtures,   | Water-in-oil emulsions, 465                                   |
| Transderm-nitro transdermal                         | 517c-518c  | Water-miscible bases, for suppositories,                      |
| therapeutic system, 349, 349                        | Vaporizers, for inhalation therapy, 627,                                     | 373–374   |
| Transplantation                                     | 627  | Waterproofing, of tablets, 285                                |
| basiliximab in, 701<br>daclizumab in, 701           | Varicella virus vaccines, 589t   | Water-removable bases, 318                                    |
| muromonab-CD3, 703                                  | Vasoconstrictors, 762  | Water-removable bases, for ointment,                          |
| Transportation, on labeling, 97                     | for ophthalmic use, 607–608, 624   | 318<br>Water-soluble bases                                    |
| Trastuzumab (Herceptin), 705–706                    | with subcutaneous injections, 194<br>Vasodilators, 762                       | for ointments, 318  |
| Tricyclic antidepressant, 762                       | with subcutaneous injections, 194  | for suppositories, 373–374                                    |
| Trituration   | Vasopressor, 762   | Weak acids, solubility, 398, 399t                             |
| particle size reduction, 216–217, 216t              | Veegum, as gelling agent, 483  | Weak bases, solubility, 399t                                  |
| in powder blending, 223–225, 224–225                | Vegetable oils, as injection vehicles, 517                                   | Wecobee bases, 373  |
| Two-compartment model, for pharmacokinetics, 204    | Vegetable powders, definition, 216   | Weight  |
| Tyrosine kinase inhibitor, 708, 708–709             | Vehicles, 155t, 769. See also Water  | body, dosage determination and, 64                            |
| Tyroome mimoe minerior, 700, 700 707                | isotonic phosphate, for ophthalmic   | of capsules, variation of, 253–254                            |
|   | preparations, 615, 615t  | measurement techniques, 777 in quantitative product strength, |
| U   | for parenterals<br>aqueous, 514  | 774–775   |
| United States Pharmacopeia                          | nonaqueous, 516–517, 517c–521c   | units   |
| compounding monographs, 86–87                       | sterile, 155t  | in apothecary system, 777                                     |
| and National Formulary, 8–11                        | Ventriculography, radiopharmaceuticals                                       | in avoirdupois system, 773                                    |
| powder particle size, 216                           | in, 642–643  | intersystem conversion of, 773                                |
| United States Pharmacopeial                         | Verapamil, metabolism, 199   | metric system, 770–773  |
| Convention, 9–11                                    | Verified documentation, definition, 78                                       | vs. volume, in solutions, 404                                 |
| Urea cyclic disorder agent, 762                     | Versacount Model automatic tablet  | Well-closed container, 88                                     |

## **794** INDEX uploaded by [stormrg]

Wet granulation method, for tablet manufacture, 276–279, 277–278 Wet gum (English) method, for emulsion preparation, 473 Wet method, for effervescent granule preparation, 234–235 Wetting agents for emulsions, 466 HLB values, 470t for suspensions, 469 Wetting solutions, for contact lenses, 623 White ointment, 317 White petrolatum, 317 Witepsol bases, 373 Women in clinical studies, 54 dosage determination for, 64

World Health Organization, International Pharmacopeia, 12 Wurster process, for tablet coating, 289

## X

Xanthine alkaloid, 762 Xanthine oxidase inhibitor, 762 Xerogels, 479 X-ray contrast medium, 762 Xylitol, in chewable tablets, 283



Yellow fever vaccine, 589t Yellow ointment, 317 Yttrium-90, 647–648, 648

## Z

Zanamivir for inhalation, 229 Zero order rate reactions, 135c–137c Zinc gelatin, 325 Zinc oxide paste, 325 Zinc oxide–eugenol mixture, 434 Z-track injection technique, 512 Zydis system, for rapidly-dissolving tablets, 267