
INDEX

Numbers preceded by A refer to Appendix pages.

A

Accuracy, 14
Addition rule, 72–73
Analysis of variance, 306–308
 assumptions, 307
 completely randomized design, 308–334
 one-way, 309–334
 procedure, 307–308
 randomized complete block design, 334–346
 repeated measures design, 346–356
 two-way, 336–346
Arithmetic mean, 38
Average hazard rate, 760

B

Backward elimination, 564
Bayes's theorem, 68, 80–83
Bernoulli process, 99–101
 β_1 , confidence interval, 438
 hypothesis test, 432–434
Binomial distribution, 99–108
 parameters, 105–107
 table, A-3–A-31
 Use of table, 104–105
Biostatistics, 3
Birth rate, crude, 15–10
Bivariate normal distribution, 445
Bonferroni's method, 324, 327
Box-and-whisker plot, 50–52

C

Case-fatality ratio, 15–14
Cause-of-death ratio, 15–8
Censored data, 752
Central limit theorem, 139–140
Central tendency, measures, 38–43

Chi-square distribution, 195–197, 600–657
 mathematical properties, 601–604
 table, A-41
 use in goodness-of-fit tests, 604–619
 small expected frequencies, 604
 use in tests of homogeneity, 630–634
 small expected frequencies, 633
 use in tests of independence, 619–630
 small expected frequencies, 625
 2 × 2 table, 625–627

Class interval, 22

Coefficient of determination, 427–428

Coefficient of multiple determination,
 501–503

Coefficient of variation, 45–46

Combination, 101

Completely randomized design, 308–334
 ANOVA table, 317
 assumptions, 311

Compound symmetry, 348

Computers:

 and analysis of variance, 308, 321–323, 326–327,
 341–343, 350–351, 355
 and biostatistical analysis, 15–16
 and chi-square, 615, 623
 and descriptive statistics, 21, 22–30, 47
 and hypothesis testing, 232–233, 243–244,
 258–259
 and interval estimation, 169–170
 and logistic regression, 573
 and multiple correlation analysis, 512–519
 and multiple regression analysis, 494–496
 and random numbers, 16
 and simple linear regression and correlation
 analysis, 450–451
 and stepwise regression, 560–563

- Confidence coefficient, 169
 - Confidence interval:
 - for β_1 , 438
 - multiple regression, 506
 - for difference between two population means, 177–185
 - nonnormal populations, 179–180
 - for difference between two population proportions, 187–188
 - for mean of Y , given X , 441–442
 - for $\mu_{y|1 \dots k}$, 508–509
 - for population mean, 165–178
 - nonnormal populations, 168–171
 - for population proportion, 185–186
 - practical interpretation, 167
 - for predicted Y , 441–442, 508–509
 - probabilistic interpretation, 167
 - for ratio of two variances, 198–201
 - for ρ , 454
 - for variance, 194–198
 - Confusion matrix, 219
 - Contingency table, 619
 - Correction for continuity, 152
 - Correlation coefficient:
 - multiple, 510–513
 - simple, 446–450
 - Correlation model:
 - multiple, 510–519
 - simple, 445–446
 - Cox regression model, 768–772
 - hazard function, 768–769
 - Critical region, 224
 - Critical value, 224
 - Cumulative frequencies, 25
 - Cumulative relative frequencies, 25
- D**
- Data, 2
 - grouped, 22–37
 - raw, 20
 - sources, 3
 - Death rate:
 - crude, 15-3
 - fetal, 15-7
 - specific, 15-3
 - standardized, 15-3
 - Death rates and ratios, 15-3 through 15-10
 - Death ratio, fetal, 15-7
 - Decision rule, 218
 - Degrees of freedom, 45
 - Density function, 115
 - Descriptive statistics, 2, 19–64
 - Dispersion, measures, 43–49
 - Distribution-free procedures, 671
 - Dummy variable, 544–559
- E**
- Epidemiology, 779
 - Estimation, 161–210
 - in simple linear regression analysis, 434, 441
 - Estimator, 165
 - robust, 170
 - Events:
 - complementary, 74
 - independent, 73–74
 - mutually exclusive, 68
 - EXCEL:
 - and binomial distribution, 106
 - Exclusive or, 73
 - Experiments, 10
 - designing, 14–15
 - Exploratory data analysis, 52
 - Extrapolation, 442, 459–460
- F**
- Factorial, 101
 - Factorial experiment, 358–369
 - ANOVA table, 364
 - assumptions, 362
 - False negative, 79
 - False positive, 79
 - Family-wise error rates, 506
 - F distribution, 199
 - table of, A-42–A-51
 - Fecundity, 15-10
 - Fertility, 15-10
 - measures, 15-10 through 15-12
 - Fertility rate:
 - age-specific, 15-11
 - cumulative, 15-12
 - general, 15-10

- standardized, 15-12
 - total, 15-12
 - Finite population correction, 141
 - Fisher exact test, 636-640
 - table for, A-55-A-85
 - Fisher's z , 453-454
 - table, A-54
 - Fixed effects model, 311
 - F-max test, 198
 - Forward selection, 563
 - Frequency distribution, 22-37
 - Frequency polygon, 27
 - Friedman test, 712-716
 - table for, A-102-A-103
 - F test, 316-317
- G**
- Goodness-of-fit tests, 604-616
 - Grouped data, 22-37
- H**
- Histogram, 25-28
 - Hypothesis, 215
 - alternative, 216
 - formulating, 14
 - null, 216
 - research, 215
 - statistical, 216
 - Hypothesis tests, 215-303
 - by means of confidence interval, 225-226
 - difference between means, 236-249
 - nonnormal populations, 242-243
 - population variances known, 236-238
 - population variances unknown, 238-243
 - for β_i , multiple regression, 504-506
 - for β_1 , simple linear regression, 427-432
 - one-sided, 226-228
 - purpose, 215, 220
 - single population mean, 222-236
 - non-normal population, 230-232
 - population variance known, 222-228
 - population variance unknown, 228-230
 - single population proportion, 257-259
 - single population variance, 264-266
 - steps in, 216
 - two population proportions, 261-264
 - two population variances, 267-272
 - two-sided, 226
- I**
- Immaturity ratio, 15-14
 - Incidence rate, 15-13
 - Inclusive or, 73
 - Inferential statistics, 2, 162
 - Interaction, 359-360, 550
 - Interpolation, 442
 - Interquartile range, 48
 - Interval estimate, 165
 - Interval scale, 6
- J**
- Joint distribution, 445
- K**
- Kaplan-Meier procedure, 756-761
 - Kolmogorov-Smirnov test, 698-703
 - advantages and disadvantages, 703
 - and StatXact computer analysis, 702
 - table for, A-99
 - Kruskal-Wallis test, 704-709
 - table for, A-100-A-101
 - Kurtosis, 48-49
- L**
- Least squares, method, 420
 - Least-squares line, 420-422
 - Levene's test, 201, 270
 - Location parameters, 47
 - Log rank test, 763-765
 - Logistic regression, 569-581
 - Loss to followup, 751
- M**
- Mann-Whitney test, 690-696
 - table for, A-95-A-98
 - Mantel-Haenszel statistic, 650-653
 - Margin of error, 168
 - Mean, 38-40
 - properties, 40
 - Measurement, 6
 - Measurement scales, 5-6
 - Median, 40
 - properties, 41

Median test, 686–689

MINITAB:

- and binomial distribution, 107
- and box-and-whisker plots, 51–52
- and chi-square, 615–616, 623, 632
- and confidence intervals for a mean, 169–170
- and descriptive measures, 47
- and dummy variables, 546–547, 550, 555
- and factorial experiment, 367–368
- and frequency distributions, 27
- and Friedman test, 716
- and histograms, 26–27
- and hypothesis testing, 253, 258–259
- and Kruskal-Wallis test, 709
- and Mann-Whitney test, 694–696
- and median test, 689
- and multiple correlation, 512, 515, 517
- and multiple regression, 495, 508
- and normal distribution, 126–127
- and one-way ANOVA, 321–322
- and ordered array, 20–21
- and Poisson distribution, 111–112
- and repeated measures ANOVA, 349–350
- and sign test, 680
- and simple linear regression, 421, 444
- and Spearman rank correlation, 724
- and stem-and-leaf displays, 29–30
- and stepwise regression, 560–563
- and two-way ANOVA, 341–342
- and Wilcoxon test, 685

Mode, 41

Morbidity, 15-13

- measures, 15-13 through 15-14

Mortality rate:

- infant, 15-7
- maternal, 15-6
- neonatal, 15-7
- perinatal, 15-7

Multicollinearity, 542

Multiple comparison, 322–326

Multiple correlation coefficient,
510–513

Multiple correlation model, 510–513

Multiplication rule, 71–72

Multivariate distribution, 510

Multivariate normal distribution, 510

N

Nominal scale, 6

Nonparametric statistics, 671–747

Nonrejection region, 218

Normal distribution, 118–127

- applications, 122–127
- characteristics, 118–119
- standard, 118–122
- table, A-38–A-39

O

Observation, 14

Observational study, 642–643

Odds ratio, 645–648

Ogive, 96

Operating characteristic curve, 277

Ordered array, 20–21

Ordinal scale, 6

Outliers, 52

P

Paired comparisons, 249–254

Parameter, 38

Partial correlation, 513–519

Partial regression coefficients, 492

Percentile, 47

Point estimate, 163

Poisson distribution, 108–113

- table of, A-32–A-37

Poisson process, 109

Population, 5

- finite, 5
- infinite, 5
- sampled, 164
- target, 164

Power, 272–279

Precision, 14, 168

Predictive value negative, 80

Predictive value positive, 80

Prospective study, 642

Prediction interval

- multiple regression, 508–509
- simple linear regression, 441–442

Prevalence rate, 15-14

Probability, 65–85

- posterior, 68

- prior, 68
 - classical, 66–67
 - conditional, 70
 - joint, 71
 - marginal, 70, 75
 - objective, 66–67
 - personalistic, 67
 - properties, 68–69
 - relative frequency, 67
 - subjective, 67–68
- Probability distributions, 92–132
 - of continuous variables, 113–128
 - of discrete variables, 93–113
 - cumulative, 96–98
 - properties, 95
- Product-limit method, *see* Kaplan-Meier procedure
- Proportional hazards model, *see* Cox regression model
- Proportional mortality ratio, 15-8
- p values, 225

Q

- Qualitative variables, 4, 543–556
- Quartile, 47–48

R

- R
 - and box-and-whisker-plots, 51
 - and confidence interval between two means, 183
- Random digits, table, A-2
 - use, 9–10
- Randomized complete block design, 334–346
 - ANOVA table, 338
 - assumptions, 337
- Range, 43–44
- Rank transformation, 672
- Rate, 15-2
- Ratio, 15-2
- Ratio scale, 6
- Regression:
 - logistic, 569–581
 - multiple, 489–510
 - assumptions, 491
 - equation, 491–492
 - model, 490–492

- nonparametric, 727–730
- resistant line, 442–444
- simple linear, 413–446
 - assumptions, 415–416
 - equation, 417–423
 - model, 414–416
 - stepwise, 560–564
- Rejection region, 218
- Relative frequencies, 24–25
- Relative risk, 643–645
- Reliability, 14
- Reliability coefficient, 167
- Repeated measures design, 346–356
 - assumptions, 347–348
 - definition, 347
- Research study, 10
- Residual, 429
- Resistant line, 442–444
- Retrospective study, 643
- Risk factor, 642

S

- Sample, 5
 - convenience, 165
 - nonrandom, 164–165
 - random, 164–165
 - simple random, 7–10
 - size for controlling Type II errors, 277–279
 - size for estimating means, 189–191
 - size for estimating proportions, 191–193
 - stratified proportional to size, 13
 - stratified random, 12
 - stratified systematic, 12
 - systematic, 11
- Sampling distribution, 135–160
 - characteristics, 136
 - construction of, 135
 - definition, 135
 - of difference between sample means, 145–150
 - nonnormal populations, 148
 - of difference between sample proportions, 154–156
 - of sample mean, 136–145
 - nonnormal populations, 139–141
 - of sample proportion, 150–153

- SAS:
 and chi-square analysis, 623–625
 and descriptive measures, 47
 and factorial experiment, 367, 368
 and hypothesis testing, 233, 244–245
 and logistic regression, 572–576
 and multiple regression, 493, 496
 and one-way ANOVA, 322
 and repeated measures ANOVA, 350–351
 and simple linear regression and correlation, 442–443, 450–451
 and Tukey's HSD test, 326
 and two-way ANOVA, 341–342
- Scatter diagram, 419–420
- Scientific methods, 13–15
- Secondary attack rate, 15–14
- Sensitivity, 80
- Significance level, 218–219
- Sign test, 673–680
- Simple random sampling, 7–10
 without replacement, 7–8
 with replacement, 7–8
- Skewness, 41–42
- Slope, 415
- Spearman rank correlation coefficient, 718–724
 table for, A-104
- Specificity, 80
- Sphericity, 348
- SPSS:
 and Fisher exact test, 640
 and frequency distribution, 25–26
 and kurtosis, 49
 and logistic regression, 577
 and Mann-Whitney test, 695–696
 and Mantel-Haenzcel test, 652–653
 and multiple regression, 493
 and odds ratio, 648
 and partial correlation, 516, 518–519
 and repeated measures ANOVA, 350–351
 and skewness, 43
 and survival analysis, 665–666
 and Tukey's HSD test, 327
- Standard deviation, 45
- Standard error of mean, 139
- Standard normal distribution, 118–122
 table of, A-38–A-39
- Statistic, 38
- Statistical inference, 7, 162
- Statistics, 2
- Stem-and-leaf-display, 28–30
- Stepwise regression, 560–569
- Studentized range, 324
 table of, A-52–A-54
- Student's distribution, 172–177
 table of, A-40
- Sturges' rule, 22
- Survival analysis, 750–776
 censored survival times, 752
 types, 752–753
 Cox regression, hazard function, 768–772
 cumulative distribution function, 754
 Kaplan–Meier procedure, 756–761
 nonparametric technique, 756
 probability of surviving, 756–757
 probability distribution function, 755
 statistical distribution functions, 753
 survival curves, comparing, 763–766
 time-to-event data, 751–756
- T**
- t* distribution, 171–177
 and difference between means, 179–183
 population variances equal, 179–180
 population variances not equal, 180–183
 properties, 172
 table of, A-40
- Time-to-event data, *see* Survival analysis
- Test statistic, 217–218
- Trimmed mean, 170
- Tukey's HSD test, 323–324
- Tukey's line, 443–444
- Type I error, 219
- Type II error, 219, 272–279
- U**
- Unbiasedness, 163
- Uniform distribution, 614–616
- Unit of association, 459
- V**
- Variable, 3
 continuous random, 4

- dependent, 415
- discrete random, 4
- dummy, 544–556
- explanatory, 490
- extraneous, 307
- independent, 415
- predictor, 417, 490
- qualitative, 4, 543–556
- quantitative, 4
- random, 4
- response, 307, 417
- treatment, 307
- Variable selection
 - procedures, 560–564
- Variance, 44–45
 - interval estimation, 194–197
- Variance ratio, 316
- Variance ratio test, 198, 268–272
- Vital statistics, 778–796

- W**
- Weibull distribution, 755–756
- Wilcoxon test, 681–686
 - table for, A-86–A-95

- Y**
- Yates' correction, 627
 - y-intercept, 415