

## REFERENCES

---

1. Lipinski, C. A., Drug-like properties and the causes of poor solubility and poor permeability, *J. Pharmacol. Tox. Meth.* **44**, 235–249 (2000).
2. Martin, E. J.; Blaney, J. M.; Siani, M. A.; Spellmeyer, D. C.; Wong, A. K.; Moos, W. H., Measuring diversity: Experimental design of combinatorial libraries for drug discovery, *J. Med. Chem.* **38**, 1431–1436 (1995).
3. Drews, J., Drug discovery: a historical perspective, *Science* **287**, 1960–1963 (2000).
4. Pickering, L., Developing drugs to counter disease, *Drug Discov. Dev.* 44–47 (Feb 2001).
5. Dennis, C. et al., The human genome, *Nature* **409**, 813–958 (2001).
6. Handen, J. S., High throughput screening challenges for the future, *Drug. Disc. World* 47–50 (Summer 2002).
7. Testa, B.; Caldwell, J., The “ad hoc” approach as a compliment to ligand design, *Med. Chem. Rev.* **16**, 233–241 (1996).
8. Stenberg, P.; Luthman, K.; Artursson, P., Virtual screening of intestinal drug permeability, *J. Control. Rel.* **65**, 231–243 (2000).
9. Baum, R. M., Combinatorial approaches provide fresh leads for medicinal chemistry, *Chem. Eng. News* 20–26 (Feb. 7 1994).
10. Halliday, R. G.; Drasco, A. L.; Lumley, C. E.; Walker, S. R., The allocation of resources for R&D in the world’s leading pharmaceutical companies, *Res. Dev. Manage.* **27**, 63–77 (1997).
11. Gaviragli, G.; Barnaby, R. J.; Pellegatti, M., Pharmacokinetic challenges in lead optimization, in Testa, B.; van de Waterbeemd, H.; Folkers, G.; Guy, R. (eds.), *Pharmacokinetic Optimization in Drug Research*, Verlag Helvetica Chimica Acta, Zürich and Wiley-VCH, Weinheim, 2001, pp. 3–14.

---

*Absorption and Drug Development: Solubility, Permeability, and Charge State.* By Alex Avdeef  
ISBN 0-471-423653. Copyright © 2003 John Wiley & Sons, Inc.

12. Lipinski, C. A.; Lombardo, F.; Dominy, B. W.; Feeney, P. J., Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings, *Adv. Drug Deliv. Rev.* **23**, 3–25 (1997).
13. Pickering, L., ADME/Tox models can speed development, *Drug Discov. Dev.* 34–36 (Jan. 2001).
14. Kaplita, P. V.; Magolda R. L.; Homon, C. A., HTS: Employing high-throughput automation to better characterize early discovery candidates, *PharmaGenomics* 34–39 (2002).
15. Lipinski, C. A., Drug-like properties and the causes of poor solubility and poor permeability, *J. Pharmacol. Tox. Meth.* **44**, 235–249 (2000).
16. Lipinski, C. A., Avoiding investment in doomed drugs—is solubility an industry wide problem? *Curr. Drug Discov.* 17–19 (April 2001).
17. Taylor, P. J., Hydrophobic properties of drugs, in Hansch, C.; Sammes, P. G.; Taylor, J. B., (eds.), *Comprehensive Medicinal Chemistry*, Vol. 4, Pergamon, Oxford, 1990, pp. 241–294.
18. Fick, A., Ueber diffusion, *Ann. Phys.* **94**, 59–86 (1855).
19. Artursson, P., Application of physicochemical properties of molecules to predict intestinal permeability, AAPS Workshop on Permeability Definitions and Regulatory Standards, Arlington, VA, Aug. 17–19, 1998.
20. van de Waterbeemd, H. Intestinal permeability: Prediction from theory, in Dressman, J. B.; Lennernäs, H. (eds.), *Oral Drug Absorption—Prediction and Assessment*, Marcel Dekker, New York, 2000, pp. 31–49.
21. McFarland, J. W., On the parabolic relationship between drug potency and hydrophobicity, *J. Med. Chem.* **13**, 1192–1196 (1970).
22. Dearden, J. C.; Townsend, M. S., Digital computer simulation of the drug transport process, in *Proc. 2nd Symp. Chemical Structure-Biological Activity Relationships: Quantitative Approaches* (Suhl), Akademie-Verlag, Berlin, 1978, pp. 387–393.
23. Kubinyi, H., Lipophilicity and biological activity, *Arzneim.-Forsch./Drug Res.* **29**, 1067–1080 (1979).
24. Kubinyi, H., Strategies and recent technologies in drug discovery, *Pharmazie* **50**, 647–662 (1995).
25. Avdeef, A., Physicochemical profiling (solubility, permeability, and charge state), *Curr. Top. Med. Chem.* **1**, 277–351 (2001).
26. Avdeef, A., High-throughput measurements of solubility profiles, in Testa, B.; van de Waterbeemd, H.; Folkers, G.; Guy, R. (eds.), *Pharmacokinetic Optimization in Drug Research*, Verlag Helvetica Chimica Acta; Zürich and Wiley-VCH, Weinheim, 2001, pp. 305–326.
27. Avdeef, A., High-throughput measurements of permeability profiles, in van de Waterbeemd, H.; Lennernäs, H.; Artursson, P. (eds.), *Drug Bioavailability. Estimation of Solubility, Permeability, Absorption and Bioavailability* Wiley-VCH: Weinheim, 2003 (in press)
28. Avdeef, A.; Testa, B., Physicochemical profiling in drug research: A brief state-of-the-art of experimental techniques, *Cell. Mol. Life Sci.* 2002 (in press).
29. Kerns, E. H., High throughput physicochemical profiling for drug discovery, *J. Pharm. Sci.* **90**, 1838–1858 (2001).
30. Kerns, E. H.; Di, L., Multivariate pharmaceutical profiling for drug discovery, *Curr. Top. Med. Chem.* **2**, 87–98 (2002).

31. Roberts, S. A., High-throughput screening approaches for investigating drug metabolism and pharmacokinetics, *Xenobiotica* **31**, 557–589 (2001).
32. van de Waterbeemd, H.; Smith, D. A.; Jones, B. C., Lipophilicity in PK design: Methyl, ethyl, futile, *J. Comp.-Aided Mol. Design* **15**, 273–286 (2001).
33. van de Waterbeemd, H.; Smith, D. A.; Beaumont, K.; Walker, D. K., Property-based design: optimization of drug absorption and pharmacokinetics, *J. Med. Chem.* **44**, 1313–1333 (2001).
34. Perrin, D. D.; Dempsey, B.; Serjeant, E. P., *pK<sub>a</sub> Prediction for Organic Acids and Bases*, Chapman & Hall, London, 1981.
35. Albert, A.; Serjeant, E. P., *The Determination of Ionization Constants*, 3rd ed., Chapman & Hall, London, 1984.
36. Sangster, J., *Octanol-Water Partition Coefficients: Fundamentals and Physical Chemistry*, Wiley Series in Solution Chemistry, Vol. 2, Wiley, Chichester, UK, 1997.
37. Grant, D. J. W.; Higuchi, T., *Solubility Behavior of Organic Compounds*, Wiley, New York, 1990.
38. Yalkowsky, S. H.; Banerjee, S., *Aqueous Solubility: Methods of Estimation for Organic Compounds*, Marcel Dekker, New York, 1992.
39. Streng, W. H., *Characterization of Compounds in Solution—Theory and Practice*, Kluwer Academic/Plenum Publishers, New York, 2001.
40. Flynn, G. L.; Yalkowsky, S. H.; Roseman, T. J., Mass transport phenomena and models: Theoretical concepts, *J. Pharm. Sci.* **63**, 479–510 (1974).
41. Weiss, T. F., *Cellular Biophysics*, Vol. I, *Transport*, MIT Press, Cambridge, MA, 1996.
42. Dawson, D. C., Principles of membrane transport, in Field, M.; Frizzell, R. A. (eds.), *Handbook of Physiology*, Sect. 6: The gastrointestinal system, Vol. IV, *Intestinal Absorption and Secretion*, Am. Physiol. Soc., Bethesda, MD, 1991, pp. 1–44.
43. Wells, J. I., *Pharmaceutical Preformulation: The Physicochemical Properties of Drug Substances*, Ellis Horwood, Chichester, UK, 1988.
44. Schanker, L. S.; Tocco, D. J.; Brodie, B. B.; Hogben, C. A. M., Absorption of drugs from the rat small intestine, *J. Am. Chem. Soc.* **123**, 81–88 (1958).
45. Dressman, J. B.; Amidon, G. L.; Fleisher, D., Absorption potential: Estimating the fraction absorbed for orally administered compounds, *J. Pharm. Sci.* **74**, 588–589 (1985).
46. Seydel, J. K.; Coats, E. A.; Cordes, H. P.; Weise, M., Drug membrane interactions and the importance for drug transport, distribution, accumulation, efficacy and resistance, *Arch. Pharm. (Weinheim)* **327**, 601–610 (1994).
47. Chan, O. H.; Stewart, B. H., Physicochemical and drug-delivery considerations for oral drug bioavailability, *Drug Discov. Today* **1**, 461–465 (1996).
48. Borchardt, R. T.; Smith, P. L.; Wilson, G., *Models for Assessing Drug Absorption and Metabolism*, Plenum Press, New York, 1996.
49. Camenisch, G.; Folkers, G.; van de Waterbeemd, H., Review of theoretical passive drug absorption models: historical background, recent developments and limitations, *Pharm. Acta Helv.* **71**, 309–327 (1996).
50. Grass, G. M., Simulation models to predict oral drug absorption from in vitro data, *Adv. Drug. Del. Rev.* **23**, 199–219 (1997).
51. Stewart, B. H.; Chan, O. H.; Jezyk, N.; Fleisher, D., Discrimination between drug candidates using models from evaluation of intestinal absorption, *Adv. Drug Del. Rev.* **23**, 27–45 (1997).

52. Dowty, M. E.; Dietsch, C. R., Improved prediction of in vivo peroral absorption from in vitro intestinal permeability using an internal standard to control for intra- & inter-rat variability, *Pharm. Res.* **14**, 1792–1797 (1997).
53. Curatolo, W. Physical chemical properties of oral drug candidates in the discovery and exploratory settings, *Pharm. Sci. Tech. Today* **1**, 387–393 (1998).
54. Camenisch, G.; Folkers, G.; van de Waterbeemd, H., Shapes of membrane permeability-lipophilicity curves: Extension of theoretical models with an aqueous pore pathway, *Eur. J. Pharm. Sci.* **6**, 321–329 (1998).
55. Ungell, A.-L.; Nylander, S.; Bergstrand, S.; Sjöberg, Å.; Lennernäs, H., Membrane transport of drugs in different regions of the intestinal tract of the rat, *J. Pharm. Sci.* **87**, 360–366 (1998).
56. Winiwarter, S.; Bonham, N. M.; Ax, F.; Hallberg, A.; Lennernäs, H.; Karlen, A., Correlation of human jejunal permeability (in vivo) of drugs with experimentally and theoretically derived parameters. A multivariate data analysis approach, *J. Med. Chem.* **41**, 4939–4949 (1998).
57. Kristl, A.; Tukker, J. J., Negative correlation of *n*-octanol/water partition coefficient and transport of some guanine derivatives through rat jejunum in vitro, *Pharm. Res.* **15**, 499–501 (1998).
58. Krämer, S. D., Absorption prediction from physicochemical parameters, *Pharm. Sci. Tech. Today* **2**, 373–380 (1999).
59. Barthe, L.; Woodley, J.; Houin, G. Gastrointestinal absorption of drugs: Methods and studies, *Funda. Clin. Pharmacol.* **13**, 154–168 (1999).
60. Raevsky, O. A.; Fetisov, V. I.; Trepalina, E. P.; McFarland, J. W.; Schaper, K.-J., Quantitative estimation of drug absorption in humans for passively transported compounds on the basis of their physico-chemical parameters, *Quant. Struct.-Act. Relat.* **19**, 366–374 (2000).
61. Dressman, J. B.; Lennernäs, H. (eds.), *Oral Drug Absorption—Prediction and Assessment*, Marcel Dekker, New York, 2000.
62. Wilson, J. P., Surface area of the small intestine in man, *Gut* **8**, 618–621 (1967).
63. Madara, J. L., Functional morphology of epithelium of the small intestine, in Field, M.; Frizzell, R. A. (eds.), *Handbook of Physiology*, Sect. 6: The gastrointestinal system, Vol. IV, *Intestinal Absorption and Secretion*, Am. Physiol. Soc., Bethesda, MD, 1991, pp. 83–120.
64. Dressman, J. B.; Bass, P.; Ritschel, W. A.; Friend, D. R.; Rubinstein, A.; Ziv, E., Gastrointestinal parameters that influence oral medications, *J. Pharm. Sci.* **82**, 857–872 (1993).
65. Gray, V. A.; Dressman, J. B., *Pharm. Forum* **22**, 1943–1945 (1996).
66. Kararli, T. T., Comparative models for studying absorption, AAPS Workshop on Permeability Definitions and Regulatory Standards for Bioequivalence, Arlington, Aug. 17–19, 1998.
67. Charman, W. N.; Porter, C. J.; Mithani, S. D.; Dressman, J. B., The effect of food on drug absorption—a physicochemical and predictive rationale for the role of lipids and pH, *J. Pharm. Sci.* **86**, 269–282 (1997).
68. Dressman, J. B.; Amidon, G. L.; Reppas, C.; Shah, V., Dissolution testing as a prognostic tool for oral drug absorption: Immediate release dosage forms, *Pharm. Res.* **15**, 11–22 (1998).

69. Berne, R. M.; Levy, M. N., *Physiology*, 4th ed., Mosby Yearbook, St. Louis, 1998, pp. 654–661.
70. Rechcemmer, G., Transport of weak electrolytes, in Field, M.; Frizzell, R. A. (eds.), *Handbook of Physiology*, Sect. 6: The gastrointestinal system, Vol. IV, *Intestinal Absorption and Secretion*, Am. Physiol. Soc., Bethesda, MD, 1991, pp. 371–388.
71. Dressman, J. B.; Reppas, C., In vitro–in vivo correlations for lipophilic, poorly water-soluble drugs, *Eur. J. Pharm. Sci.* **11** (Suppl. 2), S73–S80 (2000).
72. Larhed, A. W.; Artursson, P.; Gråsjö, J.; Björk, E., Diffusion of drugs in native and purified gastrointestinal mucus. *J. Pharm. Sci.* **86**, 660–665 (1997).
73. Shiao, Y.-F.; Fernandez, P.; Jackson, M. J.; McMonagle, S., Mechanisms maintaining a low-pH microclimate in the intestine, *Am. J. Physiol.* **248**, G608–G617 (1985).
74. Lennernäs, H., Human intestinal permeability, *J. Pharm. Sci.* **87**, 403–410 (1998).
75. Lutz, K. L.; Siahaan, T. J., Molecular structure of the apical junction complex and its contributions to the paracellular barrier, *J. Pharm. Sci.* **86**, 977–984 (1997).
76. Jackson, M. J.; Tai, C.-Y., Morphological correlates of weak electrolyte transport in the small intestine, in Dinno, M. A. (ed.), *Structure and Function in Epithelia and Membrane Biophysics*, Alan R. Liss, New York, 1981, pp. 83–96.
77. Winne, D., Shift of pH-absorption curves, *J. Pharmacokinet. Biopharm.* **5**, 53–94 (1977).
78. Said, H. M.; Blair, J. A.; Lucas, M. L.; Hilburn, M. E., Intestinal surface acid microclimate in vitro and in vivo in the rat, *J. Lab Clin. Med.* **107**, 420–424 (1986).
79. Jackson, M. J., Drug transport across gastrointestinal epithelia, in Johnson, L. R. (ed.), *Physiology of the Gastrointestinal Tract*, 2nd ed., Raven Press, New York, 1987, pp. 1597–1621.
80. Takagi, M.; Taki, Y.; Sakane, T.; Nadai, T.; Sezaki, H.; Oku, N.; Yamashita, S., A new interpretation of salicylic acid transport across the lipid bilayer: Implication of pH-dependence but not carrier-mediated absorption from the gi tract, *J. Pharmacol. Exp. Therapeut.* **285**, 1175–1180 (1998).
81. Kimura, Y.; Hosoda, Y.; Shima, M.; Adachi, S.; Matsuno, R., Physicochemical properties of fatty acids for assessing the threshold concentration to enhance the absorption of a hydrophilic substance, *Biosci. Biotechnol. Biochem.* **62**, 443–447 (1998).
82. Yamashita, S.; Furubayashi, T.; Kataoka, M.; Sakane, T.; Sezaki, H.; Tokuda, H., Optimized conditions for prediction of intestinal drug permeability using Caco-2 cells, *Eur. J. Pharm. Sci.* **10**, 109–204 (2000).
83. Asokan, A.; Cho, M. J., Exploitation of intracellular pH gradients in the cellular delivery of macromolecules, *J. Pharm. Sci.* **91**, 903–913 (2002).
84. Antonenko, Y. N.; Bulychev, A. A., Measurements of local pH changes near bilayer lipid membrane by means of a pH microelectrode and a protonophore-dependent membrane potential. Comparison of the methods, *Biochim. Biophys. Acta* **1070**, 279–282 (1991).
85. Rojanasakul, Y.; Robinson, J. R., The cytoskeleton of the cornea and its role in tight junction permeability, *Int. J. Pharm.* **68**, 135–149 (1991).
86. Schneeberger, E. E.; Lynch, R. D., Structure, function, and regulation of cellular tight junctions, *Am. J. Physiol.* **262**, L647–L661 (1992).
87. Anderberg, E. K.; Lindmark, T.; Artursson, P., Sodium caprate elicits dilations in human intestinal tight junctions and enhances drug absorption by the paracellular route, *Pharm. Res.* **10**, 857–864 (1993).

88. Bhat, M.; Toledo-Velasquez, D.; Wang, L. Y.; Malanga, C. J.; Ma, J. K. H.; Rojanasakul, Y., Regulation of tight junction permeability by calcium mediators and cell cytoskeleton in rabbit tracheal epithelium, *Pharm. Res.* **10**, 991–997 (1993).
89. Noach, A. B. J. Enhancement of paracellular drug transport across epithelia—in vitro and in vivo studies, *Pharm. World Sci.* **17**, 58–60 (1995).
90. Lutz, K. L.; Jois, S. D. S.; Siahaan, T. J., Secondary structure of the HAV peptide which regulates cadherin-cadherin interaction, *J. Biomol. Struct. Dynam.* **13**, 447–455 (1995).
91. Tanaka, Y.; Taki, Y.; Sakane, T.; Nadai, T.; Sezaki, H.; Yamashita, S., Characterization of drug transport through tight-junctional pathway in Caco-2 monolayer: Comparison with isolated rat jejunum and colon, *Pharm. Res.* **12**, 523–528 (1995).
92. Brayden, D. J.; Creed, E.; Meehan, E.; O’Malley, K. E., Passive transepithelial diltiazem absorption across intestinal tissue leading to tight junction openings, *J. Control. Rel.* **38**, 193–203 (1996).
93. Lutz, K. L.; Szabo, L. A.; Thompson, D. L.; Siahaan, T. J., Antibody recognition of peptide sequence from the cell-cell adhesion proteins: N- and E-cadherins, *Peptide Res.* **9**, 233–239 (1996).
94. Borchard, G.; Lueßen, H. L.; de Boer, A. G.; Verhoeef, J. C.; Lehr, C.-M.; Junginger, H. E., The potential of mucoadhesive polymers in enhancing intestinal peptide drug absorption. III: Effects of chitosan-glutamate and carbomer on epithelial tight junctions in vitro, *J. Control. Rel.* **39**, 131–138 (1996).
95. Lutz, K. L.; Siahaan, T. J., Molecular structure of the apical junction complex and its contributions to the paracellular barrier, *J. Pharm. Sci.* **86**, 977–984 (1997).
96. Pal, D.; Audus, K. L.; Siahaan, T. J., Modulation of cellular adhesion in bovine brain microvessel endothelial cells by a decapeptide, *Brain Res.* **747**, 103–113 (1997).
97. Gan, L.-S. L.; Yanni, S.; Thakker, D. R., Modulation of the tight junctions of the Caco-2 cell monolayers by H2-antagonists, *Pharm. Res.* **15**, 53–57 (1998).
98. Hansch, C.; Leo, A., *Substituent Constants for Correlation Analysis in Chemistry and Biology*, Wiley-Interscience, New York, 1979.
99. Iwahashi, M.; Hayashi, Y.; Hachiya, N.; Matsuzawa, H.; Kobayashi, H., Self-association of octan-1-ol in the pure liquid state and in decane solutions as observed by viscosity, self-diffusion, nuclear magnetic resonance and near-infrared spectroscopy measurements, *J. Chem. Soc. Faraday Trans.* **89**, 707–712 (1993).
100. Franks, N. P.; Abraham, M. H.; Lieb, W. R., Molecular organization of liquid n-octanol: An X-ray diffraction analysis, *J. Pharm. Sci.* **82**, 466–470 (1993).
101. Amidon, G. L.; Lennernäs, H.; Shah, V. P.; Crison, J. R., A theoretical basis for a biopharmaceutic drug classification: the correlation of in vitro drug product dissolution and in vivo bioavailability, *Pharm. Res.* **12**, 413–420 (1995).
102. Amidon, G. L., The rationale for a biopharmaceutics drug classification, in *Biopharmaceutics Drug Classification and International Drug Regulation*, Capsugel Library, 1995, pp. 179–194.
103. Amidon, G. L.; Walgreen, C. R., Rationale and implementation of a biopharmaceutics classification system (bcs) for new drug regulation, in *Biopharmaceutics Drug Classification and International Drug Regulation*, Capsugel Library, 1998, pp. 13–27.
104. Hussain, A. S., Methods for permeability determination: A regulatory perspective. AAPS Workshop on Permeability Definitions and Regulatory Standards for Bioequivalence, Arlington, VA, Aug. 17–19, 1998.

105. *CPMP Note for Guidance on the Investigation of Bioavailability and Bioequivalence* (CPMP/EWP/QWP/1401/98 Draft), Dec. 1998.
106. *FDA Guidance for Industry Waiver of in vivo Bioavailability and Bioequivalence Studies for Immediate Release Solid Oral Dosage Forms Containing Certain Active Moieties/ Active Ingredients Based on a Biopharmaceutics Classification System*, CDER-GUID2062dft.wpd Draft, Jan. 1999.
107. Blume, H. H.; Schug, B. S., The biopharmaceutics classification system (BCS): Class III drugs—better candidates for BA/BE waiver? *Eur. J. Pharm. Sci.* **9**, 117–121 (1999).
108. Lentz, K. A.; Hayashi, J.; Lucisano, L. J.; Polli, J. E., Development of a more rapid, reduced serum culture system for Caco-2 monolayers and application to the biopharmaceutics classification system, *Int. J. Pharm.* **200**, 41–51 (2000).
109. Chen, M.-L.; Shah, V.; Patnaik, R.; Adams, W.; Hussain, A.; Conner, D.; Mehta, M.; Malinowski, H.; Lazor, J.; Huang, S.-M.; Hare, D.; Lesko, L.; Sporn, D.; Williams, R., Bioavailability and bioequivalence: an FDA regulatory overview, *Pharm. Res.* **18**, 1645–1650 (2001).
110. Rege, B. D.; Yu, L. X.; Hussain, A. S.; Polli, J. E., Effect of common excipients on Caco-2 transport of low-permeability drugs, *J. Pharm. Sci.* **90**, 1776–1786 (2001).
111. Ritschel, W. A., in Fraske, D. E.; Whitney, H. A. K., Jr. (eds.), *Perspectives in Clinical Pharmacy*, Drug Intelligence Publications, Hamilton, IL, 1972, pp. 325–367.
112. Avdeef, A., *Applications and Theory Guide to pH-Metric  $pK_a$  and log P Measurement*, Sirius Analytical Instruments Ltd. Forest Row, UK, 1993.
113. Levy, R. H.; Rowland, M., Dissociation constants of sparingly soluble substances: nonlogarithmic linear titration curves, *J. Pharm. Sci.* **60**, 1155–1159 (1971).
114. Purdie, N.; Thomson, M. B.; Riemann, N., The thermodynamics of ionization of polycarboxylic acids, *J. Solut. Chem.* **1**, 465–476 (1972).
115. Streng, W. H.; Steward, D. L., Jr., Ionization constants of an amino acid as a function of temperature, *Int. J. Pharm.* **61**, 265–266 (1990).
116. Avdeef, A.; Bucher, J. J., Accurate measurements of the concentration of hydrogen ions with a glass electrode: Calibrations using the Prideaux and other universal buffer solutions and a computer-controlled automatic titrator, *Anal. Chem.* **50**, 2137–2142 (1978).
117. Avdeef, A., Weighting scheme for regression analysis using pH data from acid-base titrations, *Anal. Chim. Acta* **148**, 237–244 (1983).
118. Avdeef, A., STBLTY: Methods for construction and refinement of equilibrium models, in Leggett, D. J. (ed.), *Computational Methods for the Determination of Formation Constants*, Plenum Press, New York, 1985, pp. 355–473.
119. Avdeef, A.; Comer, J. E. A.; Thomson, S. J., pH-metric logP. 3. Glass electrode calibration in methanol-water, applied to  $pK_a$  determination of water-insoluble substances, *Anal. Chem.* **65**, 42–49 (1993).
120. Lingane, P. J.; Hugus, Z. Z. Jr., Normal equations for the gaussian least-squares refinement of formation constants with simultaneous adjustment of the spectra of the absorbing species, *Inorg. Chem.* **9**, 757–762 (1970).
121. Hugus, Z. Z., Jr.; El-Awady, A. A., The determination of the number of species present in a system: A new matrix rank treatment of spectrophotometric data, *J. Phys. Chem.* **75**, 2954–2957 (1971).

122. Alcock, R. M.; Hartley, F. R.; Rogers, D. E., A damped non-linear least-squares computer program (dalsfek) for the evaluation of equilibrium constants from spectrophotometric and potentiometric data, *J. Chem. Soc. Dalton Trans.* 115–123 (1978).
123. Maeder, M.; Gampp, H., Spectrophotometric data reduction by eigenvector analysis for equilibrium and kinetic studies and a new method of fitting exponentials, *Anal. Chim. Acta* **122**, 303–313 (1980).
124. Gampp, H.; Maeder, M.; Zuberbühler, A. D., General non-linear least-squares program for the numerical treatment of spectrophotometric data on a single-precision game computer, *Talanta* **27**, 1037–1045 (1980).
125. Kralj, Z. I.; Simeon, V., Estimation of spectra of individual species in a multisolute solution, *Anal. Chim. Acta* **129**, 191–198 (1982).
126. Gampp, H.; Maeder, M.; Meyer, C. J.; Zuberbühler, A. D., Calculation of equilibrium constants from multiwavelength spectroscopic data. I. Mathematical considerations, *Talanta* **32**, 95–101 (1985).
127. Gampp, H.; Maeder, M.; Meyer, C. J.; Zuberbühler, A. D., Calculation of equilibrium constants from multiwavelength spectroscopic data. III. Model-free analysis of spectrophotometric and ESR titrations. *Talanta* **32**, 1133–1139 (1985).
128. Alibrandi, G., Variable-concentration kinetics, *J. Chem. Soc., Chem. Commun.* 2709–2710 (1994).
129. Alibrandi, G.; Coppolino, S.; Micali, N.; Villari, A., Variable pH kinetics: an easy determination of pH-rate profile, *J. Pharm. Sci.* **90**, 270–274 (2001).
130. Saurina, J.; Hernández-Cassou, S.; Izquierdo-Ridorsa, A.; Tauler, R., pH-gradient spectrophotometric data files from flow-injection and continuous flow systems for two- and three-way data analysis, *Chem. Intell. Lab. Syst.* **50**, 263–271 (2000).
131. Allen, R. I.; Box, K. J.; Comer, J. E. A.; Peake, C.; Tam, K. Y., Multiwavelength spectrophotometric determination of acid dissociation constants of ionizable drugs, *J. Pharm. Biomed. Anal.* **17**, 699–712 (1998).
132. Tam, K. Y.; Takács-Novák, K., Multiwavelength spectrophotometric determination of acid dissociation constants. Part II. First derivative vs. target factor analysis, *Pharm. Rese.* **16**, 374–381 (1999).
133. Mitchell, R. C.; Salter, C. J.; Tam, K. Y., Multiwavelength spectrophotometric determination of acid dissociation constants. Part III. Resolution of multi-protic ionization systems, *J. Pharm. Biomed. Anal.* **20**, 289–295 (1999).
134. Tam, K. Y.; Hadley, M.; Patterson, W., Multiwavelength spectrophotometric determination of acid dissociation constants. Part IV. Water-insoluble pyridine derivatives, *Talanta* **49**, 539–546 (1999).
135. Box, K. J.; Comer, J. E. A.; Hosking, P.; Tam, K. Y.; Trowbridge, L.; Hill, A., Rapid physicochemical profiling as an aid to drug candidate selection, in Dixon, G. K.; Major, J. S.; Rice, M. J. (eds.), *High Throughput Screening: The Next Generation*, Bios Scientific Publishers, Oxford, 2000, pp. 67–74.
136. Comer, J. E. A. High-throughput pK<sub>a</sub> and log P determination, in van de Waterbeemd, H.; Lennernäs, H.; Artursson, P. (eds.), *Drug Bioavailability. Estimation of Solubility, Permeability, Absorption and Bioavailability*, Wiley-VCH, Weinheim, 2003 (in press).
137. Tam, K. Y.; Takács-Novák, K., Multi-wavelength spectroscopic determination of acid dissociation constants: A validation study, *Anal. Chim. Acta* **434**, 157–167 (2001).

138. Tam, K. Y., Multiwavelength spectrophotometric resolution of the micro-equilibria of a triprotic amphoteric drug: Methacycline, *Mikrochim. Acta* **136**, 91–97 (2001).
139. Bevan, C. D.; Hill, A. P.; Reynolds, D. P., *Patent Cooperation Treaty*, WO 99/13328, March 18, 1999.
140. Takács-Novák, K.; Tam, K. Y., Multiwavelength spectrophotometric determination of acid dissociation constants. Part V. Microconstants and tautomeric ratios of diprotic amphoteric drugs, *J. Pharm. biomed. anal.* **17**, 1171–1182 (2000).
141. Tam, K. Y., Multiwavelength spectrophotometric determination of acid dissociation constants. Part VI. Deconvolution of binary mixtures of ionizable compounds, *Anal. Lett.* **33**, 145–161 (2000).
142. Tam, K. Y.; Quéré, L., Multiwavelength spectrophotometric resolution of the micro-equilibria of cetirizine, *Anal. Sci.* **17**, 1203–1208 (2001).
143. Hendriksen, B. A.; Sanchez-Felix, M. V.; Tam, K. Y., A new multiwavelength spectrophotometric method for the determination of the molar absorption coefficients of ionizable drugs, *Spectrosc. Lett.* **35**, 9–19 (2002).
144. Cleveland, J. A. Jr.; Benko, M. H.; Gluck, S. J.; Walbroehl, Y. M., Automated pK<sub>a</sub> determination at low solute concentrations by capillary electrophoresis, *J. Chromatogr. A* **652**, 301–308 (1993).
145. Ishihama, Y.; Oda, Y.; Asakawa, N., Microscale determination of dissociation constants of multivalent pharmaceuticals by capillary electrophoresis, *J. Pharm. Sci.* **83**, 1500–1507 (1994).
146. Jia, Z.; Ramstad, T.; Zhong, M., Medium-throughput pK<sub>a</sub> screening of pharmaceuticals by pressure-assisted capillary electrophoresis, *Electrophoresis* **22**, 1112–1118 (2001).
147. Ishihama, Y.; Nakamura, M.; Miwa, T.; Kajima, T.; Asakawa, N., A rapid method for pK<sub>a</sub> determination of drugs using pressure-assisted capillary electrophoresis with photodiode array detection in drug discovery, *J. Pharm. Sci.* **91**, 933–942 (2002).
148. Oumada, F. Z.; Ràfols, C.; Rosés, M.; Bosch, E., Chromatographic determination of aqueous dissociation constants of some water-insoluble nonsteroidal antiinflammatory drugs, *J. Pharm. Sci.* **91**, 991–999 (2002).
149. Avdeef, A.; Box, K. J.; Comer, J. E. A.; Hibbert, C.; Tam, K. Y., pH-metric logP. 10. Determination of vesicle membrane–water partition coefficients of ionizable drugs, *Pharm. Res.* **15**, 208–214 (1997).
150. Slater, B.; McCormack, A.; Avdeef, A.; Comer, J. E. A., pH-metric logP. 4. Comparison of partition coefficients determined by shake-flask, hplc and potentiometric methods, *J. Pharm. Sci.* **83**, 1280–1283 (1994).
151. Avdeef, A.; Barrett, D. A.; Shaw, P. N.; Knaggs, R. D.; Davis, S. S., Octanol-, chloroform-, and PGDP-water partitioning of morphine-6-glucuronide and other related opiates, *J. Med. Chem.* **39**, 4377–4381 (1996).
152. Davies, C. W., *Ion Association*, Butterworths, London, 1962, p. 41.
153. Avdeef, A., pH-Metric logP. 2. Refinement of partition coefficients and ionization constants of multiprotic substances, *J. Pharm. Sci.* **82**, 183–190 (1993).
154. Kortnum, G.; Vogel, W.; Andrussow, K., *Dissociation Constants of Organic Acids in Aqueous Solution*, Butterworths, London, 1961.
155. Sillén, L. G.; Martell, A. E., *Stability Constants of Metal-Ion Complexes*, Special Publication 17, Chemical Society, London, 1964.

156. Perrin, D. D., *Dissociation Constants of Organic Bases in Aqueous Solution*, Butterworths, London, 1965.
157. Sillén, L. G.; Martell, A. E., *Stability Constants of Metal-Ion Complexes*, Special Publication 25, Chemical Society, London, 1971.
158. Serjeant, E. P.; Dempsey, B., *Ionization Constants of Organic Acids in Aqueous Solution*, Pergamon, Oxford, 1979.
159. Smith, R. M.; Martell, A. E., *Critical Stability Constants*, Vols. 1–6, Plenum Press, New York, 1974.
160. Smith, R. M.; Martell, A. E.; Motekaitis, R. J., *NIST Critically Selected Stability Constants of Metal Complexes Database*, Version 5, NIST Standards Reference Database 46, U. S. Dept. Commerce, Gaithersburg, MD, 1998.
161. Avdeef, A., *Sirius Technical Application Notes (STAN)*, Vol. 1, Sirius Analytical Instruments Ltd., Forest Row, UK, 1994.
162. Avdeef, A.; Box, K. J., *Sirius Technical Application Notes (STAN)*, Vol. 2, Sirius Analytical Instruments Ltd., Forest Row, UK, 1995.
163. Bjerrum, J., *Metal-Ammine Formation in Aqueous Solution*, Haase, Copenhagen, 1941.
164. Irving, H. M.; Rossotti, H. S., The calculation of formation curves of metal complexes from pH titration curves in mixed solvents, *J. Chem. Soc.* 2904–2910 (1954).
165. Avdeef, A.; Kearney, D. L.; Brown, J. A.; Chemotti, A. R. Jr., Bjerrum plots for the determination of systematic concentration errors in titration data, *Anal. Chem.* **54**, 2322–2326 (1982).
166. Takács-Novák, K.; Box, K. J.; Avdeef, A., Potentiometric pK<sub>a</sub> determination of water-insoluble compounds. Validation study in methanol/water mixtures, *Int. J. Pharm.* **151**, 235–248 (1997).
167. Avdeef, A.; Zelazowski, A. J.; Garvey, J. S., Cadmium binding by biological ligands. 3. Five- and seven-cadmium binding in metallothionein: A detailed thermodynamic study, *Inorg. Chem.* **24**, 1928–1933 (1985).
168. Dunsmore, H. S.; Midgley, D., The calibration of glass electrodes in cells with liquid junction, *Anal. Chim. Acta* **61**, 115–122 (1972).
169. Bates, R. G., The modern meaning of pH, *CRC Crit. Rev. Anal. Chem.* **10**, 247–278 (1981).
170. Sprokholt, R.; Maas, A. H. J.; Rebelo, M. J.; Covington, A. K., Determination of the performance of glass electrodes in aqueous solutions in the physiological pH range and at the physiological sodium ion concentration, *Anal. Chim. Acta* **129**, 53–59 (1982).
171. Comer, J. E. A.; Hibbert, C., pH electrode performance under automatic management conditions, *J. Auto. Chem.* **19**, 213–224 (1997).
172. Avdeef, A.; Box, K. J.; Comer, J. E. A.; Gilges, M.; Hadley, M.; Hibbert, C.; Patterson, W.; Tam, K. Y., pH-metric logP. 11. pK<sub>a</sub> determination of water-insoluble drugs in organic solvent-water mixtures, *J. Pharm. Biomed. Anal.*, **20**, 631–641 (1999).
173. Sweeton, F. H.; Mesmer, R. E.; Baes, C. F. Jr., Acidity measurements at elevated temperatures. 7. Dissociation of water, *J. Solut. Chem.* **3**, 191–214 (1974).
174. Good, N. E.; Wingert, G. D.; Winter, W.; Connolly, T. N.; Izawa, S.; Singh, R. M. M., Hydrogen ion buffers for biological research, *Biochemistry* **5**, 467–477 (1966).
175. Vega, C. A.; Bates, R. G., Buffers for the physiological pH range: Thermodynamic constants of four substituted aminoethanesulfonic acids from 5 to 50°C, *Anal. Chem.* **48**, 1293–1296 (1976).

176. Bates, R. G.; Vega, C. A.; White, D. R. Jr., Standards for pH measurements in isotonic saline media of ionic strength  $I = 0.16$ , *Anal. Chem.* **50**, 1295–1300 (1978).
177. Sankar, M.; Bates, R. G., Buffers for the physiological pH range: Thermodynamic constants of 3-(*N*-morpholino)propanesulfonic acid from 5 to 50°C, *Anal. Chem.* **50**, 1922–1924 (1978).
178. Roy, R. N.; Gibbons, J. J.; Padron, J. L.; Moeller, J., Second-stage dissociation constants of piperazine-*N,N'*-bis(2-ethanesulfonic acid) monosodium monohydrate and related thermodynamic functions in water from 5 to 55°C, *Anal. Chem.* **52**, 2409–2412 (1980).
179. Feng, D.; Koch, W. F.; Wu, Y. C., Second dissociation constant and pH of *N*-(2-hydroxyethyl)-piperazine-*N'*-2-ethanesulfonic acid from 0 to 50°C, *Anal. Chem.* **61**, 1400–1405 (1989).
180. Feng, D.; Koch, W. F.; Wu, Y. C., Investigation of the interaction of HCl and three amino acids, HEPES, MOPS and glycine, by EMF measurements, *J. Solut. Chem.* **21**, 311–321 (1992).
181. Mizutani, M., *Z. Physik. Chem.* **116**, 350–358 (1925).
182. Mizutani, M., *Z. Physik. Chem.* **118**, 318–326 (1925).
183. Mizutani, M., *Z. Physik. Chem.* **118**, 327–341 (1925).
184. Hall, N. F.; Sprinkle, M. R., Relations between the structure and strength of certain organic bases in aqueous solution, *J. Am. Chem. Soc.* **54**, 3469–3485 (1932).
185. Kolthoff, I. M.; Lingane, J. J.; Larson, W. D., The relation between equilibrium constants in water and other solvents, *J. Am. Chem. Soc.* **60**, 2512–2515 (1938).
186. Kolthoff, I. M.; Guss, L. S., Ionization constants of acid-base indicators in methanol, *J. Am. Chem. Soc.* **60**, 2516–2522 (1938).
187. Albright, P. S.; Gosting, L. J., *J. Am. Chem. Soc.* **68**, 1061–1063 (1946).
188. Grunwald, E.; Berkowitz, B. J., The measurement and correlation of acid dissociation constants for carboxylic acids in the system ethanol-water. Activity coefficients and empirical activity functions, *J. Am. Chem. Soc.* **73**, 4939–4944 (1951).
189. Gutbezahl, B.; Grunwald, E., The acidity and basicity scale in the system ethanol-water. The evaluation of degenerate activity coefficients for single ions, *J. Am. Chem. Soc.* **75**, 565–574 (1953).
190. Marshall, P. B., Some chemical and physical properties associated with histamine antagonism, *Br. J. Pharmacol.* **10**, 270–278 (1955).
191. Bacarella, A. L.; Grunwald, E.; Marshall, H. P.; Purlee, E. L., The potentiometric measurement of acid dissociation constants and pH in the system methanol-water. pK<sub>a</sub> values for carboxylic acids and anilinium ions, *J. Org. Chem.* **20**, 747–762 (1955).
192. Shedlovsky, T.; Kay, R. L., The ionization constant of acetic acid in water-methanol mixtures at 25°C from conductance measurements, *J. Am. Chem. Soc.* **60**, 151–155 (1956).
193. Edmonson, T. D.; Goyan, J. E., The effect of hydrogen ion and alcohol concentration on the solubility of phenobarbital, *J. Am. Pharm. Assoc., Sci. Ed.* **47**, 810–812 (1958).
194. Yasuda, M., Dissociation constants of some carboxylic acids in mixed aqueous solvents, *Bull. Chem. Soc. Jpn.* **32**, 429–432 (1959).
195. de Ligny, C. L.; Rehbach, M., The liquid-liquid-junction potentials between some buffer solutions in methanol and methanol-water mixtures and a saturated KCl solution in water at 25°C, *Recl. Trav. Chim. Pays-Bas* **79**, 727–730 (1960).

196. de Ligny, C. L., The dissociation constants of some aliphatic amines in water and methanol-water mixtures at 25°C, *Recl. Trav. Chim. Pays-Bas* **79**, 731–736 (1960).
197. de Ligny, C. L.; Luykx, P. F. M.; Rehbach, M.; Wienecke, A. A., The pH of some standard solutions in methanol and methanol-water mixtures at 25°C. 1. Theoretical part, *Recl. Trav. Chim. Pays-Bas* **79**, 699–712 (1960).
198. de Ligny, C. L.; Luykx, P. F. M.; Rehbach, M.; Wienecke, A. A., The pH of some standard solutions in methanol and methanol-water mixtures at 25°C. 2. Experimental part, *Recl. Trav. Chim. Pays-Bas* **79**, 713–726 (1960).
199. Yasuda, M.; Yamsaki, K.; Ohtaki, H., *Bull. Chem. Soc. Jpn.* **23**, 1067 (1960).
200. de Ligny, C. L.; Loriaux, H.; Ruiter, A., The application of Hammett's acidity function  $\text{H}_0$  to solutions in methanol-water mixtures, *Recl. Trav. Chim. Pays-Bas* **80**, 725–739 (1961).
201. Shedlovsky, T., The behaviour of carboxylic acids in mixed solvents, in Pesce, B. (ed.), *Electrolytes*, Pergamon Press, New York, 1962, pp. 146–151.
202. Long, F. A.; Ballinger, P., Acid ionization constants of alcohols in the solvents water and deuterium oxide, in Pesce, B. (ed.), *Electrolytes*, Pergamon Press, New York, 1962, pp. 152–164.
203. Bates, R. G.; Paabo, M.; Robinson, R. A., Interpretation of pH measurements in alcohol-water solvents, *J. Phys. Chem.* **67**, 1833–1838 (1963).
204. Gelsema, W. J.; de Ligny, C. L.; Remijnse, A. G.; Blijlevens, H. A., pH-measurements in alcohol-water mixtures, using aqueous standard buffer solutions for calibration, *Recl. Trav. Chim. Pays-Bas* **85**, 647–660 (1966).
205. Woolley, E. M.; Hurkot, D. G.; Hepler, L. G., Ionization constants for water in aqueous organic mixtures, *J. Phys. Chem.* **74**, 3908–3913 (1970).
206. Wooley, E. M.; Tomkins, J.; Hepler, L. G., Ionization constants for very weak organic acids in aqueous solution and apparent ionization constants for water in aqueous organic mixtures, *J. Solut. Chem.* **1**, 341–351 (1972).
207. Fisicaro, E.; Braibanti, A., Potentiometric titrations in methanol/water medium: Intertitration variability, *Talanta* **35**, 769–774 (1988).
208. Esteso, M. A.; Gonzalez-Diaz, O. M.; Hernandez-Luis, F. F.; Fernandez-Merida, L., Activity coefficients for NaCl in ethanol-water mixtures at 25°C, *J. Solut. Chem.* **18**, 277–288 (1989).
209. Papadopoulos, N.; Avranas, A., Dissociation of salicylic acid, 2,4-, 2,5- and 2,6-dihydroxybenzoic acids in 1-propanol-water mixtures at 25°C, *J. Solut. Chem.* **20**, 293–300 (1991).
210. Li, A.; Yalkowsky, S., Solubility of organic solutes in ethanol/water mixtures, *J. Pharm. Sci.* **83**, 1735–1740 (1994).
211. Halle, J.-C.; Garboriaud, R.; Schaal, R., Etude electrochimique des mélanges d'eau et de dimethylsulfoxyde. Produit ionique apparent et niveau d'acidité, *Bull. Soc. Chim.* 1851–1857 (1969).
212. Halle, J.-C.; Garboriaud, R.; Schaal, R., Sur la réalisation de mélanges tampons dans les milieux eau-DMSO, Application à l'étude de nitrodephenylamines, *Bull. Soc. Chim.* 2047–2053 (1970).
213. Woolley, E. M.; Hepler, L. G., Apparent ionization constants of water in aqueous organic mixtures and acid dissociation constants of protonated co-solvents in aqueous solution, *Anal. Chem.* **44**, 1520–1523 (1972).

214. Yakolev, Y. B.; Kul'ba, F. Y.; Zenchenko, D. A., Potentiometric measurement of the ionic products of water in water-dimethylsulphoxide, water-acetonitrile and water-dioxane mixtures, *Russ. J. Inorg. Chem.* **20**, 975–976 (1975).
215. Siow, K.-S.; Ang, K.-P., Thermodynamic of ionization of 2,4-dinitrophenol in water-dimethylsulfoxide solvents, *J. Solut. Chem.* **18**, 937–947 (1989).
216. Van Uitert, L. G.; Haas, C. G., Studies on coordination compounds. 1. A method for determining thermodynamic equilibrium constants in mixed solvents, *J. Am. Chem. Soc.* **75**, 451–455 (1953).
217. Grunwald, E., Solvation of electrolytes in dioxane-water mixtures, in Pesce, B. (ed.), *Electrolytes*, Pergamon Press, New York, 1962, pp. 62–76.
218. Marshall, W. L., Complete equilibrium constants, electrolyte equilibria, and reaction rates, *J. Phys. Chem.* **74**, 346–355 (1970).
219. Sigvartsen, T.; Songstadt, J.; Gestblom, B.; Noreland, E., Dielectric properties of solutions of tetra-isopentylammonium nitrate in dioxane-water mixtures, *J. Solut. Chem.* **20**, 565–582 (1991).
220. Casassas, E.; Fonrodona, G.; de Juan, A., Solvatochromic parameters for binary mixtures and a correlation with equilibrium constants. 1. Dioxane-water mixtures, *J. Solut. Chem.* **21**, 147–162 (1992).
221. Cavill, G. W. K.; Gibson, N. A.; Nyholm, R. S., *J. Chem. Soc.* 2466–2470 (1949).
222. Garrett, E. R., Variation of  $pK'_a$ -values of tetracyclines in dimethylformamide-water solvents, *J. Pharm. Sci.* **52**, 797–799 (1963).
223. Hawes, J. L.; Kay, R. L., Ionic association of potassium and cesium chloride in ethanol-water mixtures from conductance measurements at 25°, *J. Phys. Chem.* **69**, 2420–2431 (1965).
224. Takács-Novák, K.; Avdeef, A.; Podányi, B.; Szász, G., Determination of protonation macro- and microconstants and octanol/water partition coefficient of anti-inflammatory niflumic acid, *J. Pharm. Biomed. Anal.* **12**, 1369–1377 (1994).
225. Avdeef, A.; Takács-Novák, K.; Box, K. J., pH-metric logP. 6. Effects of sodium, potassium, and N-CH<sub>3</sub>-D-glucamine on the octanol-water partitioning with prostaglandins E1 and E2, *J. Pharm. Sci.* **84**, 523–529 (1995).
226. Leggett, D. J., SQUAD: Stability quotients from absorbance data, in Leggett, D. J. (ed.), *Computational Methods for the Determination of Formation Constants*, Plenum Press, New York, 1985, pp. 158–217.
227. Lawson, C. L.; Hanson, R. J., *Solving Least Squares Problems*, Prentice-Hall, Englewood Cliffs, NJ, 1974.
228. Tam, K. Y. BUFMAKE: A computer program to calculate the compositions of buffers of defined buffer capacity and ionic strength for ultra-violet spectrophotometry, *Comp. Chem.* **23**, 415–419 (1999).
229. Comer, J.; Tam, K., Lipophilicity profiles: theory and measurement, in Testa, B.; van de Waterbeemd, H.; Folkers, G.; Guy, R. (eds.), *Pharmacokinetic Optimization in Drug Research*, Verlag Helvetica Chimica Acta, Zürich and Wiley-VCH, Weinheim, 2001, pp. 275–304.
230. Niebergall, P. J.; Schnaare, R. L.; Sugita, E. T., Spectral determination of microdissociation constants, *J. Pharm. Sci.* **61**, 232–234 (1972).
231. Streng, W. H., Microionization constants of commercial cephalosporins, *J. Pharm. Sci.* **67**, 666–669 (1978).

232. Noszál, B., Microspeciation of polypeptides, *J. Phys. Chem.* **90**, 6345–6349 (1986).
233. Noszál, B., Group constant: A measure of submolecular basicity, *J. Phys. Chem.* **90**, 4104–4110 (1986).
234. Noszál, B.; Osztás, E., Acid-base properties for each protonation site of six corticotropin fragments, *Int. J. Peptide Protein Res.* **33**, 162–166 (1988).
235. Noszál, B.; Sándor, P., Rota-microspeciation of aspartic acid and asparagine, *Anal. Chem.* **61**, 2631–2637 (1989).
236. Nyéki, O.; Osztás, E.; Noszál, B.; Burger, K., Acid-base properties and microspeciation of six angiotensin-type octapeptides, *Int. J. Peptide Protein Res.* **35**, 424–427 (1990).
237. Takács-Novák, K.; Noszál, B.; Hermecz, I.; Keresztúri, G.; Podányi, B.; Szász, G., Protonation equilibria of quinolone antibacterials, *J. Pharm. Sci.* **79**, 1023–1028 (1990).
238. Noszál, B.; Guo, W.; Rabenstein, D. L., Rota-microspeciation of serine, cysteine and selenocysteine, *J. Phys. Chem.* **95**, 9609–9614 (1991).
239. Noszál, B.; Rabenstein, D. L., Nitrogen-protonation microequilibria and C(2)-deprotonation microkinetics of histidine, histamine and related compounds, *J. Phys. Chem.* **95**, 4761–4765 (1991).
240. Noszál, B.; Kassai-Tanczos, R., Microscopic acid-base equilibria of arginine, *Talanta* **38**, 1439–1444 (1991).
241. Takács-Novák, K.; Józan, M.; Hermecz, I.; Szász, G., Lipophilicity of antibacterial fluoroquinolones, *Int. J. Pharm.* **79**, 89–96 (1992).
242. Noszál, B.; Guo, W.; Rabenstein, D. L., Characterization of the macroscopic and microscopic acid-base chemistry of the native disulfide and reduced dithiol forms of oxytocin, arginine-vasopressin and related peptides, *J. Org. Chem.* **57**, 2327–2334 (1992).
243. Takács-Novák, K.; Józan, M.; Szász, G., Lipophilicity of amphoteric molecules expressed by the true partition coefficient, *Int. J. Pharm.* **113**, 47 (1995).
244. Cannon, J. B.; Krill, S. L.; Porter, W. R., Physicochemical properties of A-75998, an antagonist of luteinizing hormone releasing hormone, *J. Pharm. Sci.* **84**, 953–958 (1995).
245. Dearden, J. C.; Bresnen, G. M., The measurement of partition coefficients, *Quant. Struct.-Act. Relat.* **7**, 133–144 (1988).
246. Hersey, A.; Hill, A. P.; Hyde, R. M.; Livingstone, D. J., Principles of method selection in partition studies, *Quant. Struct. Act. Relat.* **8**, 288–296 (1989).
247. Tsai, R.-S.; Fan, W.; El Tayar, N.; Carrupt, P.-A.; Testa, B.; Kier, L. B., Solute-water interactions in the organic phase of a biphasic system. 1. Structural influence of organic solutes on the “water-dragging” effect, *J. Am. Chem. Soc.* **115**, 9632–9639 (1993).
248. Fan, W.; Tsai, R. S.; El Tayar, N.; Carrupt, P.-A.; Testa, B., Solute-water interactions in the organic phase of a biphasic system. 2. Effects of organic phase and temperature on the “water-dragging” effect, *J. Phys. Chem.* **98**, 329–333 (1994).
249. Leahy, D. E.; Taylor, P. J.; Wait, A. R., Model solvent systems for QSAR. 1. Propylene glycol dipelargonate (PGDP). A new standard for use in partition coefficient determination, *Quant. Struct. Act. Relat.* **8**, 17–31 (1989).
250. Seiler, P., The simultaneous determination of partition coefficients and acidity constant of a substance, *Eur. J. Med. Chem. Chim. Therapeut.* **9**, 665–666 (1974).

251. van de Waterbeemd, H.; Kansy, M., Hydrogen-bonding capacity and brain penetration, *Chimia* **46**, 299–303 (1992).
252. von Geldern, T. W.; Hoffman, D. J.; Kester, J. A.; Nellans, H. N.; Dayton, B. D.; Calzadilla, S. V.; Marsh, K. C.; Hernandez, L.; Chiou, W.; Dixon, D. B.; Wu-wong, J. R.; Opgenorth, T. J., Azole endothelin antagonists. 3. Using  $\Delta\log P$  as a tool to improve absorption, *J. Med. Chem.* **39**, 982–991 (1996).
253. Kubinyi, H., Hydrogen bonding: The last mystery in drug design? in Testa, B.; van de Waterbeemd, H.; Folkers, G.; Guy, R. (eds.), *Pharmacokinetic Optimization in Drug Research*, Verlag Helvetica Chimica Acta, Zürich and Wiley-VCH, Weinheim, 2001, pp. 513–524.
254. Raevsky, O.; Grigor'ev, V. Y.; Kireev, D.; Zefirov, N., Complete thermodynamic description of H-bonding in the framework of multiplicative approach, *Quant. Struct. Act. Relat.*, **11**, 49–63 (1992).
255. El Tayar, N.; Testa, B.; Carrupt, P.-A., Polar intermolecular interactions encoded in partition coefficients: An indirect estimation of hydrogen-bond parameters of polyfunctional solutes, *J. Phys. Chem.* **96**, 1455–1459 (1992).
256. Chikhale, E. G.; Ng, K. Y.; Burton, P. S.; Borchardt, R. T., Hydrogen bonding potential as a determinant of the in vitro and in situ blood-brain barrier permeability of peptides, *Pharm. Res.* **11**, 412–419 (1994).
257. Abraham, M.; Chadha, H.; Whiting, G.; Mitchell, R., Hydrogen bonding. 32. An analysis of water-octanol and water-alkane partitioning and the log P parameter of Seiler, *J. Pharm. Sci.* **83**, 1085–1100 (1994).
258. ter Laak, A. M.; Tsai, R.-S.; den Kelder, G. M., D.-O.; Carrupt, P.-A.; Testa, B., Lipophilicity and hydrogen-bonding capacity of H<sub>1</sub>-antihistaminic agents in relation to their central sedative side effects, *Eur. J. Pharm. Sci.* **2**, 373–384 (1994).
259. Potts, R. O.; Guy, R. H., A predictive algorithm for skin permeability: The effects of molecular size and hydrogen bonding, *Pharm. Res.* **12**, 1628–1633 (1995).
260. Abraham, M. H.; Martins, F.; Mitchell, R. C., Algorithms for skin permeability using hydrogen bond descriptors: The problems of steroids, *J. Pharm. Pharmacol.* **49**, 858–865 (1997).
261. Reymond, F.; Steyaert, G.; Carrupt, P.-A.; Morin, D.; Tillement, J. P.; Girault, H.; Testa, B., The pH-partition profile of the anti-ischemic drug trimetazidine may explain its reduction of intracellular acidosis, *Pharm. Res.* **16**, 616–624 (1999).
262. Carrupt, P.-A.; Testa, B.; Bechalany, A.; El Tayar, N.; Descas, P.; Perrissoud, D., Morphine 6-glucuronide and morphine 3-glucuronide as molecular chameleons with unexpectedly high lipophilicity, *J. Med. Chem.* **34**, 1272–1275 (1991).
263. Gaillard, P.; Carrupt, P.-A.; Testa, B., The conformation-dependent lipophilicity of morphine glucuronides as calculated from the molecular lipophilicity potential, *Bioorg. Med. Chem. Lett.* **4**, 737–742 (1994).
264. Poulin, P.; Schoenlein, K.; Theil, F.-P., Prediction of adipose tissue: Plasma partition coefficients for structurally unrelated drugs, *J. Pharm. Sci.* **90**, 436–447 (2001).
265. Chamberlain, K.; Evans, A. A.; Bromilow, R. H., 1-Octanol/water partition coefficients ( $K_{ow}$ ) and pK<sub>a</sub> for ionizable pesticides measured by a pH-metric method, *Pest. Sci.* **47**, 265–271 (1996).
266. Pliska, V.; Testa, B.; van de Waterbeemd, H. (eds.), *Methods and Principles in Medicinal Chemistry*, Vol. 4, VCH, Weinheim, Germany, 1996.

267. Testa, B.; van de Waterbeemd, H.; Folkers, G.; Guy, R. (eds.), *Pharmacokinetic Optimization in Drug Research*, Verlag Helvetica Chimica Acta, Zürich and Wiley-VCH: Weinheim, 2001.
268. Krämer, S. D.; Gautier, J.-C.; Saudemon, P., Considerations on the potentiometric log P determination, *Pharm. Res.* **15**, 1310–1313 (1998).
269. Bouchard, G.; Pagliara, A.; van Balen, G. P.; Carrupt, P.-A.; Testa, B.; Gobry, V.; Girault, H. H.; Caron, G.; Ermondi, G.; Fruttero, R., Ionic partition diagram of the zwitterionic antihistamine cetirizine, *Helv. Chim. Acta* **84**, 375–387 (2001).
270. Scherrer, R. A.; Howard, S. M., Use of distribution coefficients in quantitative structure-activity relationships, *J. Med. Chem.* **20**, 53–58 (1977).
271. Schaper, K.-J., Simultaneous determination of electronic and lipophilic properties [pK<sub>a</sub>, P(ion), P(neutral)] for acids and bases by nonlinear regression analysis of pH-dependent partition measurements, *J. Chem. Res.* (S) 357 (1979).
272. Taylor, P. J.; Cruickshank, J. M., Distribution coefficients of atenolol and sotalol, *J. Pharm. Pharmacol.* **36**, 118–119 (1984).
273. Taylor, P. J.; Cruickshank, J. M., Distribution coefficients of atenolol and sotalol, *J. Pharm. Pharmacol.* **37**, 143–144 (1985).
274. Manners, C. N.; Payling, D. W.; Smith, D. A., Distribution coefficient, a convenient term for the relation of predictable physico-chemical properties to metabolic processes, *Xenobiotica* **18**, 331–350 (1988).
275. Avdeef, A., Assessment of distribution-pH profiles, in Pliska, V.; Testa, B.; van de Waterbeemd, H. (eds.), *Methods and Principles in Medicinal Chemistry*, Vol. 4, VCH, Weinheim, Germany, 1996, pp. 109–139.
276. Scherrer, R. A., Biolipid pK<sub>a</sub> values in the lipophilicity of ampholytes and ion pairs, in Testa, B.; van de Waterbeemd, H.; Folkers, G.; Guy, R. (eds.), *Pharmacokinetic Optimization in Drug Research*, Verlag Helvetica Chimica Acta, Zürich and Wiley-VCH, Weinheim, 2001, pp. 351–381.
277. Murthy, K. S.; Zografi, G., Oil-water partitioning of chlorpromazine and other phenothiazine derivatives using dodecane and n-octanol, *J. Pharm. Sci.* **59**, 1281–1285 (1970).
278. Tomlinson, E.; Davis, S. S., Interactions between large organic ions of opposite and unequal charge. II. Ion pair and ion triplet formation, *J. Colloid Interface Sci.* **74**, 349–357 (1980).
279. van der Giesen, W. F.; Janssen, L. H. M., *Int. J. Pharm.* **12**, 231–249 (1982).
280. Scherrer, R. A., The treatment of ionizable compounds in quantitative structure-activity studies with special consideration of ion partitioning, in Magee, P. S.; Kohn, G. K.; Menn, J. J. (eds.), *Pesticide Syntheses through Rational Approaches*, ACS Symposium Series 225, ACS (American Chemical Society), Washington, DC, 1984, pp. 225–246.
281. Scherrer, R. A.; Crooks, S. L., Titrations in water-saturated octanol: A guide to partition coefficients of ion pairs and receptor-site interactions, *Quant. Struct.-Act. Relat.* **8**, 59–62 (1989).
282. Akamatsu, M.; Yoshida, Y.; Nakamura, H.; Asao, M.; Iwamura, H.; Fujita, T., Hydrophobicity of di- and tripeptides having unionizable side chains and correlation with substituent and structural parameters, *Quant. Struct. -Act. Relat.* **8**, 195–203 (1989).

283. Manners, C. N.; Payling, D. W.; Smith, D. A., Lipophilicity of zwitterionic suphate conjugates of tiaramide, propanolol and 4'-hydroxypropranolol, *Xenobiotica* **19**, 1387–1397 (1989).
284. Westall, J. C.; Johnson, C. A.; Zhang, W., Distribution of LiCl, NaCl, KCl, HCl, MgCl<sub>2</sub>, and CaCl<sub>2</sub> between octanol and water, *Environ. Sci. Technol.* **24**, 1803–1810 (1990).
285. Chmelík, J.; Hudeček, J.; Putyera, K.; Makovička, J.; Kalous, V.; Chmelíková, J., Characterization of the hydrophobic properties of amino acids on the basis of their partition and distribution coefficients in the 1-octanol-water system, *Collect. Czech. Chem. Commun.* **56**, 2030–2040 (1991).
286. Tsai, R.-S.; Testa, B.; El Tayar, N.; Carrupt, P.-A., Structure-lipophilicity relationships of zwitterionic amino acids, *J. Chem. Soc. Perkin Trans. 2*, 1797–1802 (1991).
287. Akamatsu, M.; Fujita, T., Quantitative analysis of hydorphobicity of di- to pentapeptides having un-ionizable side chains with substituent and structural parameters, *J. Pharm. Sci.* **81**, 164–174 (1992).
288. Akamatsu, M.; Katayama, T.; Kishimoto, D.; Kurokawa, Y.; Shibata, H., Quantitative analysis of the structure-hydophobicity relationships for n-acetyl di- and tripeptide amides, *J. Pharm. Sci.* **83**, 1026–1033 (1994).
289. Abraham, M. H.; Takács-Novák, K.; Mitchell, R. C., On the partition of ampholytes: Application to blood-brain distribution, *J. Pharm. Sci.* **86**, 310–315 (1997).
290. Bierer, D. E.; Fort, D. M.; Mendez, C. D.; Luo, J.; Imbach, P. A.; Dubenko, L. G.; Jolad, S. D.; Gerber, R. E.; Litvak, J.; Lu, Q.; Zhang, P.; Reed, M. J.; Waldeck, N.; Bruening, R. C.; Noamesi, B. K.; Hector, R. F.; Carlson, T. J.; King, S. R., Ethnobotanical-directed discovery of the antihyperglycemic properties of cryptolepine: Its isolation from Cryptolepis sanguinolenta, synthesis, and in vitro and in vivo activities, *J. Med. Chem.* **41**, 894–901 (1998).
291. Takács-Novák, K.; Szász, G., Ion-pair partition of quaternary ammonium drugs: The influence of counter ions of different lipophilicity, size, and flexibility, *Pharm. Res.* **16**, 1633–1638 (1999).
292. Valkó, K.; Slégl, P., New chromatographic hydrophobicity index ( $\phi_0$ ) based on the slope and the intercept of the log  $k'$  versus organic phase concentration plot, *J. Chromatogr.* **631**, 49–61 (1993).
293. Valkó, K.; Bevan, C.; Reynolds, D., Chromatographic hydrophobicity index by fast-gradient RP-HPLC: A high-throughput alternative to logP/logD, *Anal. Chem.* **69**, 2022–2029 (1997).
294. Hitzel, L.; Watt, A. P.; Locker, K. L., An increased throughput method for the determination of partition coefficients, *Pharm. Res.* **17**, 1389–1395 (2000).
295. Schräder, W.; Andersson, J. T., Fast and direct method for measuring 1-octanol-water partition coefficients exemplified for six local anesthetics, *J. Pharm. Sci.* **90**, 1948–1954 (2001).
296. Lombardo, F.; Shalaeva, M. Y.; Tupper, K. A.; Gao, F.; Abraham, M. H., E logP<sub>oct</sub>: A tool for lipophilicity determination in drug discovery, *J. Med. Chem.* **43**, 2922–2928 (2000).
297. Lombardo, F.; Shalaeva, M. Y.; Tupper, K. A.; Gao, F., E logD<sub>oct</sub>: A tool for lipophilicity determination in drug discovery. 2. Basic and neutral compounds, *J. Med. Chem.* **43**, 2922–2928 (2000).

298. Valkó, K., Measurements of physical properties for drug design in industry, in Valkó, K. (ed.), *Separation Methods in Drug Synthesis and Purification*, Elsevier, Amsterdam, 2001, Ch. 12.
299. Pidgeon, C.; Venkataram, U. V., Immobilized artificial membrane chromatography: Supports composed of membrane lipids, *Anal. Biochem.* **176**, 36–47 (1989).
300. Pidgeon, C.; Ong, S.; Choi, H.; Liu, H., Preparation of mixed ligand immobilized artificial membranes for predicting drug binding to membranes, *Anal. Chem.* **66**, 2701–2709 (1994).
301. Pidgeon, C.; Ong, S.; Liu, H.; Qui, X.; Pidgeon, M.; Dantzig, A. H.; Munroe, J.; Hornback, W. J.; Kasher, J. S.; Glunz, L.; Szczerba, T., IAM chromatography: An in vitro screen for predicting drug membrane permeability, *J. Med. Chem.* **38**, 590–594 (1995).
302. Ong, S.; Liu, H.; Qiu, X.; Bhat, G.; Pidgeon, C., Membrane partition coefficients chromatographically measured using immobilized artificial membrane surfaces, *Anal. Chem.* **67**, 755–762 (1995).
303. Ong, S.; Pidgeon, C., Thermodynamics of solute partitioning into immobilized artificial membranes, *Anal. Chem.* **67**, 2119–2128 (1995).
304. Ong, S.; Liu, H.; Pidgeon, C., Immobilized-artificial-membrane chromatography: Measurements of membrane partition coefficient and predicting drug membrane permeability, *J. Chromatogr. A* **728**, 113–128 (1996).
305. Barton, P.; Davis, A. M.; McCarthy, D. J.; Webborn, P. J. H., Drug-phospholipid interactions. 2. Predicting the sites of drug distribution using n-octanol/water and membrane/water distribution coefficients. *J. Pharm. Sci.* **86**, 1034–1039 (1997).
306. Barbato, F.; La Rotonda, M. I.; Quaglia, F., Interactions of nonsteroidal antiinflammatory drugs with phospholipids: comparison between octanol/buffer partition coefficients and chromatographic indexes on immobilized artificial membranes, *J. Pharm. Sci.* **86**, 225–229 (1997).
307. Yang, C. Y.; Cai, S. J.; Liu, H.; Pidgeon, C., Immobilized artificial membranes—screens for drug-membrane interactions, *Adv. Drug Deliv. Rev.* **23**, 229–256 (1997).
308. Stewart, B. H.; Chung, F. Y.; Tait, B.; John, C.; Chan, O. H., Hydrophobicity of HIV protease inhibitors by immobilized artificial membrane chromatography, *Pharm. Res.* **15**, 1401–1406 (1998).
309. Stewart, B. H.; Chan, O. H., Use of immobilized artificial membrane chromatography for drug transport applications, *J. Pharm. Sci.* **87**, 1471–1478 (1998).
310. Ottiger, C.; Wunderli-Allenspach, H., Immobilized artificial membrane (IAM)-HPLC for partition studies of neutral and ionized acids and bases in comparison with the liposomal partition system, *Pharm. Res.* **16**, 643–650 (1999).
311. Taillardat-Bertschinger, A.; Martinet, C. A. M.; Carrupt, P.-A.; Reist, M.; Caron, G.; Fruttero, R.; Testa, B., Molecular factors influencing retention on immobilized artificial membranes (IAM) compared to partitioning in liposomes and n-octanol, *Pharm. Res.* **19**, 729–737 (2002).
312. Genty, M.; González, G.; Clere, C.; Desangle-Gouty, V.; Legendre, J.-Y., Determination of the passive absorption through the rat intestine using chromatographic indices and molar volume, *Eur. J. Pharm. Sci.* **12**, 223–229 (2001).
313. Miyake, K.; Kitaura, F.; Mizuno, N., Phosphatidylcholine-coated silica as a useful stationary phase for high-performance liquid chromatographic determination of partition coefficients between octanol and water, *J. Chromatogr.* **389**, 47–56 (1987).

314. Yang, Q.; Lundahl, P., Binding of lysozyme on the surface of entrapped phosphatidylserine-phosphatidylcholine vesicles and an example of high-performance lipid vesicle surface chromatography, *J. Chromatogr.* **512**, 377–386 (1990).
315. Zhang, Y.; Zeng, C.-M.; Li, Y.-M.; Stellan, H.; Lundahl, P., Immobilized liposome chromatography of drugs on capillary continuous beds for model analysis of drug-membrane interactions, *J. Chromatogr. A* **749**, 13–18 (1996).
316. Lundahl, P.; Beigi, F., Immobilized liposome chromatography of drugs for model analysis of drug-membrane interactions, *Adv. Drug Deliv. Rev.* **23**, 221–227 (1997).
317. Österberg, T.; Svensson, M.; Lundahl, P., Chromatographic retention of drug molecules on immobilized liposomes prepared from egg phospholipids and from chemically pure phospholipids, *Eur. J. Pharm. Sci.* **12**, 427–439 (2001).
318. Danelian, E.; Karlén, A.; Karlsson, R.; Winiwarter, S.; Hansson, A.; Löfås, S.; Lennernäs, H.; Hämäläinen, M. D., SPR biosensor studies of the direct interaction between 27 drugs and a liposome surface: Correlation with fraction absorbed in humans, *J. Med. Chem.* **43**, 2083–2086 (2000).
319. Loidl-Stahlhofen, A.; Hartmann, T.; Schöttner, M.; Röhrling, C.; Brodowsky, H.; Schmitt, J.; Keldenich, J., Multilamellar liposome and solid-supported lipid membranes (TRAN-SIL): Screening of lipid-water partitioning toward a high-throughput scale, *Pharm. Res.* **18**, 1782–1788 (2001).
320. Ishihama, Y.; Oda, Y.; Uchikawa, K.; Asakawa, N., Evaluation of solute hydrophobicity by microemulsion electrokinetic chromatography, *Anal. Chem.* **67**, 1588–1595 (1995).
321. Razak, J. L.; Cutak, B. J.; Larive, C. K.; Lunte, C. E., Correlation of the capacity factor in vesicular electrokinetic chromatography with the octanol:water partition coefficient for charged and neutral analytes, *Pharm. Res.* **18**, 104–111 (2001).
322. Burns, S. T.; Khaledi, M. G., Rapid determination of liposome-water partition coefficient ( $K_{lw}$ ) using liposome electrokinetic chromatography (LEKC), *J. Pharm. Sci.* **91**, 1601–1612 (2002).
323. Schräder, W.; Andersson, J. T., Fast and direct method for measuring 1-octanol-water partition coefficients exemplified for six local anesthetics, *J. Pharm. Sci.* **90**, 1948–1954 (2001).
324. Dyrssen, D., Studies on the extraction of metal complexes. IV. The dissociation constants and partition coefficients of 8-quinolinol (oxine) and N-nitroso-N-phenylhydroxylamine (cupferron), *Sv. Kem. Tidsks.* **64**, 213–224 (1952).
325. Rydberg, J., Studies on the extraction of metal complexes. IX. The distribution of acetylacetone between chloroform or hexane and water, *Sv. Kem. Tidskr.* **65**, 37–43 (1953).
326. Hök, B., Studies on the extraction of metal complexes. XV. The dissociation constants of salicylic acid, 3,5-dinitrobenzoic acid, and cinnamic acid and the distribution between chloroform-water and methyl isobutyl ketone (hexane)-water, *Sv. Kem. Tidskr.* **65**, 182–194 (1953).
327. Dyrssen, D., Studies on the extraction of metal complexes. XVIII. The dissociation constant and partition coefficient of tropolone, *Acta Chem. Scand.* **8**, 1394–1397 (1954).
328. Dyrssen, D.; Dyrssen, M.; Johansson, E., Studies on the extraction of metal complexes. XXXI. Investigation with some 5,7-dihalogen derivatives of 8-quinolinol, *Acta Chem. Scand.* **10**, 341–352 (1956).

329. Hök-Bernström, B., Studies on the extraction of metal complexes. XXIII. On the complex formation of thorium with salicylic acid, methoxybenzoic acid and cinnamic acid, *Acta Chem. Scand.* **10**, 174–186 (1956).
330. Dyrssen, D., Studies on the extraction of metal complexes. XXXII. N-phenylbenzo-hydroxamic acid, *Acta Chem. Scand.* **10**, 353–359 (1957).
331. Dyrssen, D., Studies on the extraction of metal complexes. XXX. The dissociation, distribution, and dimerization of di-n-butyl phosphate (DBP), *Acta Chem. Scand.* **11**, 1771–1786 (1957).
332. Brändström, A., A rapid method for the determination of distribution coefficient of bases for biological purposes, *Acta Chem. Scand.* **17**, 1218–1224 (1963).
333. Korenman, I. M.; Gur'ev, I. A., Acid-base titration in the presence of extractants, *J. Anal. Chem. USSR* **30**, 1601–1604 (Engl.) (1975).
334. Kaufman, J. J.; Semo, N. M.; Koski, W. S., Microelectrometric titration measurement of the  $pK_a$ 's and partition and drug distribution coefficients of narcotics and narcotic antagonists and their pH and temperature dependence, *J. Med. Chem.* **18**, 647–655 (1975).
335. Johansson, P.-A.; Gustavii, K., Potentiometric titration of ionizable compounds in two phase systems. 2. Determination of partition coefficients of organic acids and bases, *Acta Pharm. Suecica* **13**, 407–420 (1976).
336. Johansson, P.-A.; Gustavii, K., Potentiometric titration of ionizable compounds in two phase systems. 3. Determination of extraction constants, *Acta Pharm. Suecica* **14**, 1–20 (1977).
337. Gur'ev, I. A.; Korenman, I. M.; Aleksandrova, T. G.; Kutsovskaya, V. V., Dual-phase titration of strong acids, *J. Anal. Chem. USSR* **32**, 192–195 (Engl.) (1977).
338. Gur'ev, I. A.; Gur'eva, Z. M., Dual-phase titration of aromatic acids, *J. Anal. Chem. USSR* **32**, 1933–1935 (Engl.) (1977).
339. Gur'ev, I. A.; Kiseleva, L. V., Dual-phase titration of aliphatic acids in the presence of neutral salts, *J. Anal. Chem. USSR* **33**, 427–430 (Engl.) (1978).
340. Gur'ev, I. A.; Zinina, O. B., Improving conditions for titrating acids by using dual-phase systems, *J. Anal. Chem. USSR* **33**, 1100–1102 (Engl.) (1978).
341. Gur'ev, I. A.; Gushchina, E. A.; Mitina, E. N., Dual-phase titration of phenols with an ion selective electrode, *J. Anal. Chem. USSR* **34**, 913–916 (Engl.) (1979).
342. Gur'ev, I. A.; Gushchina, E. A.; Gadashovich, M. Z., Dual-phase potentiometric titration of aromatic carboxylic acids with liquid ion selective electrodes, *J. Anal. Chem. USSR* **36**, 803–810 (Engl.) (1981).
343. Gur'ev, I. A.; Gushchina, E. A., Use of extraction parameters to predict the results of the dual-phase titration of phenolate ions with a liquid ion-selective electrode, *J. Anal. Chem. USSR* **37**, 1297–1301 (Engl.) (1982).
344. Clarke, F. H., Ionization constants by curve fitting. Application to the determination of partition coefficients, *J. Pharm. Sci.* **73**, 226–230 (1984).
345. Clarke, F. H.; Cahoon, N. M., Ionization constants by curve fitting: Determination of partition and distribution coefficients of acids and bases and their ions, *J. Pharm. Sci.* **76**, 611–620 (1987).
346. Avdeef, A., Fast simultaneous determination of  $\log P$  and  $pK_a$  by potentiometry: Para-alkoxyphenol series (methoxy to pentoxy), in Silipo, C.; Vittoria, A. (eds.), *QSAR: Rational Approaches to the Design of Bioactive Compounds*, Elsevier, Amsterdam, 1991, pp. 119–122.

347. Avdeef, A., pH-metric log P. I. Difference plots for determining ion-pair octanol-water partition coefficients of multiprotic substances, *Quant Struct. Act. Relat.* **11**, 510–517 (1992).
348. Avdeef, A.; Comer, J. E. A., Measurement of  $pK_a$  and  $\log P$  of water-insoluble substances by potentiometric titration, in Wermuth, C. G. (ed.), *QSAR and Molecular Modelling*, Escom, Leiden, 1993, pp. 386–387.
349. Comer, J. E. A., The acid test; ionization and lipophilicity of drugs; influence on biological activity, *Chem. Britain* **30**, 983–986 (1994).
350. Clarke, F. H.; Cahoon, N. M., Ionization constants by curve fitting: determination of partition and distribution coefficients of acids and bases and their ions, *J. Pharm. Sci.* **83**, 1524 (1994).
351. Clarke, F. H.; Cahoon, N. M., Potentiometric determination of the partition and distribution coefficients of dianionic compounds, *J. Pharm. Sci.* **84**, 53–54 (1995).
352. Comer, J. E. A.; Avdeef, A.; Box, K. J., Limits for successful measurement of  $pK_a$  and  $\log P$  by pH-metric titration, *Am. Lab.* **4**, 36c–36i (1995).
353. Herbette, L. G.; Vecchiarelli, M.; Trumltz, G., *NSAID Mechanism of Action: Membrane Interactions in the Role of Intracellular Pharmacokinetics*, Vane, Botting, Botting, pp. 85–102 (1995).
354. Karagiannis, H.; van de Waterbeemd, H., The prediction of the lipophilicity of peptidomimetics. A comparison between experimental and theoretical lipophilicity values of renin inhibitors and the building blocks, *Pharm. Acta Helv.* **70**, 67–77 (1995).
355. Danielsson, L.-G.; Zhang, Y.-H., Methods for determining n-octanol-water partition constants, *Trends Anal. Chem.* **15**, 188–196 (1996).
356. Caron, G.; Pagliara, A.; Gaillard, P.; Carrupt, P.-A.; Testa, B., Ionization and partitioning profiles of zwitterions: The case of the anti-inflammatory drug azapropazone, *Helv. Chim. Acta* **79**, 1683–1695 (1996).
357. Takács-Novák, K.; Avdeef, A., Interlaboratory study of log P determination by shake-flask and potentiometric methods, *J. Pharm. Biomed. Anal.* **14**, 1405–1413 (1996).
358. McFarland, J. W.; Berger, C. M.; Froshauer, S. A.; Hayashi, S. F.; Hecker, S. J.; Jaynes, B. H.; Jefson, M. R.; Kamicker, B. J.; Lipinski, C. A.; Lundy, K. M.; Reese, C. P.; Vu, C. B., Quantitative structure-activity relationships among macrolide antibacterial agents: In vitro and in vivo potency against pasteurella multocida, *J. Med. Chem.* **40**, 1340–1346 (1997).
359. Caron, G.; P. Gaillard, P.; Carrupt, P.-A.; Testa, B., Lipophilicity behavior of model and medicinal compounds containing a sulfide, sulfoxide, or sulfone moiety, *Helv. Chim. Acta* **80**, 449–462 (1997).
360. Herbette, L. G.; Vecchiarelli, M.; Leonardi, A., Lercanidipine: Short plasma half-life, long duration of action, *J. Cardiovasc. Pharmacol.*, **29** (Suppl 1), S19–S24 (1997).
361. Morgan, M. E.; Liu, K.; Anderson, B. D., Microscale titrimetric and spectrophotometric methods for determination of ionization constants and partition coefficients of new drug candidates, *J. Pharm. Sci.* **87**, 238–245 (1998).
362. Caron, G.; Steyaert, G.; Pagliara, A.; Reymond, F.; Crivori, P.; Gaillard, P.; Carrupt, P.-A.; Avdeef, A.; Comer, J.; Box, K. J.; Girault, H. H.; Testa, B., Structure-lipophilicity relationships of neutral and protonated  $\beta$ -blockers. Part I: Intra and intermolecular effects in isotropic solvent systems, *Helv. Chim. Acta* **82**, 1211–1222 (1999).

363. Franke, U.; Munk, A.; Wiese, M., Ionization constants and distribution coefficients of phenothiazines and calcium channel antagonists determined by a pH-metric method and correlation with calculated partition coefficients, *J. Pharm. Sci.* **88**, 89–95 (1999).
364. Leo, A.; Hansch, C.; Elkins, D., Partition coefficients and their uses, *Chem. Rev.* **71**, 525–616 (1971).
365. Sangster, J., Log KOW Databank, Montreal, Quebec, Canada, Sangster Research Laboratories, 1994.
366. Howard, P. H.; Meylan, W., *PHYSPROP Database*, Syracuse Research Corp., Syracuse, NY, 2000.
367. *Physicochemical Parameter Database*, Medicinal Chemistry Project, Pomona College, Claremont, CA, USA ([www.biobyte.com/bb/prod/cqsar.html](http://www.biobyte.com/bb/prod/cqsar.html)).
368. Krämer, S. D.; Jakits-Deiser, C.; Wunderli-Allenspach, H., Free-fatty acids cause pH-dependent changes in the drug-lipid membrane interactions around the physiological pH, *Pharm. Sci.* **14**, 827–832 (1997).
369. Jonkman, J. H. G.; Hunt, C. A., Ion pair absorption of ionized drugs—fact or fiction? *Pharm. Weekblad Sci. Ed.* **5**, 41–47 (1983).
370. Dollery, C. (ed.), *Therapeutic Drugs*, 2nd ed., Vols. 1 and 2, Churchill Livingstone, Edinburgh, 1999.
371. El Tayar, N.; Tsai, R.-S.; Carrupt, P.-A.; Testa, B., Octan-1-ol-water partition coefficients of zwitterionic  $\alpha$ -amino acids. Determination by centrifugal partition chromatography and factorization into steric/hydrophobic and polar components, *J. Chem. Soc. Perkin Trans. 2*, 79–84 (1992).
372. Wilson, C. G.; Tomlinson, E.; Davis, S. S.; Olejnik, O., Altered ocular absorption and disposition of sodium cromoglycate upon ion-pair and complex coacervate formation with dodecylbenzyldimethyl-ammonium chloride, *J. Pharm. Pharmacol.* **31**, 749–753 (1981).
373. Megwa, S. A.; Cross, S. E.; Benson, H. A. E.; Roberts, M. S., Ion-pair formation as a strategy to enhance topical delivery of salicylic acid, *J. Pharm. Pharmacol.* **52**, 919–928 (2000).
374. M. S. Mirrlees, M. S.; S. J. Moulton, S. J.; C. T. Murphy; C. T.; P. J. Taylor, P. J., Direct measurement of octanol-water partition coefficients by high-pressure liquid chromatography, *J. Med. Chem.* **19**, 615–619 (1976).
375. Unger, S. H.; Cook, J. R.; Hollenberg, J. S., Simple procedure for determining octanol-aqueous partition, distribution, and ionization coefficients by reverse-phase high-pressure liquid chromatography, *J. Pharm. Sci.* **67**, 1364–1366 (1978).
376. Loidl-Stahlhofen, A.; Eckert, A.; Hartmann, T.; Schöttner, M., Solid-supported lipid membranes as a tool for determination of membrane affinity: High-throughput screening of a physicochemical parameter, *J. Pharm. Sci.* **90**, 599–606 (2001).
377. Terada, H.; Murayama, W.; Nakaya, N.; Nunogaki, Y.; Nunogaki, K.-I., Correlation of hydrophobic parameters of organic compounds determined by centrifugal partition chromatography with partition coefficients between octanol and water, *J. Chromatogr.* **400**, 343–351 (1987).
378. El Tayar, N.; Tsai, R.-S.; Vallat, P.; Altomare, C.; Testa, B., Measurement of partition coefficients by various centrifugal partition chromatographic techniques. A comparative evaluation, *J. Chromatogr.* **556**, 181–194 (1991).

379. Tsai, R.-S.; El Tayar; N.; Carrupt, P.-A.; Testa, B., Physicochemical properties and transport behavior of piribedil: Considerations on its membrane-crossing potential, *Int. J. Pharm.* **80**, 39–49 (1992).
380. Cevc, G. (ed.), *Phospholipid Handbook*, Marcel Dekker, New York, 1993.
381. Miyoshi, H.; Maeda, H.; Tokutake, N.; Fujita, T., Quantitative analysis of partition behavior of substituted phenols from aqueous phase into liposomes made of lecithin and various lipids, *Bull. Chem. Soc. Jpn.* **60**, 4357–4362 (1987).
382. Escher, B. I.; Schwarzenbach, R. P., Partitioning of substituted phenols in liposome-water, biomembrane-water, and octanol-water systems, *Environ. Sci. Tech.* **30**, 260–270 (1996).
383. Escher, B. I.; Snozzi, M.; Schwarzenbach, R. P., Uptake, speciation, and uncoupling activity of substituted phenols in energy transducing membranes, *Environ. Sci. Technol.* **30**, 3071–3079 (1996).
384. Escher, B. I.; Hunziker, R.; Schwarzenbach, R. P.; Westall, J. C., Kinetic model to describe the intrinsic uncoupling activity of substituted phenols in energy transducing membranes, *Environ. Sci. Technol.* **33**, 560–570 (1999).
385. Balon, K.; Mueller, B. W.; Riebesehl, B. U., Determination of liposome partitioning of ionizable drugs by titration, *Pharm. Res.* **16**, 802–806 (1999).
386. Balon, K.; Mueller, B. W.; Riebesehl, B. U., Drug liposome partitioning as a tool for the prediction of human passive intestinal absorption, *Pharm. Res.* **16**, 882–888 (1999).
387. Fruttero, R.; Caron, G.; Fornatto, E.; Boschi, D.; Ermondi, G.; Gasco, A.; Carrupt, P.-A.; Testa, B., Mechanism of liposome/water partitioning of (p-methylbenzyl)alkylamines, *Pharm. Res.* **15**, 1407–1413 (1998).
388. Allan, D., Mapping the lipid distribution in the membranes of BHK cells, *Molec. Membr. Biol.* **13**, 81–84 (1996).
389. Hope, M. J.; Bally, M. B.; Webb, G.; Cullis, P. R., Production of large unilamellar vesicles by a rapid extrusion procedure. Characterization of size distribution, trapped volume and ability to maintain a membrane potential, *Biochim. Biophys. Acta* **812**, 55–65 (1985).
390. Mayer, L. D.; Hope, M. J.; Cullis, P. R.; Janoff, A. S., Solute distributions and trapping efficiencies observed in freeze-thawed multilamellar vesicles, *Biochim. Biophys. Acta* **817**, 193–196 (1985).
391. Mayer, L. D.; Bally, M. B.; Hope, M. J.; Cullis, P. R., Techniques for encapsulating bioactive agents into liposomes, *Chem. Phys. Lipids* **40**, 333–345 (1986).
392. Perkins, W. R.; Minchey, S. R.; Ahl, P. H.; Janoff, A. S., The determination of liposome captured volume, *Chem. Phys. Lipids* **64**, 197–217 (1993).
393. Lasic, D. D.; Needham, D., The “stealth” liposome. A prototypical biomaterial, *Chem. Rev.* **95**, 2601–2628 (1995).
394. Davis, S. S.; James, M. J.; Anderson, N. H., The distribution of substituted phenols in lipid vesicles, *Faraday Discuss. Chem. Soc.* **81**, 313–327 (1986).
395. Bäuerle, H.-D.; Seelig, J., Interaction of charged and uncharged calcium channel antagonists with phospholipid membranes. Binding equilibrium, binding enthalpy, and membrane location, *Biochemistry* **30**, 7203–7211 (1991).
396. Miyazaki, J.; Hideg, K.; Marsh, D., Interfacial ionization and partitioning of membrane-bound local anesthetics, *Biochim. Biophys. Acta* **1103**, 62–68 (1992).
397. Thomas, P. G.; Seelig, J., Binding of the calcium antagonist flunarizine to phosphatidyl-choline bilayers: Charge effects and thermodynamics, *Biochem. J.* **291**, 397–402 (1993).

398. Wenk, M. R.; Fahr, A.; Reszka, R.; Seelig, J., Paclitaxel partitioning into lipid bilayers, *J. Pharm. Sci.* **85**, 228–231 (1996).
399. Yeagle, P. L.; Hutton, W. C.; Huang, C.-H.; Martin, R. B., Headgroup conformation and lipid-cholesterol association in phosphatidylcholine vesicles: A  $^{31}\text{P}\{^1\text{H}\}$  nuclear Overhauser effect study, *Proc. Natl. Acad. Sci.* **72**, 3477–3481 (1975).
400. Gur, Y.; Ravina, I.; Babchin, A. J., On the electrical double layer theory. II. The Poisson-Boltzman equation including hydration forces, *J. Colloid Inter. Sci.* **64**, 333–341 (1978).
401. Brown, M. F.; Seelig, J., Influence of cholesterol on the polar region of phosphatidylcholine and phosphatidylethanolamine bilayers, *Biochemistry* **17**, 381–384 (1978).
402. Eisenberg, M.; Gresalfi, T.; Riccio, T.; McLaughlin, S., Adsorption of monovalent cations to bilayer membranes containing negative phospholipids, *Biochemistry* **18**, 5213–5223 (1979).
403. Cevc, G.; Watts, A.; Marsh, D., Titration of the phase transition of phosphatidylserine bilayer membranes. Effects of pH, surface electrostatics, ion binding, and head group hydration, *Biochemistry* **20**, 4955–4965 (1981).
404. Rooney, E. K.; Lee, A. G., Binding of hydrophobic drugs to lipid bilayers and to the  $(\text{Ca}^{2+} + \text{Mg}^{2+})$ -ATPase, *Biochim. Biophys. Acta* **732**, 428–440 (1983).
405. Cevc, G.; Marsh, D., Properties of the electrical double layer near the interface between a charged bilayer membrane and electrolyte solution: Experiment vs. theory, *J. Phys. Chem.* **87**, 376–379 (1983).
406. Cevc, G., Membrane electrostatics, *Biochim. Biophys. Acta* **1031**, 311–382 (1990).
407. McLaughlin, S., Electrostatic potentials at membrane-solution interfaces, *Curr. Topics Membr. Transport* **9**, 71–144 (1977).
408. Shnitzky, M. (ed.), *Biomembranes: Physical Aspects*, VCH, Weinheim, 1993.
409. Biegel, C. M.; Gould, J. M., Kinetics of hydrogen ion diffusion across phospholipid vesicle membranes, *Biochemistry* **20**, 3474–3479 (1981).
410. Perkins, W. R.; Cafiso, D. S., An electrical and structural characterization of  $\text{H}^+/\text{OH}^-$  currents in phospholipid vesicles, *Biochemistry* **25**, 2270–2276 (1986).
411. Meier, E. M.; Schummer, D.; Sandhoff, K., Evidence for the presence of water within the hydrophobic core of membranes, *Chem. Phys. Lipids* **55**, 103–113 (1990).
412. Nagle, J. F., Theory of passive proton conductance in lipid bilayers, *J. Bioenerg. Biomem.* **19**, 413–426 (1987).
413. Gutknecht, J., Proton conductance through phospholipid bilayers: Water wires or weak acids, *J. Bioenerg. Biomem.* **19**, 427–442 (1987).
414. Redelmeier, T. E.; Mayer, L. D.; Wong, K. F.; Bally, M. B.; Cullis, P. R., Proton flux in large unilamellar vesicles in response to membrane potentials and pH gradients, *Biophys. J.* **56**, 385–393 (1989).
415. Norris, F. A.; Powell, G. L., The apparent permeability coefficient for proton flux through phosphatidylcholine vesicles is dependent on the direction of flux, *Biochim. Biophys. Acta* **1030**, 165–171 (1990).
416. Madden, T. D.; Harrigan, P. R.; Tai, L. C. L.; Bally, M. B.; Mayer, L. D.; Redelmeier, T. E.; Loughrey, H. C.; Tilcock, C. P. S.; Reinish, L. W.; Cullis, P. R., The accumulation of drugs within large unilamellar vesicles exhibiting a proton gradient: A survey, *Chem. Phys. Lipids* **53**, 37–46 (1990).

417. Chakraborti, A. C.; Deamer, D. W., Permeability of lipid bilayers to amino acids and phosphate, *Biochim. Biophys. Acta* **1111**, 171–177 (1992).
418. Romanowski, M.; Zhu, X.; Kim, K.; Hruby, V.; O'Brien, D. F., Interactions of enkaphalin peptides with anionic model membranes, *Biochim. Biophys. Acta* **1558**, 45–53 (2002).
419. Lee, R. J.; Wang, S.; Low, P. S., Measurement of endosome pH following folate receptor-mediated endocytosis, *Biochim. Biophys. Acta* **1312**, 237–242 (1996).
420. Boulanger, Y.; Schreier, S.; Leitch, L. C.; Smith, I. C. P., Multiple binding sites for local anesthetics in membranes: Characterization of the sites and their equilibria by deuterium NMR of specifically deuterated procaine and tetracaine, *Can. J. Biochem.* **58**, 986–995 (1980).
421. Boulanger, Y.; Schreier, S.; Smith, I. C. P., Molecular details of anesthetic-lipid interaction as seen by deuterium and phosphorus-31 nuclear magnetic resonance, *Biochemistry* **20**, 6824–6830 (1981).
422. Westman, J.; Boulanger, Y.; Ehrenberg, A.; Smith, I. C. P., Charge and pH dependent drug binding to model membranes: A  $^2\text{H}$ -NMR and light absorption study, *Biochim. Biophys. Acta* **685**, 315–328 (1982).
423. Kelusky, E. C.; Smith, I. C. P., Anesthetic-membrane interaction: A  $^2\text{H}$  nuclear magnetic resonance study of the binding of specifically deuterated tetracaine and procaine to phosphatidylcholine, *Can. J. Biochem. Cell Biol.* **62**, 178–184 (1984).
424. Schreier, S.; Frezzatti, W. A., Jr.; Araujo, P. S.; Chaimovich, H.; Cuccovia, I. M., Effect of lipid membranes on the apparent pK of the local anesthetic tetracaine: Spin label and titration studies, *Biochim. Biophys. Acta* **769**, 231–237 (1984).
425. Trumbore, M.; Chester, D. W.; Moring, J.; Rhodes, D.; Herbette, L. G., Structure and location of amiodarone in a membrane bilayer as determined by molecular mechanics and quantitative X-ray diffraction, *Biophys. J.* **54**, 535–543 (1988).
426. Mason, R. P.; Campbell, S. F.; Wang, S.-D.; Herbette, L. G., Comparison of location and binding for the positively charged 1,4-dihydropyridine calcium channel antagonist amlopidine with uncharged drugs of this class in cardiac membranes, *Molec. Pharmacol.* **36**, 634–640 (1989).
427. Mason, R. P.; Rhodes, D. G.; Herbette, L. G., Reevaluating equilibrium and kinetic binding parameters for lipophilic drugs based on a structural model for drug interaction with biological membranes, *J. Med. Chem.* **34**, 869–877 (1991).
428. Herbette, L. G.; Rhodes, D. R.; Mason, R. P., New approaches to drug design and delivery based on drug-membrane interactions, *Drug Design Deliv.* **7**, 75–118 (1991).
429. Chatelain, P.; Laruel, R., Amiodarone partitioning with phospholipid bilayers and erythrocyte membranes, *J. Pharm. Sci.* **74**, 783–784 (1985).
430. Betageri, G. V.; Rogers, J. A., Thermodynamics of partitioning of  $\beta$ -blockers in the n-octanol buffer and liposome systems, *Int. J. Pharm.* **36**, 165–173 (1987).
431. Choi, Y. W.; Rogers, J. A., The liposome as a model membrane in correlations of partitioning with  $\alpha$ -adrenoreceptor against activities, *Pharm. Res.* **7**, 508–512 (1990).
432. Rogers, J. A.; Choi, Y. W., The liposome partitioning system for correlating biological activities for imidazolidine derivatives, *Pharm. Res.* **10**, 913–917 (1993).
433. Alcorn, C. J.; Simpson, R. J.; Leahy, D. E.; Peters, T. J., Partition and distribution coefficients of solutes and drugs in Brush Border membrane vesicles, *Biochem. Pharmacol.* **45**, 1775–1782 (1993).

434. Smejtek, P.; Wang, S., Distribution of hydrophobic ionizable xenobiotics between water and lipid membrane: Pentachlorophenol and pentachlorophenone. A comparison with octanol-water partition, *Arch. Environ. Contam. Toxicol.* **25**, 394–404 (1993).
435. Pauletti, G. M.; Wunderli-Allenspach, H., Partition coefficients in vitro: Artificial membranes as a standardized distribution model, *Eur. J. Pharm. Sci.* **1**, 273–281 (1994).
436. Krämer, S. D.; Jakits-Dieser, C.; Wunderli-Allenspach, H., Free fatty acids cause pH-dependent changes in drug-lipid membrane interactions around physiological pH, *Pharm. Res.* **14**, 827–832 (1997).
437. Ottiger, C.; Wunderli-Allenspach, H., Partition behavior of acids and bases in a phosphatidylcholine liposome-buffer equilibrium dialysis system, *Eur. J. Pharm. Sci.* **5**, 223–231 (1997).
438. Krämer, S. D.; Braun, A.; Jakits-Dieser, C.; Wunderli-Allenspach, H., Towards the predictability of drug-lipid membrane interactions: The pH-dependent affinity of propranolol to phosphatidylinositol containing liposomes, *Pharm. Rev.* **15**, 739–744 (1998).
439. Xiang, T.-X.; Anderson, B. D., Phospholipid surface density determines the partitioning and permeability of acetic acid in DMPC: cholesterol bilayers, *J. Membrane Biol.* **148**, 157–167 (1995).
440. Xiang, T.-X.; Anderson, B. D., Development of a combined NMR paramagnetic ion-induced line-broadening dynamic light scattering method for permeability measurements across lipid bilayer membranes, *J. Pharm. Sci.* **84**, 1308–1315 (1995).
441. Austin, R. P.; Davis, A. M.; Manners, C. N., Partitioning of ionized molecules between aqueous buffers and phospholipid vesicles, *J. Pharm. Sci.* **54**, 1180–1183 (1995).
442. Austin, R. P.; Barton, P.; Davis, A. D.; Manners, C. N.; Stansfield, M. C., The effect of ionic strength on liposome-buffer and 1-octanol-buffer distribution coefficients, *J. Pharm. Sci.* **87**, 599–607 (1998).
443. Tatulian, S. A., Ionization and ion binding, in Cevc, G. (ed.), *Phospholipid Handbook*, Marcel Dekker, New York, 1993, pp. 511–550.
444. Miller, K. W.; Yu, S.-C. T., The dependence of the lipid bilayer membrane: Buffer partition coefficient of pentobarbitone on pH and lipid composition, *Br. J. Pharm.* **61**, 57–63 (1977).
445. Herbette, L. G., “Pharmacokinetic” and “pharmacodynamic” design of lipophilic drugs based on a structural model for drug interactions with biological membranes, *Pest. Sci.* **35**, 363–368 (1992).
446. Bellemare, F.; Fragata, M., *J. Colloid Interface Sci.* **77**, 243 (1980).
447. Colbow, K.; Chong, C. S., in Colbow, K. (ed.), *Biological Membranes*, Simon Fraser Univ. Burnaby, Canada, 1975, p. 145.
448. Fernández, M. S.; Fromherz, P., Lipoid pH indicators as probes of electrical potential and polarity in micelles, *J. Phys. Chem.* **81**, 1755–1761 (1977).
449. Thomas, J. K., *Chem. Rev.* **80**, 283 (1980).
450. Iwamoto, K.; Sunamoto, J., *Bull. Chem. Soc. Jpn.* **54**, 399 (1981).
451. Laver, D. R.; Smith, J. R.; Coster, H. G. L., *Biochim. Biophys. Acta* **772**, 1 (1984).
452. Kanashina, S.; Kamaya, H.; Ueda, I., *Biochim. Biophys. Acta* **777**, 75 (1984).
453. Lessard, J. G.; Fragata, M., Micropolarities of lipid bilayers and micelles. 3. Effect of monovalent ions on the dielectric constant of the water-membrane interface of unilamellar phosphatidylcholine vesicles, *J. Phys. Chem.* **90**, 811–817 (1986).

454. Kariv, I.; Cao, H.; Oldenburg, K. R., Development of a high-throughput equilibrium dialysis method, *J. Pharm. Sci.* **90**, 580–587 (2001).
455. Lee, A. G., Effects of charged drugs on the phase transition temperature of phospholipid bilayers, *Biochim. Biophys. Acta* **514**, 95–104 (1978).
456. Davis, M. G.; Manners, C. N.; Payling, D. W.; Smith, D. A.; Wilson, C. A., GI absorption for the strongly acidic drug proxicromil, *J. Pharm. Sci.* **73**, 949–953 (1984).
457. Bunton, C. A.; Ohmenzetter, K.; Sepulvida, L., Binding of hydrogen ions to anionic micelles, *J. Phys. Chem.* **81**, 2000–2004 (1977).
458. Garcia-Soto, J.; Fernández, M. S., The effect of neutral and charged micelles on the acid-base dissociation of the local anesthetic tetracaine, *Biochim. Biophys. Acta* **731**, 275–281 (1983).
459. Saad, H. Y.; Higuchi, T., Water solubility of cholesterol, *J. Pharm. Sci.* **54**, 1205–1206 (1965).
460. Mosharraf, M.; Nyström, C., Solubility characterization of practically insoluble drugs using the coulter counter principle, *Int. J. Pharm.* **122**, 57–67 (1995).
461. Quartermain, C. P.; Bonham, N. M.; Irwin, A. K., Improving the odds—high-throughput techniques in new drug selection, *Eur. Pharm. Rev.* **18**, 27–32 (1998).
462. Bevan, C. D.; Lloyd, R. S., A high-throughput screening method for the determination of aqueous drug solubility using laser nephelometry in microtitre plates, *Anal. Chem.* **72**, 1781–1787 (2000).
463. Pan, L.; Ho, Q.; Tsutsui, K.; Takahashi, L., Comparison of chromatographic and spectroscopic methods used to rank compounds for aqueous solubility, *J. Pharm. Sci.* **90**, 521–529 (2001).
464. Hancock, B. C.; Zografi, G., Characterization and significance of the amorphous state in pharmaceutical systems, *J. Pharm. Sci.* **86**, 1–12 (1997).
465. Feitknecht, W.; Schindler, P., *Solubility Constant of Metal Oxides, Metal Hydroxides and Metal Hydroxide Salts in Aqueous Solution*, Butterworths, London, 1963.
466. Brittain, H. G., Spectral methods for the characterization of polymorphs and solvates, *J. Pharm. Sci.* **86**, 405–411 (1997).
467. Levy, R. H.; Rowland, M., Dissociation constants of sparingly soluble substances: Nonlogarithmic linear titration curves, *J. Pharm. Sci.* **60**, 1155–1159 (1971).
468. Charykov, A. K.; Tal'nikova, T.V., pH-metric method of determining the solubility and distribution ratios of some organic compounds in extraction systems, *J. Anal. Chem. USSR* **29**, 818–822 (Engl.) (1974).
469. Kaufman, J. J.; Semo, N. M.; Koski, W. S., Microelectrometric titration measurement of the  $pK_{as}$  and partition and drug distribution coefficients of narcotics and narcotic antagonists and their pH and temperature dependence, *J. Med. Chem.* **18**, 647–655 (1975).
470. Streng, W. H.; Zoglio, M. A., Determination of the ionization constants of compounds which precipitate during potentiometric titration using extrapolation techniques, *J. Pharm. Sci.* **73**, 1410–1414 (1984).
471. Todd, D.; Winnike, R. A., A rapid method for generating pH-solubility profiles for new chemical entities, Abstr. 9th Ann. Mtng., Amer. Assoc. Pharm. Sci., San Diego, 1994.
472. Avdeef, A., pH-metric solubility. 1. Solubility-pH profiles from Bjerrum plots. Gibbs buffer and  $pK_a$  in the solid state, *Pharm. Pharmacol. Commun.* **4**, 165–178 (1998).

473. Avdeef, A.; Berger, C. M.; Brownell, C., pH-metric solubility. 2. Correlation between the acid-base titration and the saturation shake-flask solubility-pH methods, *Pharm. Res.* **17**, 85–89 (2000).
474. Avdeef, A.; Berger, C. M., pH-metric solubility. 3. Dissolution titration template method for solubility determination, *Eur. J. Pharm. Sci.* **14**, 281–291 (2001).
475. Chowhan, Z. T., pH-solubility profiles of organic carboxylic acids and their salts, *J. Pharm. Sci.* **67**, 1257–1260 (1978).
476. Bogardus, J. B.; Blackwood, R. K. Jr., Solubility of doxycycline in aqueous solution, *J. Pharm. Sci.* **68**, 188–194 (1979).
477. Ahmed, B. M.; Jee, R. D., The acidity and solubility constants of tetracyclines in aqueous solution, *Anal. Chim. Acta* **166**, 329–333 (1984).
478. Chiarini, A.; Tartarini, A.; Fini, A., pH-solubility relationship and partition coefficients for some anti-inflammatory arylaliphatic acids, *Arch. Pharm.* **317**, 268–273 (1984).
479. Anderson, B. D., Conradi, R. A., Predictive relationships in the water solubility of salts of a nonsteroidal anti-inflammatory drug, *J. Pharm. Sci.* **74**, 815–820 (1985).
480. Streng, W. H.; Tan, H. G. H., General treatment of pH solubility profiles of weak acids and bases. II. Evaluation of thermodynamic parameters from the temperature dependence of solubility profiles applied to a zwitterionic compound, *Int. J. Pharm.* **25**, 135–145 (1985).
481. Zimmermann, I., Determination of overlapping pK<sub>a</sub> values from solubility data, *Int. J. Pharm.* **31**, 69–74 (1986).
482. Garren, K. W., Pyter, R. A., Aqueous solubility properties of a dibasic peptide-like compound, *Int. J. Pharm.* **63**, 167–172 (1990).
483. Islam, M. H.; Narurkar, M. M., Solubility, stability and ionization behavior of famotidine, *J. Pharm. Pharmacol.* **45**, 682–686 (1993).
484. Streng, W. H.; Yu, D. H.-S., Precision tests of a pH-solubility profile computer program, *Int. J. Pharm.* **164**, 139–145 (1998).
485. Roseman, T. J.; Yalkowsky, S. H., Physical properties of prostaglandin F<sub>2α</sub> (tromethamine salt): Solubility behavior, surface properties, and ionization constants, *J. Pharm. Sci.* **62**, 1680–1685 (1973).
486. Attwood, D.; Gibson, J., Aggregation of antidepressant drugs in aqueous solution, *J. Pharm. Pharmacol.* **30**, 176–180 (1978).
487. Streng, W. H.; Yu, D. H.-S.; Zhu, C., Determination of solution aggregation using solubility, conductivity, calorimetry, and pH measurements, *Int. J. Pharm.* **135**, 43–52 (1996).
488. Zhu, C.; Streng, W. H., Investigation of drug self-association in aqueous solution using calorimetry, conductivity, and osmometry, *Int. J. Pharm.* **130**, 159–168 (1996).
489. Ritschel, W. A.; Alcorn, G. C.; Streng, W. H.; Zoglio, M. A., Cimetidine-theophylline complex formation, *Meth. Find. Exp. Clin. Pharmacol.* **5**, 55–58 (1983).
490. Li, P.; Tabibi, S. E.; Yalkowsky, S. H., Combined effect of complexation and pH on solubilization, *J. Pharm. Sci.* **87**, 1535–1537 (1998).
491. Nuñez, F. A. A.; Yalkowsky, S. H., Solubilization of diazepam, *J. Pharm. Sci. Tech.* 33–36 (1997).
492. Meyer, J. D.; Manning, M. C., Hydrophobic ion pairing: Altering the solubility properties of biomolecules, *Pharm. Res.* **15**, 188–193 (1998).

493. Narisawa, S.; Stella, V. J., Increased shelf-life of fosphenytoin: solubilization of a degradant, phenytoin, through complexation with (SBE)<sub>7m</sub>- $\beta$ -CD, *J. Pharm. Sci.* **87**, 926–930 (1998).
494. Higuchi, T.; Shih, F.-M. L.; Kimura, T.; Rytting, J. H., Solubility determination of barely aqueous soluble organic solids, *J. Pharm. Sci.* **68**, 1267–1272 (1979).
495. Venkatesh, S.; Li, J.; Xu, Y.; Vishnuvajjala, R.; Anderson, B. D., Intrinsic solubility estimation and pH-solubility behavior of cosalane (NSC 658586), and extremely hydrophobic diprotic acid, *Pharm. Res.* **13**, 1453–1459 (1996).
496. Badwan, A. A.; Alkaysi, H. N.; Owais, L. B.; Salem, M. S.; Arafat, T. A., Terfenadine, *Anal. Profiles Drug Subst.* **19**, 627–662 (1990).
497. Streng, W. H.; Hsi, S. K.; Helms, P. E.; Tan, H. G. H., General treatment of pH-solubility profiles of weak acids and bases and the effect of different acids on the solubility of a weak base, *J. Pharm. Sci.* **73**, 1679–1684 (1984).
498. Miyazaki, S.; Oshiba, M.; Nadai, T., Precaution on use of hydrochloride salts in pharmaceutical formulation, *J. Pharm. Sci.* **70**, 594–596 (1981).
499. Ledwidge, M. T.; Corrigan, O. I., Effects of surface active characteristics and solid state forms on the pH solubility profiles of drug-salt systems, *Int. J. Pharm.* **174**, 187–200 (1998).
500. Jinno, J.; Oh, D.-M.; Crison, J. R.; Amidon, G. L., Dissolution of ionizable water-insoluble drugs: The combined effect of pH and surfactant, *J. Pharm. Sci.* **89**, 268–274 (2000).
501. Mithani, S. D.; Bakatselou, V.; TenHoor, C. N.; Dressman, J. B., Estimation of the increase in solubility of drugs as a function of bile salt concentration, *Pharm. Res.* **13**, 163–167 (1996).
502. Anderson, B. D.; Flora, K. P., Preparation of water-soluble compounds through salt formation, in Wermuth, C. G. (ed.), *The Practice of Medicinal Chemistry*, Academic Press, London, 1996, pp. 739–754.
503. Engel, G. L.; Farid, N. A.; Faul, M. M.; Richardson, L. A.; Winneroski, L. L., Salt selection and characterization of LY333531 mesylate monohydrate, *Int. J. Pharm.* **198**, 239–247 (2000).
504. McFarland, J. W.; Avdeef, A.; Berger, C. M.; Raevsky, O. A., Estimating the water solubilities of crystalline compounds from their chemical structure alone, *J. Chem. Inf. Comput. Sci.* **41**, 1355–1359 (2001).
505. McFarland, J. W.; Du, C. M.; Avdeef, A., Factors influencing the water solubility of crystalline drugs, in van de Waterbeemd, H.; Lennernäs, H.; Artursson, P. (eds.), *Drug Bioavailability. Estimation of Solubility, Permeability, Absorption and Bioavailability*, Wiley-VCH, Weinheim, 2003 (in press).
506. Bergström, C. A. S.; Strafford, M.; Lazarova, L.; Avdeef, A.; Luthman, K.; Artursson, P., Absorption classification of oral drugs based on molecular surface properties, *J. Med. Chem.* **46**, 558–570 (2003).
507. Yalkowsky, S. H.; Dannenfelser, R.-M. (eds.), *AQUASOL dATABASE of Aqueous Solubility*, 5th ed., College of Pharmacy, Univ. Arizona, Tucson, 1998.
508. Abraham, M. H.; Le, J., The correlation and prediction of the solubility of compounds in water using an amended solvation energy relationship, *J. Pharm. Sci.* **88**, 868–880 (1999).
509. Faller, B.; Wohnsland, F., Physicochemical parameters as tools in drug discovery and lead optimization, in Testa, B.; van de Waterbeemd, H.; Folkers, G.; Guy, R. (eds.), *Pharmacokinetic Optimization in Drug Research*, Verlag Helvetica Chimica Acta, Zürich and Wiley-VCH, Weinheim, 2001, pp. 257–274.

510. Hidalgo, I. J.; Kato, A.; Borchardt, R. T., Binding of epidermal growth factor by human colon carcinoma cell (Caco-2) monolayers, *Biochem. Biophys. Res. Commun.* **160**, 317–324 (1989).
511. Artursson, P., Epithelial transport of drugs in cell culture. I: A model for studying the passive diffusion of drugs over intestinal absorptive (Caco-2) cells, *J. Pharm. Sci.* **79**, 476–482 (1990).
512. Karlsson, J. P.; Artursson, P., A method for the determination of cellular permeability coefficients and aqueous boundary layer thickness in monolayers of intestinal epithelial (Caco-2) cells grown in permeable filter chambers, *Int. J. Pharm.* **7**, 55–64 (1991).
513. Hilgers, A. R.; Conradi, R. A.; Burton, P. S., Caco-2 cell monolayers as a model for drug transport across the intestinal mucosa, *Pharm. Res.* **7**, 902–910 (1990).
514. Ho, N. F. H.; Raub, T. J.; Burton, P. S.; Barsuhn, C. L.; Adson, A.; Audus, K. L.; Borchardt, R. T., Quantitative approaches to delineate passive transport mechanisms in cell culture monolayers, in Amidon, G. L.; Lee, P. I.; Topp, E. M. (eds.), *Transport Processes in Pharmaceutical Systems*, Marcel Dekker, New York, 2000, pp. 219–317.
515. Hidalgo, I. J.; Hillgren, K. M.; Grass, G. M.; Borchardt, R. T., A new side-by-side diffusion cell for studying transport across epithelial cell monolayers, *In Vitro Cell Dev. Biol.* **28A**, 578–580 (1992).
516. Mueller, P.; Rudin, D. O.; Tien, H. T.; Westcott, W. C., Reconstitution of cell membrane structure in vitro and its transformation into an excitable system, *Nature* **194**, 979–980 (1962).
517. Tanford, C., *Ben Franklin Stilled the Waves: An Informal History of Pouring Oil on Water with Reflections on the Ups and Downs of Scientific Life in General*, Duke Univ. Press: Durham, NC, 1989.
518. Tien, T. H.; Ottova, A. L., The lipid bilayer concept and its experimental realization: From soap bubbles, kitchen sink, to bilayer lipid membranes, *J. Membr. Sci.* **189**, 83–117 (2001).
519. Ottova, A.; Tien, T. H., The 40th anniversary of bilayer lipid membrane research, *Bioelectrochemistry* **5693** (2002).
520. Hooke, R., Royal Society Meeting, in Birch, T. (ed.). *The History of the Royal Society of London*, Vol. 3, No. 29, A., Miller, London, 1672, p. (1757).
521. Newton, I., *Optics*, Dover, New York, 1952 (1704), pp. 215–232 (reprinted).
522. Franklin, B., Of the stilling of waves by means of oil, *Phil. Trans. (Roy. Soc.)* **64**, 445–460 (1974).
523. Tien, H. T.; Ottova, A. L., *Membrane Biophysics: As Viewed from Experimental Bilayer Lipid Membranes (Planar Lipid Bilayers and Spherical Liposomes)*, Elsevier: Amsterdam, 2000.
524. Overton, E., *Vjschr. Naturforsch. Ges. Zurich* **44**, 88 (1899).
525. Collander, R.; Bärlund, H., Permeabilitätsstudien an Chara eratophylla, *Acta Bot. Fenn.* **11**, 72–114 (1932).
526. Langmuir, I., The constitution and fundamental properties of solids and liquids. II. Liquids, *J. Am. Chem. Soc.* **39**, 1848–1906 (1917).
527. Gorter, E.; Grendel, F., On biomolecular layers of lipids on the chromocytes of the blood, *J. Exp. Med.* **41**, 439–443 (1925).
528. Danielli, J. F.; Davson, H., A contribution to the theory of permeability of thin films, *J. Cell Comp. Physiol.* **5**, 495–508 (1935).

529. Singer, S. J.; Nicolson, G. L., The fluid mosaic model of the structure of cell membranes, *Science* **175**, 720–731 (1972).
530. Singer, S. J., The molecular organization of membranes, *Ann. Rev. Biochem.* **43**, 805–834 (1974).
531. Fishman, P. (ed.), in *Proc. Symp. Plasma Membrane*, Am. Heart Assoc. and NY Heart Assoc., New York City, Dec 8–9, 1961, *Circulation* **26**(5) (1962) (suppl.).
532. Mueller, P.; Rudin, D. O.; Tien, H. T.; Wescott, W. C., Reconstitution of cell membrane structure in vitro and its transformation into an excitable system, *J. Phys. Chem.* **67**, 534 (1963).
533. Bangham, A. D., Surrogate cells or Trojan horses, *BioEssays* 1081–1088 (1995).
534. Barry, P. H.; Diamond, J. M., Effects of the unstirred layers on membrane phenomena, *Physiol. Rev.* **64**, 763–872 (1984).
535. Gutknecht, J.; Tosteson, D. C., Diffusion of weak acids across lipid membranes: Effects of chemical reactions in the unstirred layers, *Science* **182**, 1258–1261 (1973).
536. Gutknecht, J.; Bisson, M. A.; Tosteson, F. C., Diffusion of carbon dioxide through lipid bilayer membranes. Effects of carbonic anhydrase, bicarbonate, and unstirred layers, *J. Gen. Physiol.* **69**, 779–794 (1977).
537. Walter, A.; Gutknecht, J., Monocarboxylic acid permeation through lipid bilayer membranes, *J. Membr. Biol.* **77**, 255–264 (1984).
538. Xiang, T.-X.; Anderson, B. D., Substituent contributions to the transport of substituted p-toluic acids across lipid bilayer membranes, *J. Pharm. Sci.* **83**, 1511–1518 (1994).
539. Pohl, P.; Saparov, S. M.; Antonenko, Y. N., The size of the unstirred water layer as a function of the solute diffusion coefficient, *Biophys. J.* **75**, 1403–1409 (1998).
540. Antonenko, Y. N.; Denisov, G. A.; Pohl, P., Weak acid transport across bilayer lipid membrane in the presence of buffers, *Biophys. J.* **64**, 1701–1710 (1993).
541. Cotton, C. U.; Reuss, L., Measurement of the effective thickness of the mucosal unstirred layer in *Necturus* gallbladder epithelium, *J. Gen. Physiol.* **93**, 631–647 (1989).
542. Mountz, J. M.; Tien, H. T., Photoeffects of pigmented lipid membranes in a microporous filter, *Photochem. Photobiol.* **28**, 395–400 (1978).
543. Thompson, M.; Lennox, R. B.; McClelland, R. A., Structure and electrochemical properties of microfiltration filter-lipid membrane systems, *Anal. Chem.* **54**, 76–81 (1982).
544. O'Connell, A. M.; Koeppe, R. E., II; Andersen, O. S., Kinetics of gramicidin channel formation in lipid bilayers: Transmembrane monomer association, *Science* **250**, 1256–1259 (1990).
545. Cools, A. A.; Janssen, L. H. M., Influence of sodium ion-pair formation on transport kinetics of warfarin through octanol-impregnated membranes, *J. Pharm. Pharmacol.* **35**, 689–691 (1983).
546. Camenisch, G.; Folkers, G.; van de Waterbeemd, H., Comparison of passive drug transport through Caco-2 cells and artificial membranes, *Int. J. Pharm.* **147**, 61–70 (1994).
547. Kansy, M.; Senner, F.; Gubernator, K., Physicochemical high throughput screening: Parallel artificial membrane permeability assay in the description of passive absorption processes, *J. Med. Chem.* **41**, 1070–1110 (1998).
548. Ghosh, R., Novel membranes for simulating biological barrier transport, *J. Membr. Sci.* **192**, 145–154 (2001).

549. Zhu, C.; Chen, T.-M.; Hwang, K., A comparative study of parallel artificial membrane permeability assay for passive absorption screening, in CPSA2000: The Symposium on Chemical and Pharmaceutical Structure Analysis. Milestone Development Services. Princeton, NJ, Sept. 26–28, 2000.
550. Kansy, M.; Fischer, H.; Kratzat, K.; Senner, F.; Wagner, B.; Parrilla, I., High-throughput artificial membrane permeability studies in early lead discovery and development, in Testa, B.; van de Waterbeemd, H.; Folkers, G.; Guy, R. (eds.), *Pharmacokinetic Optimization in Drug Research*, Verlag Helvetica Chimica Acta, Zürich and Wiley-VCH: Weinheim, 2001, pp. 447–464.
551. Krishna, G.; Chen, K.-J.; Lin, C.-C.; Nomeir, A. A., Permeability of lipophilic compounds in drug discovery using in-vitro human absorption model, Caco-2, *Int. J. Pharm.* **222**, 77–89 (2001).
552. Adson, A.; Burton, P. S.; Raub, T. J.; Barsuhn, C. L.; Audus, K. L.; Ho, N. F. H., Passive diffusion of weak organic electrolytes across Caco-2 cell monolayers: Uncoupling the contributions of hydrodynamic, transcellular, and paracellular barriers, *J. Pharm. Sci.* **84**, 1197–1204 (1995).
553. Pontier, C.; Pachot, J.; Botham, R.; Lefant, B.; Arnaud, P., HT29-MTX and Caco-2/TC7 monolayers as predictive models for human intestinal absorption: Role of mucus layer, *J. Pharm. Sci.* **90**, 1608–1619 (2001).
554. Wohnsland, F.; Faller, B., High-throughput permeability pH profile and high-throughput alkane/water log P with artificial membranes, *J. Med. Chem.* **44**, 923–930 (2001).
555. Hwang, K., Predictive artificial membrane technology for high throughput screening, in *New Technologies to Increase Drug Candidate Survivability Conf.*, SR Institute, May 17–18, 2001, Somerset, NJ.
556. Avdeef, A.; Strafford, M.; Block, E.; Balogh, M. P.; Chambliss, W.; Khan, I., Drug absorption in vitro model: Filter-immobilized artificial membranes. 2. Studies of the permeability properties of lactones in piper methysticum forst, *Eur. J. Pharm. Sci.* **14**, 271–280 (2001).
557. Ruell, J. A., Membrane-based drug assays. *Modern Drug Disc.* Jan., 28–30 (2003).
558. Ruell, J. A.; Tsinman, K. L.; Avdeef, A., PAMPA—a drug absorption in vitro model. 4. Unstirred water layer in iso-pH mapping assays in  $pK_a$  flux–optimized design (pOD-PAMPA) (submitted).
559. Avdeef, A., PAMPA—a drug absorption in vitro model. 16. Gradient-pH permeability (in preparation).
560. Avdeef, A.; Takács-Novák, K., PAMPA—a drug absorption in vitro model. 23. Permeability of quaternary ammonium drugs and other charged substances (in preparation).
561. Sugano, K.; Hamada, H.; Machida, M.; Ushio, H., High throughput prediction of oral absorption: Improvement of the composition of the lipid solution used in parallel artificial membrane permeability assay, *J. Biomolec. Screen.* **6**, 189–196 (2001).
562. Sugano, K.; Hamada, H.; Machida, M.; Ushio, H.; Saitoh, K.; Terada, K., Optimized conditions of biomimetic artificial membrane permeability assay, *Int. J. Pharm.* **228**, 181–188 (2001).
563. Zhu, C.; Jiang, L.; Chen, T.-M.; Hwang, K.-K., A comparative study of artificial membrane permeability assay for high-throughput profiling of drug absorption potential, *Eur. J. Med. Chem.* **37**, 399–407 (2002).

564. Veber, D. F.; Johnson, S. R.; Cheng, H.-Y; Smith, B. R.; Ward, K. W.; Kopple, K. D., Molecular properties that influence the oral bioavailability of drug candidates, *J. Med. Chem.* **45**, 2615–2623 (2002).
565. www.pampa2002.com.
566. Haase, R. et al., The phospholipid analogue hexadecylphosphatidylcholine inhibits phosphatidyl biosynthesis in Madin-Darby Canine Kidney Cells, *FEBS Lett.* **288**, 129–132 (1991).
567. Rawn, J. D., *Biochemistry*, Niel Patterson Publishers, Burlington, NC, 1989, pp. 223–224.
568. Hennesthal, C.; Steinem, C., Pore-spanning lipid bilayers visualized by scanning force microscopy. *J. Am. Chem. Soc.* **122**, 8085–8086 (2000).
569. Lichtenberg, D., Micelles and liposomes, in Shinitzky, M. (ed.), *Biomembranes: Physical Aspects*, VCH, Weinheim, 1993, pp. 63–96.
570. Lamson, M. J.; Herbette, L. G.; Peters, K. R.; Carson, J. H.; Morgan, F.; Chester, D. C.; Kramer, P. A., Effects of hexagonal phase induction by dolichol on phospholipid membrane permeability and morphology, *Int. J. Pharm.* **105**, 259–272 (1994).
571. Proulx, P., Structure-function relationships in intestinal brush border membranes, *Biochim. Biophys. Acta* **1071**, 255–271 (1991).
572. Krämer, S. D.; Begley, D. J.; Abbott, N. J., Relevance of cell membrane lipid composition to blood-brain barrier function: Lipids and fatty acids of different BBB models, *Am. Assoc. Pharm. Sci. Ann. Mtg.*, 1999.
573. Krämer, S. D.; Hurley, J. A.; Abbott, N. J.; Begley, D. J., Lipids in blood-brain barrier models in vitro I: TLC and HPLC for the analysis of lipid classes and long polyunsaturated fatty acids, *J. Lipid Res.* (2003), in press.
574. Sawada, G. A.; Ho, N. F. H.; Williams, L. R.; Barsuhn, C. L.; Raub, T. J., Transcellular permeability of chlorpromazine demonstrating the roles of protein binding and membrane partitioning, *Pharm. Res.* **11**, 665–673 (1994).
575. Sawada, G. A.; Barsuhn, C. L.; Lutzke, B. S.; Houghton, M. E.; Padbury, G. E.; Ho, N. F. H.; Raub, T. J., Increased lipophilicity and subsequent cell partitioning decrease passive transcellular diffusion of novel, highly lipophilic antioxidants, *J. Pharmacol. Exp. Ther.* **288**, 1317–1326 (1999).
576. Sawada, G. A.; Williams, L. R.; Lutzke, B. S.; Raub, T. J., Novel, highly lipophilic antioxidants readily diffuse across the blood-brain barrier and access intracellular sites, *J. Pharmacol. Exp. Ther.* **288**, 1327–1333 (1999).
577. Helenius, A.; Simons, K., Solubilization of membranes by detergents, *Biochim. Biophys. Acta* **413**, 29–79 (1975).
578. Palm, K.; Luthman, K.; Ros, J.; Gråsjö, J.; Artursson, P., Effect of molecular charge on intestinal epithelial drug transport: pH-dependent transport of cationic drugs, *J. Pharmacol. Exp. Ther.* **291**, 435–443 (1999).
579. Kakemi, K.; Arita, T.; Hori, R.; Konishi, R., Absorption and excretion of drugs. XXX. Absorption of barbituric acid derivatives from rat stomach, *Chem. Pharm. Bull.* **17**, 1534–1539 (1967).
580. Wagner, J. G.; Sedman, A. J., Quantitation of rate of gastrointestinal and buccal absorption of acidic and basic drugs based on extraction theory, *J. Pharmacokinet. Biopharm.* **1**, 23–50 (1973).

581. Garrigues, T. M.; Pérez-Varona, A. T.; Climent, E.; Bermejo, M. V.; Martin-Villodre, A.; Plá-Delfina, J. M., Gastric absorption of acidic xenobiotics in the rat: Biophysical interpretation of an apparently atypical behaviour, *Int. J. Pharm.* **64**, 127–138 (1990).
582. Lee, A. J.; King, J. R.; Barrett, D. A., Percutaneous absorption: A multiple pathway model, *J. Control. Release* **45**, 141–151 (1997).
583. Ho, N. F. H.; Park, J. Y.; Morozowich, W.; Higuchi, W. I., Physical model approach to the design of drugs with improved intestinal absorption, in Roche, E. B. (ed.), *Design of Biopharmaceutical Properties through Prodrugs and Analogs* APhA/APS, Washington, DC, pp. 136–227.
584. Ho, N. F. H.; Park, J. Y.; Ni, P. F.; Higuchi, W. I., Advancing quantitative and mechanistic approaches in interfacing gastrointestinal drug absorption studies in animals and man, in Crouthamel, W. G.; Sarapu, A. (eds.), *Animal Models for Oral Drug Delivery in Man: In Situ and In Vivo Approaches*, APhA/APS, Washington, DC, pp. 27–106.
585. Hansch, C., A quantitative approach to biochemical structure-activity relationships, *Acc. Chem. Res.* **2**, 232–239 (1969).
586. Kubinyi, H., Quantitative structure-activity relationships. IV. Nonlinear dependence of biological activity on hydrophobic character: a new model, *Arzneim. Forsch. (Drug Res.)* **26**, 1991–1997 (1976).
587. Collander, R., The partition of organic compounds between higher alcohols and water, *Acta Chem. Scand.* **5**, 774–780 (1951).
588. van de Waterbeemd, H.; van Boeckel, S.; Jansen, A. C. A.; Gerritsma, K., Transport in QSAR. II: Rate-equilibrium relationships and the interfacial transfer of drugs, *Eur. J. Med. Chem.* **3**, 279–282 (1980).
589. van de Waterbeemd, H.; van Boeckel, S.; de Sevaux, R.; Jansen, A.; Gerritsma, K., Transport in QSAR. IV: The interfacial transfer model. Relationships between partition coefficients and rate constants of drug partitioning, *Pharm. Weekbl. Sci. Ed.* **3**, 224–237 (1981).
590. van de Waterbeemd, H.; Jansen, A., Transport in QSAR. V: Application of the interfacial drug transfer model, *Pharm. Weekbl. Sci. Ed.* **3**, 587–594 (1981).
591. Wils, P.; Warnery, A.; Phung-Ba, V.; Legrain, S.; Scherman, D., High lipophilicity decreases drug transport across intestinal epithelial cells, *J. Pharmacol. Exp. Ther.* **269**, 654–658 (1994).
592. Poulin, P.; Thiel, F.-P., Prediction of pharmacokinetics prior to in vivo studies. 1. Mechanism-based prediction of volume of distribution, *J. Pharm. Sci.* **91**, 129–156 (2002).
593. Vilà, J. I.; Calpena, A. C.; Obach, R.; Domenech, J., Gastric, intestinal and colonic absorption of a series of β-blockers in the rat, *J. Clin. Pharmacol. Ther.* **30**, 280–286 (1992).
594. Cussler, E. L. *Diffusion*, 2nd ed., Cambridge Univ. Press, Cambridge, UK, 1997, p. 112.
595. Chen, Y.; Pant, A. C.; Simon, S. M., P-glycoprotein does not reduce substrate concentration from the extracellular leaflet of the plasma membrane in living cells, *Cancer Res.* **61**, 7763–7768 (2001).
596. Higgins, C. F.; Callaghan, R.; Linton, K. J.; Rosenberg, M. F.; Ford, R. C., Structure of the multidrug resistance P-glycoprotein, *Cancer Biol.* **8**, 135–142 (1997).
597. Regev, R.; Eytan, G. D., Flip-flop of doxorubicin across erythrocyte and lipid membranes, *Biochem. Pharmacol.* **54**, 1151–1158 (1997).

598. Langguth, P.; Kubis, A.; Krumbiegel, G.; Lang, W.; Merkle, H. P.; Wächter, W.; Spahn-Langguth, H.; Weyhenmeyer, R., Intestinal absorption of the quaternary trospium chloride: Permeability-lowering factors and bioavailabilities for oral dosage forms, *Eur. J. Pharm. Biopharm.* **43**, 265–272 (1997).
599. Young, R. C.; Mitchell, R. C.; Brown, T. H.; Ganellin, C. R.; Griffiths, R.; Jones, M.; Rana, K. K.; Saunders, D.; Smith, I. R.; Sore, N. E.; Wilks, T. J., Development of a new physicochemical model for brain penetration and its application to the design of centrally acting H<sub>2</sub> receptor histamine antagonists, *J. Med. Chem.* **31**, 565–671 (1988).
600. Garrone, A.; Marengo, E.; Fornatto, E.; Gasco, A., A study on pK<sub>a</sub><sup>app</sup> and partition coefficient of substituted benzoic acids in SDS anionic micellar system, *Quant. Struct.-Act. Relat.* **11**, 171–175 (1992).
601. Henis, Y. I., Lateral and rotational diffusion in biological membranes, in Shinitzky, M. (ed.), *Biomembranes: Physical Aspects*, VCH, Weinheim, 1993, pp. 279–340.
602. Yazdanian, M.; Glynn, S. L.; Wright, J. L.; Hawi, A., Correlating partitioning and Caco-2 cell permeability of structurally diverse small molecular weight compounds, *Pharm. Res.* **15**, 1490–1494 (1998).
603. Karlsson, J.; Ungell, A.-L.; Grasjo, J.; Artursson, P., Paracellular drug transport across intestinal epithelia: Influence of charge and induced water flux, *Eur. J. Pharm. Sci.* **9**, 47–56 (1999).
604. Irvine, J. D.; Takahashi, L.; Lockhart, K.; Cheong, J.; Tolan, J. W.; Selick, H. E.; Grove, J. R., MDCK (Madin-Darby canine kidney) cells: A tool for membrane permeability screening, *J. Pharm. Sci.* **88**, 28–33 (1999).
605. Adson, A.; Raub, T. J.; Burton, P. S.; Barsuhn, C. L.; Hilgers, A. R.; Audus, K. L.; Ho, N. F. H., Quantitative approaches to delineate paracellular diffusion in cultured epithelial cell monolayers, *J. Pharm. Sci.* **83**, 1529–1536 (1994).
606. Conradi, R. A., Pharmacia & Upjohn, personal correspondence, Feb. 2000.
607. Stenberg, P.; Norinder, U.; Luthman, K.; Artursson, P., Experimental and computational screening models for the prediction of intestinal drug absorption, *J. Med. Chem.* **44**, 1927–1937 (2001).
608. Walter, E.; Janich, S.; Roessler, B. J.; Hilfinger, J. M.; Amidon, G. L., HT29-MTX/Caco-2 cocultures as an in vitro model for the intestinal epithelium: In vitro-in vivo correlation with permeability data from rats and humans, *J. Pharm. Sci.* **85**, 1070–1076 (1996).
609. Hilgendorf, C.; Spahn-Langguth, H.; Regardh, C. G.; Lipka, E.; Amidon, G. L.; Langguth, P., Caco-2 vs Caco-2/HT29-MTX co-cultured cell lines: Permeabilities via diffusion, inside- and outside-directed carrier-mediated transport, *J. Pharm. Sci.* **89**, 63–75 (2000).
610. Aungst, B. J.; Nguyen, N. H.; Bulgarelli, J. P.; Oates-Lenz, K., The influence of donor and reservoir additives on Caco-2 permeability and secretory transport of HIV protease inhibitors and other lipophilic compounds, *Pharm. Res.* **17**, 1175–1180 (2000).
611. Collett, A.; Sims, E.; Walker, D.; He, Y.-L.; Ayrton, J.; Rowland, M.; Warhurst, G., Comparison of HT29-18-C1 and Caco-2 cell lines as models for studying intestinal paracellular drug absorption, *Pharm. Res.* **13**, 216–221 (1996).
612. Partridge, W. M., *Peptide Drug Delivery to the Brain*, Raven Press, New York, 1991, pp. 52–88.

# INDEX

---

- Absorption, 242–246  
Absorption-distribution-metabolism-excretion (ADME)  
medicinal chemists, 4  
multimechanism, 3, 6  
the ‘A’ in, 5
- Aggregates, effect on solubility  
charged, 112  
uncharged, 112
- Amphotericin B, 125
- Apparent partition coefficient,  $\log D$ , 45–53
- Aqueous dilution solubility method, 107
- Bile acids, effects on permeability, 135, 226–228
- Bile salts, 136, 228
- Biological membranes, lipid compositions, 132
- Biopharmaceutics Classification System, 20
- Bjerrum plot, 25–28, 57–58, 103–106, 198
- Black lipid membrane (BLM), 123
- Caco-2, 3, 131, 134, 136, 138, 142, 170, 221, 238, 242
- Charged species transport  
discrimination, 17  
evidence of transport, 215, 218–224
- Chemical sink, 139
- Cosolvent, effect on permeability, 135, 226–228  
Cosolvent solubility method, 108–111  
Cosolvent titrations, 29
- Cyclodextrin  
effect on permeability, 228  
effect on solubility, 113
- Danielli and Davson, 121  
 $\Delta \log P_e$  concept, 222–226  
 $\Delta$  shift method, 113
- diff* 1-2, liposome-water rule, 81–83  
*diff* 3-4, octanol-water rule, 53
- Diffusivity,  $D_{aq}$ , 9, 207
- Dissolution template titration method, 101
- DMSO effect on solubility, 111
- Double-sink PAMPA, 151
- Double-sink  $\Sigma-P_m$  PAMPA GIT model, 244–246, 249
- Effects of cosolvents, bile acids, and surfactants on permeability, 135, 226–228
- Egg lecithin from different sources, 183
- Epithelial cells, 13, 14
- Fick’s law, 7–11
- Focused libraries, 2
- Franklin, Benjamin, experiment, 119

- Gastrointestinal tract (GIT), 11–17  
 food effect, 12, 13  
 pH, 12  
 surface area, 12, 13  
 transit time, 13
- Glycocalyx, 14, 15
- Gorter and Grendel, 120
- Gradient-pH  
 equilibrium concentrations, 151  
 permeability, 211–215, 241
- Henderson-Hasselbalch equation, 23
- Human absorption prediction, 242–246
- Human jejunal permeability, 237–239, 246
- Intrinsic permeability, 11, 201
- Intrinsic permeability vs. alkane partition coefficient, 124
- Ion-pair absorption  
 liposome, experimental methods, 75  
 octanol, fact or fiction, 53
- Ion-pair partition, 45
- Langmuir, I., 120
- Lateral diffusion, 16, 17
- Lipid bilayer concept, history, 118–121
- Lipophilicity profiles  
 liposome-water, 80  
 octanol-water, 47–52
- Liposomes  
 charged, 85  
 partition coefficients, 67–90. *See also* Partition coefficients
- $\log D$ , apparent partition coefficient, 45–53
- $\log S$  vs. pH plots, 92–97
- Maximum chemical diversity, 1
- Membrane retention, %R, 117, 142, 169–171, 196
- Membrane permeability, 199, 205
- Metallothionein, 28
- Microfilters as artificial membrane supports, 124
- Microvilli, 15
- Mucus layer, 14, 15
- Octanol  
 partition coefficients, 42–66. *See also* Partition coefficients  
 structure, 19
- Octanol-impregnated filters, 126–128
- Olive oil, 43, 119, 167
- Overton, 119
- Parallel artificial membrane permeability assay (PAMPA), 116–249  
 brush-border lipid (Chugai model), 130, 181  
 dodecane, 168  
 DOPC, 166  
 effect of buffer, 229–231  
 egg lecithin (Roche model), 128  
 five-component lipid model, 181  
 hexadecane (Novartis model), 129  
 hydrophilic filters (Aventis model), 131  
 intra- and inter-plate reproducibility, 232–234  
 membrane retention %R, 117, 142, 169–171, 196  
 models for gastrointestinal tract, 156–160  
 neutral lipid models, 160–169  
 olive oil, 167  
 optimized double-sink model, 236–238  
 retention-gradient-sink (*p*ION model), 131  
 two-component anionic lipid models, 171–181  
 UV spectral data quality, 233–235
- PAMPA lipid models table, 157, 240
- Partition coefficients  
 charged liposome, 85–87  
 entropy-/enthalpy-driven, 70  
 immobilized artificial membrane (IAM)  
 chromatography, 54, 83
- liposomes  
 charged compounds, 79–83, 85  
 databases, 69  
 dielectric medium, 71–73  
 drug location in bilayer, 69  
 Gouy-Chapman, 81–82  
 one-pH measurement, 84  
 preparation, 74  
 prediction of absorption, 90  
 prediction from octanol, 76–79  
 table of values, 87–89  
 water wires, 73
- liposome-water, 67–90
- octanol  
 Bjerrum plot, 57, 58  
 charged-species partitioning, 53  
 chromatographic, 54  
 common-ion effect, 52  
 conditional constants, 45  
 cyclic voltammetry, 55  
 databases, 45  
 high-throughput, 59  
 $\log D$ , 45–50  
 microconstants, 54  
 peptides, 50  
 potentiometric, 55–58  
 table of values, 59–66
- octanol-water, 42–66

- Permeability, 9–11, 116–249  
 effect of stirring, 205, 231  
 gastrointestinal tract and blood-brain barrier, 116–118  
 mass balance, 143  
 Permeability equation  
 Caco-2, 142  
 gradient pH, 148–150  
 iso-pH with membrane retention, 142–147  
 precipitate in donor compartment, 147  
 sink condition in acceptor well, 147  
 without retention, 139–142  
 Permeability-lipophilicity, nonlinear relations, 153–156  
 Permeability-pH, 132–134  
 Permeability-solubility product, 10, 11  
 pH electrode calibration, 27  
 pH microclimate, 17, 133, 249  
 pH partition hypothesis, 7  
 pH, intracellular, 18  
 $pK_a$  (ionization constant)  
 Bjerrum plot, 25–26, 28  
 capillary electrophoresis, 32  
 chromatographic, 33  
 databases, 24  
 equilibria and quotients, 23  
 ionic strength dependence, 24  
 microconstants, 33  
 poorly-soluble compounds, 29  
 potentiometric, 25  
 spectroscopic, 31  
 table of values, 35–41  
 $pK_a^{\text{flux}}$ , 202, 206, 216  
 $pK_a^{\text{gibbs}}$ , 93, 98  
 $pK_a^{\text{mem}}$ , 67, 83–84  
 $pK_a^{\text{oct}}$ , 44, 83–84  
 Pockels, Agnes, 119  
 Salt solubility, 97–98  
 Screening, target vs. ADME, 2  
 $sdiff$  3–4 rule, 97, 99  
 Serum proteins, role in permeability assays, 135  
 Sialic acid, 15  
 Sink condition, 9, 138–139, 147, 150, 171, 177, 196, 228, 235  
 Solubility, 10–11, 91–115  
 databases, 100  
 equilibria and quotients, 92–93  
 fast UV plate spectrophotometer method, 107  
 high-throughput, 107–111  
 HPLC method, 101  
 potentiometric method, 101–107  
 saturation shake-flask method, 101  
 table of values, dissolution template method, 114–115  
 turbidity method, 101  
 use of Bjerrum plots, 103–107  
 Solubility product, 97  
 Soy lecithin, 187–196  
 Stirring in PAMPA, 231–232  
 Structure of phospholipid membranes, 131  
 Summary of simple prediction rules, 247–249  
 Surface ion-pair (SIP), 82  
 Surfactants, effect on permeability, 135, 226–228  
 Tetrad of equilibria  
 liposome partition coefficients, 67  
 octanol partition coefficients, 43  
 solubility, 99  
 Tight junction pores 15, 18, 248  
 Unstirred water layer (UWL), 15, 135, 177, 179, 199–208, 216, 236, 241  
 thickness, 199, 205, 207–208, 236  
 UV spectra, 235  
 Vancomycin, 28  
 Villi, 13  
 Yasuda-Shedlovsky procedure, 29