

Definition of Standards and Fundamental and Derived Physical Constants*

APPENDIX A

The definition of primary standards is by agreement within the International Conference of Weights and Measures whose most recent general meeting was October, 1964 in Paris. Measured and derived values of the fundamental physical constants summarize hundreds of physical measurements made over the years by scientists in all parts of the world. They have been subjected to exhaustive statistical analysis and, with their accompanying error limits (which are given as three standard deviations), represent the best values to date (1963). For most problem work three significant figures suffice, and the "computational" (rounded) values may be used. The data presented are based largely on values given in the *National Bureau of Standards Technical News Bulletin*, Vol. 47, No. 10 (October 1963).

*See "A Pilgrim's Progress in Search of the Fundamental Constants," by J. W. M. Du Mond, *Physics Today*, October 1965.

DEFINITION OF STANDARDS AND EQUIVALENTS

Standard	Abbre- viation	Equivalent
Meter	m	1,650,763.73 wavelengths in vacuo of the unperturbed transition $2p_{10} - 5d_5$ in Kr^{86}
Kilogram	kg	mass of the international kilogram at Sèvres, France
Second	sec	9,192,631,770 vibrations of the unperturbed hyperfine transition $4,0 - 3,0$ of the fundamental state 2S_1 in Cs^{133s}
Degree Kelvin	$^{\circ}K$	defined in the thermodynamic scale by assigning $273.16\ ^{\circ}K$ to the triple point of water
Unified atomic mass unit	amu	$\frac{1}{12}$ the mass of an atom of the C^{12} nuclide
Mole	mol	amount of substance containing the same number of atoms as 12 gm (exactly) of pure C^{12}
Standard acceleration of free fall	g _n	9.80665 meter/sec ²
Normal atmospheric pressure	atm	101,325 nt/meter ²
Thermochemical calorie	cal	4.1840 joules
Liter	l	0.001,000,028 meter ³
Inch	in.	0.0254 meter
Pound (avdp.)	lb	0.453,592,37 kg

* There is no measurable difference between this and the previous standard of time, 1/31,556,925.9747 of the tropical year at 12^h ET, 0 January 1900. For this reason and because even more accurate maser standards may soon be available, the Cs standard was adopted provisionally rather than "permanently."

FUNDAMENTAL AND DERIVED CONSTANTS

Name	Symbol	Computational Value	Best Experimental Value ^b
Speed of light	c	3.00×10^8 meters/sec	2.997925 ± 0.000003
Permeability constant	μ_0	1.26×10^{-6} henry/meter	$4\pi \times 10^{-7}$ exactly
Permittivity constant	ϵ_0	8.85×10^{-12} farad/meter	8.85418 ± 0.00002
Elementary charge	e	1.60×10^{-19} coul	1.60210 ± 0.00007
Avogadro constant	N_A	6.02×10^{23} /mole	6.02252 ± 0.00028
Electron rest mass	m_e	9.11×10^{-31} kg	9.1091 ± 0.0004
Proton rest mass	m_p	1.67×10^{-27} kg	1.67252 ± 0.00008
Neutron rest mass	m_n	1.67×10^{-27} kg	1.67482 ± 0.00008
Faraday constant	F	9.65×10^4 coul/mole	9.64870 ± 0.00016
Planck constant	h	6.63×10^{-34} joule sec	6.6256 ± 0.0005
Fine structure constant	α	7.30×10^{-3}	7.29720 ± 0.00010
Electron charge/mass ratio	e/m_e	1.76×10^{11} coul/kg	1.758796 ± 0.000019
Quantum/charge ratio	\hbar/e	4.14×10^{-16} joule sec/coul	4.13556 ± 0.00012
Electron Compton wavelength	λ_C	2.43×10^{-12} meter	2.42621 ± 0.00006
Proton Compton wavelength	λ_{Cp}	1.32×10^{-15} meter	1.32140 ± 0.00004
Rydberg constant	R_∞	1.10×10^7 /meter	1.0973731 ± 0.0000003
Bohr radius	a_0	5.29×10^{-11} meter	5.29167 ± 0.00007
Bohr magneton	μ_B	9.27×10^{-24} joule/tesla ^a	9.2732 ± 0.0006
Nuclear magneton	μ_N	5.05×10^{-27} joule/tesla ^a	5.0505 ± 0.0004
Proton magnetic moment	μ_p	1.41×10^{-26} joule/tesla ^a	1.41040 ± 0.00013
Universal gas constant	R	8.31 jouls/ $^\circ$ K mole	8.3143 ± 0.0012
Standard volume of ideal gas	—	2.24×10^{-2} meter ³ /mole	2.24136 ± 0.00030
Boltzmann constant	k	1.38×10^{-23} joule/ $^\circ$ K	1.38054 ± 0.00018
First radiation constant $\tau = 2 hc^2$	c_1	3.74×10^{-16} watt/meter ²	3.7405 ± 0.0003
Second radiation constant hc/k	c_2	1.44×10^{-2} meter $^\circ$ K	1.43379 ± 0.00019
Wien displacement constant	b	2.90×10^{-3} meter $^\circ$ K	2.8978 ± 0.0004
Stefan-Boltzmann constant	σ	5.67×10^{-8} watt/meter ² $^\circ$ K ⁴	5.6697 ± 0.0029
Gravitational constant	G	6.67×10^{-11} nt meter ² /kg ²	6.670 ± 0.015

^a Tesla = weber/meter².^b Same units and power of ten as the computational value.

Miscellaneous Terrestrial Data

APPENDIX B

Standard atmosphere	1.013×10^5 nt/meter ² 14.70 lb/in ² 760.0 mm-Hg
Density of dry air at STP*	1.293 kg/meter ³ 2.458×10^{-3} slug/ft ³
Speed of sound in dry air at STP	331.4 meters/sec 1089 ft/sec 742.5 miles/hr
Acceleration of gravity, <i>g</i> (standard value) ^b	9.80665 meters/sec ² 32.1740 ft/sec ²
Solar constant ^c	1340 watts/m ² 1.92 cal/cm ² -min
Mean total solar radiation	3.92×10^{26} watts
Equatorial radius of earth	6.378×10^6 meters 3963 miles
Polar radius of earth	6.357×10^6 meters 3950 miles
Volume of earth	1.087×10^{21} meter ³ 3.838×10^{22} ft ³
Radius of sphere having same volume	6.371×10^6 meters 3959 miles 2.090×10^7 ft

Mean density of earth	5522 kg/meter ³
Mass of earth	5.983×10^{24} kg
Mean orbital speed of earth	29,770 meters/sec 18.50 miles/sec
Mean angular speed of rotation of earth	7.29×10^{-5} radians/sec
Earth's magnetic field, B (at Washington, D. C.)	5.7×10^{-5} tesla ^d
Earth's magnetic dipole moment	6.4×10^{21} amp-m ²

* STP = standard temperature and pressure = 0° C and 1 atm.

^b This value, used for barometer corrections, legal weights, etc., was adopted by the International Committee on Weights and Measures in 1901. It approximates 45° latitude at sea level.

^c The solar constant is the solar energy falling per unit time at normal incidence on unit area of the earth's surface.

^d Tesla = weber/meter².

The Solar System^c

APPENDIX C

Planet	Mercury ♀	Venus ♀	Earth ⊕, Θ, δ	Mars ♂	Jupiter ♀	Saturn ♀	Uranus ♀	Neptune ♀	Pluto ♀
Mean diameter km.	5,000	12,400	12,742	6,870	139,760	115,100	51,000	50,000	12,700 ^f
Earth diameters	0.39	0.973	1.000	0.532	10.97	9.03	4.00	3.90	0.46
Volume (earth volumes)	0.06	0.92	1.00	0.15	1,318	736	64	39	0.10
Mass (earth masses)	0.04	0.82	1.00	0.11	318.3	95.3	14.7	17.3	1.07
Density (earth densities)	0.69	0.89	1.00	0.70	0.24	0.13	0.23	0.29	?
Mean density gm./cm. ³	3.8	4.86	5.52	3.96	1.33	0.71	1.26	1.6	?
Surface gravity (earth's)	0.27	0.86	1.00	0.37	2.64	1.17	0.92	1.44	?
Velocity of escape, km/sec	3.6	10.2	11.2	5.0	60	36	21	23	11?
Length of day (earth days)	58.6 ^d	30 ^d ? ^e	1 ^d	1 ^d 37 ^m 23 ^s	9 ^h 55 ^m	10 ^h 38 ^m	10.7 ^a	15.8 ^a	?

Planet	Mercury ♀	Venus ♀	Earth, ⊕, θ, δ	Mars σ	Jupiter 2	Saturn ♄	Uranus ♆, ♋	Neptune ♌	Pluto ♎
Period, sidereal days	87.97	224.70	365.26	686.98	4,332.59	10,759.20	30,685.93	60,187.64	90,885
Inclination of equator to orbit	—	0°?	23°27'	25°12'	3°7'	26°45'	98.0°	29°	?
Oblateness	0.00	0.00	1/296	1/192	1/15.4	1/9.5	1/14	1/45	?
Atmosphere, main constituents	none	N ₂ , CO ₂ , A	N ₂ , O ₂	N ₂ , CO ₂ , H ₂ O	CH ₄ , NH ₃	none			
Maximum surface temperature, °K	700	700	350	320	153	138	110?	90?	80?
Distance from Sun, 10 ⁶ km	58	108	149	228	778	1426	2869	4495	5900

The Sun ○ 329,390 earth masses, mean density 1.42, mean diameter 1,390,600 km, surface gravity 28 (earth's).

The Moon ☽ 0.01228 earth masses, mean density 3.36, mean diameter 3,476 km, surface gravity 0.17 (earth's), distance from earth 38×10^4 km.

* Adapted from Payne-Gaposchkin and *Handbook of Chemistry and Physics*.

Periodic Table of the Elements

APPENDIX D

Atomic weights are expressed in **atomic mass units** (amu), one atom of the isotope C¹² being defined to have a mass of (exactly) 12 amu. For unstable elements the mass number of the most stable or best known isotope is given in brackets.

Group →	Period Series	I	II	III	IV	V	VI	VII	VIII	0
1	1	1 H 1.00797	3 Li 6.939	4 Be 9.0122	5 B 10.811	6 C 12.01115	7 N 14.0067	8 O 15.9994	9 F 18.9984	2 He 4.0026
2	2	11 Na 22.9898	12 Mg 24.312	13 Al 26.9815	14 Si 28.086	15 P 30.9738	16 S 32.064	17 Cl 35.453	18 A 39.948	10 Ne 20.183
3	3	19 K 39.102	20 Ca 40.08	21 Sc 44.956	22 Ti 47.90	23 V 50.942	24 Cr 51.996	25 Mn 54.9380	26 Fe 55.847	27 Co 58.9332
4	4	29 Cu 63.54	30 Zn 65.37	31 Ga 69.72	32 Ge 72.59	33 As 74.9216	34 Se 78.96	35 Br 79.909	36 Kr 83.80	10 Ne 20.183
5	5	37 Rb 85.47	38 Sr 87.62	39 Y 88.905	40 Zr 91.22	41 Nb 92.906	42 Mo 95.94	43 Tc [99]	44 Ru 101.07	45 Rh 102.905
6	6	47 Ag 107.870	48 Cd 112.40	49 In 114.82	50 Sn 118.69	51 Sb 121.75	52 Te 127.60	53 I 126.9044	46 Pd 106.4	54 Xe 131.30
7	7	55 Cs 132.905	56 Ba 137.34	57-71 Lanthanide series*	72 Hf 178.49	73 Ta 180.948	74 W 183.85	75 Re 186.2	76 Os 190.2	77 Ir 192.2
8	8	79 Au 196.967	80 Hg 200.59	81 Tl 204.37	82 Pb 207.19	83 Bi 208.980	84 Po [210]	85 At [210]	78 Pt 195.09	86 Ra [222]
9	9	87 Fr [223]	88 Ra [226]	89 Actinide series ^b						
10	10									

* Lanthanide series:

57 La 138.91	58 Ce 140.12	59 Pr 140.907	60 Nd 144.24	61 Pm 145.1	62 Sm 150.35	63 Eu 151.96	64 Gd 157.26	65 Tb 165.924	66 Dy 162.50	67 Ho 164.930	68 Er 167.26	69 Tm 168.934	70 Yb 173.04	71 Lu 174.97
89 Ac [232]	90 Th [231]	91 Pa [233]	92 U [233]	93 Np [237]	94 Pu [242]	95 Am [243]	96 Cm [245]	97 Bk [249]	98 Cf [249]	99 Es [254]	100 Fm [252]	101 Md [256]	102 Lw [257]	

^b Actinide series:

- $$9. \frac{d}{dx} \cosh x = \sinh x \quad 9. \int \sinh x \, dx = \cosh x$$
- $$10. \frac{d}{dx} \arctan x = \frac{1}{1+x^2} \quad 10. \int \frac{dx}{1+x^2} = \arctan x$$
- $$11. \frac{d}{dx} \arcsin x = \frac{1}{\sqrt{1-x^2}} \quad 11. \int \frac{dx}{\sqrt{1-x^2}} = \arcsin x$$
- $$12. \frac{d}{dx} \text{arcsec } x = \frac{1}{x\sqrt{x^2-1}} \quad 12. \int \frac{dx}{x\sqrt{x^2-1}} = \text{arcsec } x$$
- $$13. \frac{d}{dx} \cos x = -\sin x \quad 13. \int \sin x \, dx = -\cos x$$
- $$14. \frac{d}{dx} \sin x = \cos x \quad 14. \int \cos x \, dx = \sin x$$
- $$15. \frac{d}{dx} \tan x = \sec^2 x \quad 19. \int \tan x \, dx = \ln |\sec x|$$
- $$16. \frac{d}{dx} \cot x = -\csc^2 x \quad 20. \int \cot x \, dx = \ln |\sin x|$$
- $$17. \frac{d}{dx} \sec x = \tan x \sec x \quad 21. \int \sec x \, dx = \ln |\sec x + \tan x|$$
- $$18. \frac{d}{dx} \csc x = -\cot x \csc x \quad 22. \int \csc x \, dx = \ln |\csc x - \cot x|$$

Vector products

Let $\mathbf{i}, \mathbf{j}, \mathbf{k}$ be unit vectors in the x, y, z directions. Then

$$\mathbf{i} \cdot \mathbf{i} = \mathbf{j} \cdot \mathbf{j} = \mathbf{k} \cdot \mathbf{k} = 1, \quad \mathbf{i} \cdot \mathbf{j} = \mathbf{j} \cdot \mathbf{k} = \mathbf{k} \cdot \mathbf{i} = 0,$$

$$\mathbf{i} \times \mathbf{i} = \mathbf{j} \times \mathbf{j} = \mathbf{k} \times \mathbf{k} = 0,$$

$$\mathbf{i} \times \mathbf{j} = \mathbf{k}, \quad \mathbf{j} \times \mathbf{k} = \mathbf{i}, \quad \mathbf{k} \times \mathbf{i} = \mathbf{j}.$$

Any vector \mathbf{a} with components a_x, a_y, a_z along the x, y, z axes can be written

$$\mathbf{a} = a_x \mathbf{i} + a_y \mathbf{j} + a_z \mathbf{k}.$$

Let $\mathbf{a}, \mathbf{b}, \mathbf{c}$ be arbitrary vectors with magnitudes a, b, c . Then

$$\mathbf{a} \times (\mathbf{b} + \mathbf{c}) = \mathbf{a} \times \mathbf{b} + \mathbf{a} \times \mathbf{c}$$

$$(\mathbf{s}\mathbf{a}) \times \mathbf{b} = \mathbf{a} \times (\mathbf{s}\mathbf{b}) = \mathbf{s}(\mathbf{a} \times \mathbf{b}) \quad (\mathbf{s} \text{ a scalar}).$$

Let θ be the smaller of the two angles between \mathbf{a} and \mathbf{b} . Then

$$\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a} = a_x b_x + a_y b_y + a_z b_z = ab \cos \theta$$

$$\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{vmatrix} = (a_y b_z - b_y a_z) \mathbf{i} + (a_z b_x - b_z a_x) \mathbf{j} + (a_x b_y - b_x a_y) \mathbf{k}$$

$$|\mathbf{a} \times \mathbf{b}| = ab \sin \theta$$

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \mathbf{b} \cdot (\mathbf{c} \times \mathbf{a}) = \mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})$$

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \cdot \mathbf{c})\mathbf{b} - (\mathbf{a} \cdot \mathbf{b})\mathbf{c}$$

Values of Trigonometric Functions

APPENDIX J

TRIGONOMETRIC FUNCTIONS

Radians	Degrees	Sines	Cosines	Tangents	Cotangents		
.0000	0	.0000	1.0000	.0000	∞	90	1.5708
.0175	1	.0175	.9998	.0175	57.29	89	1.5533
.0349	2	.0349	.9994	.0349	28.64	88	1.5359
.0524	3	.0523	.9986	.0524	19.08	87	1.5184
.0698	4	.0698	.9976	.0699	14.30	86	1.5010
.0873	5	.0872	.9962	.0875	11.430	85	1.4835
.1047	6	.1045	.9945	.1051	9.514	84	1.4661
.1222	7	.1219	.9925	.1228	8.144	83	1.4486
.1396	8	.1392	.9903	.1405	7.115	82	1.4312
.1571	9	.1564	.9877	.1584	6.314	81	1.4137
.1745	10	.1736	.9848	.1763	5.671	80	1.3963
.1920	11	.1908	.9816	.1944	5.145	79	1.3788
.2094	12	.2079	.9781	.2126	4.705	78	1.3614
.2269	13	.2250	.9744	.2309	4.332	77	1.3439
.2443	14	.2419	.9703	.2493	4.011	76	1.3265
.2618	15	.2588	.9659	.2679	3.732	75	1.3090
.2793	16	.2756	.9613	.2867	3.487	74	1.2915
.2967	17	.2924	.9563	.3057	3.271	73	1.2741
.3142	18	.3090	.9511	.3249	3.078	72	1.2566
.3316	19	.3256	.9455	.3443	2.904	71	1.2392

TRIGONOMETRIC FUNCTIONS (*Continued*)

Radians	Degrees	Sines	Cosines	Tangents	Cotangents		
.3491	20	.3420	.9397	.3640	2.748	70	1.2217
.3665	21	.3584	.9336	.3839	2.605	69	1.2043
.3840	22	.3746	.9272	.4040	2.475	68	1.1868
.4014	23	.3907	.9205	.4245	2.356	67	1.1694
.4189	24	.4067	.9135	.4452	2.246	66	1.1519
.4363	25	.4226	.9063	.4663	2.144	65	1.1345
.4538	26	.4384	.8988	.4877	2.050	64	1.1170
.4712	27	.4540	.8910	.5095	1.963	63	1.0996
.4887	28	.4695	.8829	.5317	1.881	62	1.0821
.5061	29	.4848	.8746	.5543	1.804	61	1.0647
.5236	30	.5000	.8660	.5774	1.732	60	1.0472
.5411	31	.5150	.8572	.6009	1.664	59	1.0297
.5585	32	.5299	.8480	.6249	1.600	58	1.0123
.5760	33	.5446	.8387	.6494	1.540	57	0.9948
.5934	34	.5592	.8290	.6745	1.483	56	0.9774
.6109	35	.5736	.8192	.7002	1.428	55	0.9599
.6283	36	.5878	.8090	.7265	1.376	54	0.9425
.6458	37	.6018	.7986	.7536	1.327	53	0.9250
.6632	38	.6157	.7880	.7813	1.280	52	0.9076
.6807	39	.6293	.7771	.8098	1.235	51	0.8901
.6981	40	.6428	.7660	.8391	1.192	50	0.8727
.7156	41	.6561	.7547	.8693	1.150	49	0.8552
.7330	42	.6691	.7431	.9004	1.111	48	0.8378
.7505	43	.6820	.7314	.9325	1.072	47	0.8203
.7679	44	.6947	.7193	.9657	1.036	46	0.8029
.7854	45	.7071	.7071	1.0000	1.000	45	0.7854
		Cosines	Sines	Cotangents	Tangents	Degrees	Radians

Nobel Prize Winners in Physics^a

APPENDIX K

1901	Wilhelm Konrad Röntgen	1845-1923	German	Discovery of X-rays.
1902	Hendrik Antoon Lorentz	1853-1928	Dutch	Influence of magnetism on the phenomena of atomic radiation.
	Pieter Zeeman	1865-1943	Dutch	
1903	Henri Becquerel	1852-1908	French	Discovery of natural radioactivity and of the radioactive elements radium and polonium.
	Pierre Curie	1850-1906	French	
	Marie Curie	1867-1934	French	
1904	Baron Rayleigh	1842-1919	English	Discovery of argon.
1905	Philipp Lenard	1862-1947	German	Research in cathode rays.
1906	Sir Joseph John Thomson	1856-1940	English	Conduction of electricity through gases.
1907	Albert A. Michelson	1852-1931	U. S.	Invention of interferometer and spectroscopic and metrological investigations.
1908	Gabriel Lippmann	1845-1921	French	Photographic reproduction of colors.
1909	Guglielmo Marconi	1874-1937	Italian	Development of wireless telegraphy.
	Karl Ferdinand Braun	1850-1918	German	
1910	Johannes Diderik van der Waals	1837-1923	Dutch	Equations of state of gases and fluids.
1911	Wilhelm Wien	1864-1928	German	Laws of heat radiation.
1912	Nils Gustaf Dalen	1869-1937	Swedish	Automatic coastal lighting.
1913	Heike Kamerlingh-Onnes	1853-1926	Dutch	Properties of matter at low temperatures; production of liquid helium.
1914	Max von Laue	1879-1960	German	Diffraction of X-rays in crystals.
1915	Sir William Henry Bragg	1862-1942	English	Study of crystal structure by means of X-rays.
	Sir William Lawrence Bragg	1890-	English—his son	
1916	(No award)			
1917	Charles Glover Barkla	1877-1944	English	Discovery of the characteristic X-rays of elements.
1918	Max Planck	1858-1947	German	Discovery of the elemental quantum.
1919	Johannes Stark	1874-1957	German	Discovery of the Doppler effect in canal rays and the splitting of spectral lines in the electric field.

APPENDIX K

1920	Charles Edouard Guillaume	1861-1938	Swiss	Discovery of the anomalies of nickel-steel alloys.
1921	Albert Einstein	1879-1955	German	Discovery of the law of the photoelectric effect.
1922	Niels Bohr	1885-1963	Danish	Study of structure and radiations of atoms.
1923	Robert Andrews Millikan	1868-1953	U. S.	Work on elementary electric charge and the photoelectric effect.
1924	Manne Siegbahn	1886-	Swedish	Discoveries in the area of X-ray spectra.
1925	James Franck	1882-1964	German	Laws governing collision between electron and atom.
	Gustav Hertz	1887	German	
1926	Jean Perrin	1870-1942	French	Discovery of the equilibrium of sedimentation.
1927	Arthur H. Compton	1892-1962	U. S.	Discovery of the scattering of X-rays by charged particles
	Charles T. R. Wilson	1869-1959	English	Invention of the cloud chamber, a device to make visible the paths of charged particles.
1928	Sir Owen Williams Richardson	1879-1959	English	Discovery of the law known by his name (the dependency of the emission of electrons on temperature).
1929	Louis-Victor de Broglie	1892-	French	Wave nature of electrons.
1930	Sir Chandrasekhara Raman	1888-	Indian	Work on the scattering of light and discovery of the effect known by his name.
1931	(No award)			
1932	Werner Heisenberg	1901-	German	Creation of quantum mechanics.
1933	Paul Adrien Maurice Dirac	1902-	English	Discovery of new fertile forms of the atomic theory.
	Erwin Schrödinger	1887-1961	Austrian	
1934	(No award)			
1935	James Chadwick	1891-	English	Discovery of the neutron.
1936	Victor Hess	1883-1964	Austrian	Discovery of cosmic radiation.
	Carl David Anderson	1905-	U. S.	Discovery of the positron.
1937	Clinton Joseph Davission	1881-1958	U. S.	Discovery of diffraction of electrons by crystals.
	George P. Thomson	1892-	English	
1938	Enrico Fermi	1901-1954	Italian	Artificial radioactive elements from neutron irradiation.
1939	E. O. Lawrence	1901-1958	U. S.	Invention of the cyclotron.
1940-				
1942	(No awards)			
1943	Otto Stern	1888	U. S. ^a	Work with molecular beams and magnetic moment of proton.
1944	Isidor Isaac Rabi	1898-	U. S.	Nuclear magnetic resonance.
1945	Wolfgang Pauli	1900-1958	Austrian	Discovery of quantum exclusion principle.
1946	Percy Williams Bridgman	1882-1961	U. S.	High-pressure physics.
1947	Sir Edward Appleton	1892-	English	Upper atmosphere physics and discovery of Appleton layer.
1948	Patrick Maynard Stuart Blackett	1897-	English	Discoveries in cosmic radiation and nuclear physics.
1949	Hideki Yukawa	1907-	Japanese	Prediction of existence of meson.
1950	Cecil Frank Powell	1903-	English	Photographic method of studying nuclear processes; discoveries about mesons.
1951	Sir John Douglas Cockcroft	1897-	English	Transmutation of atomic nuclei by artificially accelerated atomic particles.
	Ernest Thomas Sinton Walton	1903-	Irish	
1952	Felix Bloch	1905-	U. S.	Measure of magnetic fields in atomic nuclei.
	Edward Mills Purcell	1912-	U. S.	
1953	Frits Zernike	1888-	Dutch	Invention of phase contrast microscopy.
1954	Max Born	1882-	English ^b	Work in quantum mechanics and statistical interpretation of wave function.

	Walther Bothe	1891-1957	German	Analysis of cosmic radiation using the coincidence method.
1955	Willis E. Lamb, Jr.	1913	U. S.	Fine structure of hydrogen.
	Polykarp Kusch	1911-	U. S.	Magnetic moment of electron.
1956	John Bardeen	1908-	U. S.	Invention and development of transistor.
	Walter H. Brattain	1902-	U. S.	
	William B. Shockley	1910-	U. S.*	
1957	Chen Ning Yang	1922-	Chinese ^d	Non-conservation of parity and work in elementary particle theory.
	Tsung Dao Lee	1926-	Chinese ^d	
1958	Pavel A. Čerenkov	1904-	Russian	Discovery and interpretation of Čerenkov effect of radiation by fast charged particles in matter.
	Ilya M. Frank	1908-	Russian	
	Igor Y. Tamm	1895-	Russian	Discovery of the antiproton.
1959	Owen Chamberlain	1920-	U. S.	
	Emilio Gino Segré	1905-	U. S.*	
1960	Donald A. Glaser	1926-	U. S.	Invention of bubble chamber.
1961	Robert L. Hofstadter	1915-	U. S.	Electromagnetic structure of nucleons from high-energy electron scattering.
	Rudolf L. Mössbauer	1929	German	Discovery of recoilless resonance absorption of gamma rays in nuclei.
1962	Lev D. Landau	1908-	Russian	Theory of condensed matter; phenomena of superfluidity and superconductivity.
1963	Eugene B. Wigner	1902-	U. S. ^f	Contributions to theoretical atomic and nuclear physics.
	Maria Goeppert-Mayer	1906-	U. S. ^e	Shell model theory and magic numbers for the atomic nucleus.
	J. H. D. Jensen	1907-	German	
1964	C. H. Townes	1915-	U. S.	Invention of the maser and theory of coherent atomic radiation.
	Nikolai Basov	1922-	Russian	
	Aleksandr Prokhorov	1916-	Russian	
1965	Richard Feynman	1918	U. S.	Development of quantum electrodynamics
	Julian Schwinger	1918	U. S.	
	Shin-Ichiro Tomonaga	1906	Japanese	

* See Nobel: *The Man and His Prizes*, by Schück et al., Elsevier, N. Y.

^b Born in Germany; naturalized British citizen.

^c Born in England; naturalized U. S. citizen.

^d Both have permanent U. S. resident status.

^e Born in Italy; naturalized U. S. citizen.

^f Born in Hungary; naturalized U. S. citizen.

^g Born in Germany; naturalized U. S. citizen.

Quantity	Rationalized mks	Gaussian
Permittivity constant	ϵ_0	$1/4\pi$
Permeability constant	μ_0	$4\pi/c^2$
Electric displacement	D	$D/4\pi$
Magnetic induction	B	B/c
Magnetic flux	Φ_B	Φ_B/c
Magnetic field strength	H	$cH/4\pi$
Magnetization	M	cM
Magnetic dipole moment	μ	$c\mu$

The Gaussian System of Units

APPENDIX L

Much of the literature of physics is written, and continues to be written, in the Gaussian system of units. In electromagnetism many equations have slightly different forms depending on whether it is intended, as in this book, that mks variables be used or that Gaussian variables be used. Equations in this book can be cast in Gaussian form by replacing the symbols listed below under "rationalized mks" by those listed under "Gaussian." For example, Eq. 37-26,

$$\mathbf{B} = \mu_0(\mathbf{H} + \mathbf{M})$$

becomes

$$\frac{\mathbf{B}}{c} = \left(\frac{4\pi}{c^2} \right) \left(\frac{c}{4\pi} \mathbf{H} + c\mathbf{M} \right)$$

or

$$\mathbf{B} = \mathbf{H} + 4\pi\mathbf{M}$$

in Gaussian form. Symbols used in this book that are not listed below remain unchanged. The quantity c is the speed of light.

Quantity	Rationalized mks	Gaussian
Permittivity constant	ϵ_0	$1/4\pi$
Permeability constant	μ_0	$4\pi/c^2$
Electric displacement	D	$D/4\pi$
Magnetic induction	B	B/c
Magnetic flux	Φ_B	Φ_B/c
Magnetic field strength	H	$cH/4\pi$
Magnetization	M	cM
Magnetic dipole moment	μ	$c\mu$

In addition to casting the equations in the proper form it is of course necessary to use a consistent set of units in those equations. Below we list some equivalent quantities in mks and Gaussian units. This table can be used to transform units from one system to the other.

Quantity	Symbol	Mks system	Gaussian system
Length	l	1 meter	10^2 cm
Mass	m	1 kg	10^3 gm
Time	t	1 sec	1 sec
Force	F	1 newton	10^5 dynes
Work or Energy	W, E	1 joule	10^7 ergs
Power	P	1 watt	10^7 ergs/sec
Charge	q	1 coulomb	3×10^9 statcoul
Current	i	1 ampere	3×10^9 statamp
Electric field strength	E	1 volt/meter	$\frac{1}{3} \times 10^{-4}$ statvolt/cm
Electric potential	V	1 volt	$\frac{1}{300}$ statvolt
Electric polarization	P	1 coul/meter ²	3×10^5 statcoul/cm ²
Electric displacement	D	1 coul/meter ²	$12\pi \times 10^5$ statvolt/cm
Resistance	R	1 ohm	$\frac{1}{3} \times 10^{-11}$ sec cm ⁻¹
Capacitance	C	1 farad	9×10^{11} cm
Magnetic flux	Φ_B	1 weber	10^8 maxwells
Magnetic induction	B	1 tesla = 1 weber/meter ²	10^4 gauss
Magnetic field strength	H	1 amp-turn/meter	$4\pi \times 10^{-3}$ oersted
Magnetization	M	1 weber/meter ²	$1/4\pi \times 10^4$ gauss
Inductance	I	1 henry	$\frac{1}{3} \times 10^{-11}$

All factors of 3 in the above table, apart from exponents, should be replaced by (2.997925 ± 0.000003) for accurate work; this arises from the numerical value of the speed of light. For example the mks unit of capacitance ($= 1$ farad) is actually 8.98758×10^{11} cm rather than $9 (= 3^2) \times 10^{11}$ cm as listed above. This example also shows that not only units but also the dimensions of physical quantities may differ between the two systems. In the mks system (see Appendix F) the dimensions of capacitance are $M^{-1}L^{-2}T^2Q^2$; in the Gaussian system they are simply L , the Gaussian standard unit of capacitance being 1 cm.

The student should consult *Classical Electromagnetism*, p. 611, by J. D. Jackson (John Wiley and Sons, 1962) for a fuller treatment of units and dimensions.

Answers to Odd-Numbered Problems

Chapter 1

1. $6.00 \text{ ft} = 1.83 \text{ meters}$.
 3. 186 miles.
 5. (a) $d_{\text{sun}}/d_{\text{moon}} = 400$.
(b) $V_{\text{sun}}/V_{\text{moon}} = 6.4 \times 10^9$.
We assume spherical shapes and precise eclipsing.
 7. 2.03 hr.
 9. (a) $\sim 10^6$ meters (see Table 1-1).
(b) ~ 10 sec (see Table 1-2).
 11. C, D, A, B, E (best to worse). The criteria are first the constancy and second the magnitude of the daily variation.
-
13. $r_z = 2.0 \text{ miles}; r_y = r_z = 4.0 \text{ miles}$.
 15. (a) 21 ft.
(b) Can be greater but not less.
(c) $10i + 12j + 14k$ for a particular choice of axes.
 17. 6950 miles, pointing through the earth from Washington to Manila.
 21. (a) Scalar of magnitude 30 units 2 .
(b) Vector of magnitude 52 units 2 perpendicular to the plane formed by \mathbf{a} and \mathbf{b} .
 31. (a) $d_x = d_y = a^2; d_z = -a^2$.
(b) $\mathbf{b} \cdot \mathbf{c} = a^2; \mathbf{d} \cdot \mathbf{c} = \mathbf{d} \cdot \mathbf{b} = 0$.

Chapter 2

3. The displacements should be:
(a) parallel, (b) antiparallel,
(c) perpendicular.
5. The magnitudes are: 5, 10, 11.2, 11.2, 11.2. The angles made with the x -axis are: 323° , 53.1° , 26.5° , 79.7° , and 260° .
7. 81.0 miles; 39.5° N of E.
9. (a) $a_x = -2.8 \text{ meters}$;
 $a_y = -2.8 \text{ meters}$.
 $b_x = 5.0 \text{ meters}; b_y = 0$.
 $c_x = 3.0 \text{ meters}; c_y = 5.2 \text{ meters}$.
(b) $d_x = 5.17 \text{ meters}$;
 $d_y = 2.37 \text{ meters}$.

Chapter 3

1. (a) 5.7 ft/sec
(b) 7.0 ft/sec.
3. (a) Infinite number.
(b) 60 miles.
5. 3000 ft/sec^2 , upwards.
7. $OA AB BC CD$
 $v_x + 0 + +$
 $a_x - 0 + 0$
Intervals OA and BC
9. $8.0 \times 10^{14} \text{ meters/sec}^2$.
11. 10 cm; no time.
13. (a) 5.0 ft/sec^2 .
(b) 4.0 sec.

- (c) 6.0 sec.
 (d) 90 ft.
15. (a) 15 ft/sec.
 (b) 5.0 ft/sec.
 (c) 23 ft.
17. No.
19. (a) 57 ft/sec.
 (b) 3.6 sec.
21. 3.41 sec; 187 ft.
23. (a) 3.5×10^5 ft.
 (b) 330 sec.
25. 40 ft/sec.
27. $\frac{1}{8}$ ft.
29. (a) 17 sec.
 (b) 290 meters.
31. (a) $a: LT^{-2}$; ft/sec².
 b: LT^{-3} ; ft/sec³.
 (b) 2.0 sec.
 (c) 24 ft.
 (d) -16 ft.
 (e) 3.0, 0.0, -9.0, -24 ft/sec.
 (f) 0.0, -6.0, -12, -18 ft/sec.²
- Chapter 4**
3. The third step.
7. 76°.
9. 1.9 in.
11. Yes.
13. (a) 2.0 mm.
 (b) $v_{\text{hor}} = 1.0 \times 10^9$ cm/sec.
 $v_{\text{vert}} = -0.20 \times 10^9$ cm/sec.
15. Electron 5.5×10^{-15} meter.
 Neutron 1.0×10^{-6} meter.
 Neon 1.4×10^{-5} meter.
 Oxygen 2.3×10^{-5} meter.
 Golf ball 4.9×10^{-4} meter (assuming 100 meters/sec).
19. (a) $v_0 \cos \theta_0$.
 (b) g.
 (c) $\mathbf{v} \perp \mathbf{a}$.
 (d) $(v_0^2/g) \cos^2 \theta_0$.
21. 6.7×10^6 meters/sec.
23. 2400 meters/sec².
25. (a) 4.2 meters at 45°;
 5.5 meters at 68°;
 6.0 meters at 90°.
 (b) 4.2 meters at 135°.
 (c) 0.85 meter/sec at 135°.
 (d) 0.94 meter/sec at 90°;
 0.94 meter/sec at 180°.
 (e) 0.27 meter/sec² at 225°.
 (f) 0.30 meter/sec² at 180°;
 0.30 meter/sec² at 270°.
 All angles are measured counter-clockwise from a line extending horizontally to the right from O in Fig. 4-15.
27. 6.0×10^{-3} meter/sec².
29. (a) $\mathbf{r} = i(r \sin \omega t) + j(r \cos \omega t)$
31. $\mathbf{u}_r = i \cos \theta + j \sin \theta$.
 $\mathbf{u}_{\theta} = -i \sin \theta + j \cos \theta$.
33. 2.2 meters/sec and 1.8 meters/sec.
35. (a) He should head the boat 25.4° upstream.
 (b) 12.7 min.
37. (a) Wind is blowing from a direction 75° E of S.
 (b) 30° E of N.
- Substituting W for E in the above yields another solution.
- Chapter 5**
1. $a_1/a_2 = m_2/m_1$.
3. 1.0 meters/sec², 37° from \mathbf{F}_2 toward \mathbf{F}_1 .
5. 1300 lb, 5.5 sec, 50 ft, 2.7 sec.
7. $a_{\text{hor}} = 65$ ft/sec², $a_{\text{down}} = 32$ ft/sec².
 $v_{\text{hor}} = 65t$ ft/sec²,
 $v_{\text{down}} = 32t$ ft/sec².
9. (a) 740 nt.
 (b) 610 nt.
 (c) Zero.
H is mass is 75 kg at each location.
13. Lower it with an acceleration of 4.2 ft/sec² or greater.
15. (a) 3.2 ft/sec².
 (b) 58 lb.
17. (a) 2.0 meters/sec².
 (b) 4.0 meters/sec.
 (c) 4.0 meters.
19. (a) $g \sin \theta$ down the plane.
 (b) $g \sin \theta$ down the plane.
 (c) $(g - a) \sin \theta$ down the plane.
 (d) $(g + a) \sin \theta$ down the plane.
 (e) Zero.
21. (a) 0.50 slugs.
 (b) 20 lb.
23. (a) 32 lb, 55 lb.
 (b) 16 ft/sec².
25. 3.3 meters/sec², 6.5 nt.
27. (a) 19 ft/sec².
 (b) $(M + m) \times 19$ ft/sec².
- Chapter 6**
1. 110 lb.
3. (a) 0.38 ft/sec², 0.79 ft/sec².
 (b) 0.041, 0.028.
5. (a) 0.031 lb.
 (b) 0.13.
7. 0.75.

9. 40 lb.
 11. (a) 15 lb.
 (b) 6.4 ft/sec^2 .
 13. (a) $v_0^2/4g \sin \varphi$.
 (b) No.
 15. (a) 1.06 nt (in tension).
 (b) 3.62 meters/sec².
 (c) 1.06 nt (in compression),
 3.62 meters/sec².
 17. $g(\sin \theta - \sqrt{2} \mu_k \cos \theta)$.
 19. (a) 2700 ft.
 (b) 5000 lb, upward.
 21. (a) 0.0338 nt.
 (b) 9.77 nt.
 23. (a) 15° .
 (b) 0.27.
 25. $v^2/r = Mg/m$.
 27. (a) 68 ft.
 (b) 18° .
29. $p_{\min} = \frac{1}{2\pi} \sqrt{\frac{g}{r}} \left(\frac{\sin \theta - \mu \cos \theta}{\cos \theta + \mu \sin \theta} \right)$
 $p_{\max} = \frac{1}{2\pi} \sqrt{\frac{g}{r}} \left(\frac{\sin \theta + \mu \cos \theta}{\cos \theta - \mu \sin \theta} \right)$

Chapter 7

1. (a) 52 lb.
 (b) -260 ft-lb.
 (c) +300 ft-lb.
 (d) -40 ft-lb.
 (e) Zero.
 (f) Zero.
 3. (a) 50 lb.
 (b) No.
 (c) Yes; 100 ft-lb.
 (d) No.
 5. (a) Zero.
 (b) 30.1 joules.
 (c) -30.1 joules.
 (d) 0.225.
 9. Boy: 4.8 meters/sec.
 Man: 2.4 meters/sec.
 11. (a) 2.9×10^7 meters/sec.
 (b) 1.3×10^6 ev.
 15. (a) 135 nt.
 (b) 60.0 joules.
 17. 18 ft-lb.
 21. (a) 5.4×10^{10} ft-lb.
 (b) 1.6×10^6 hp.
 23. (a) 1.8×10^6 ft-lb.
 (b) 0.55 hp.
 25. 0.27 hp.
 27. (b) $m(v_f/t_f)^2 t$.
 (c) 140 hp.

Chapter 8

5. (a) $v = 2 \sqrt{gl}$, $T = 5 mg$.
 (b) 71° .
 7. 2d.
 9. (a) $v_B = v_0$; $v_C = \sqrt{v_0^2 + gh}$.
 (b) $(v_0^2 + 2gh)/2L$.
 (c) It will never reach B.
 11. (a) $\sqrt{5gR}$.
 (b) P is above the horizontal by $\sin^{-1}(\frac{1}{3})$.
 13. (a) 4.0 meters.
 (b) 4.5 meters/sec.
 15. (a) $U(x) = -km_1 m_2/x$ if $U(\infty) = 0$.
 (b) $\frac{km_1 m_2 d}{x_1(d+x_1)}$.
 17. (a) $F_x = -kx$, $F_y = -ky$; F always points radially inward.
 (b) $F_r = -kr$, $F_\theta = 0$.
 21. (a) 31.0 joules.
 (b) 5.33 meters/sec.
 (c) Conservative.
 23. (a) $\sqrt{2gl(\sin \theta - \mu \cos \theta)}$.
 (b) $l(\sin \theta - \mu \cos \theta)/\mu$.
 25. 7.2 meters/sec.
 27. (a) 260 ft-lb.
 (b) 45 ft-lb; 1.7 ft up the plane.
 29. (a) 6700 ft-lb/sec.
 (b) 2000 ft-lb.
 (c) No.
 31. (a) 5.8×10^{-13} joule.
 (b) 0.08.
 33. (a) $\cong 0.010$ kg; 0.023 kg.
 35. 27 Mev.
- Chapter 9
3. 6.46×10^{-11} meter, along the line of symmetry.
 5. 6.75×10^{-13} meter below, on the line of symmetry.
 7. (a) Center of mass remains at rest.
 (b) 0.75 meter.
 9. 5500 slug-ft/sec; 6.0 miles/hr; 13 miles/hr.
 11. 13.6 ft.
 13. $10\sqrt{2}$ meters/sec, 135° from either.
 15. 1.1×10^6 ft.
 17. (a) The casing and the capsule both move forward with speeds of 24,000 and 27,000 ft/sec respectively.
 (b) The energy increases from 9.38×10^9 ft-lb before separation to 9.40×10^9 ft-lb after, the in-

crease coming from energy stored in the spring.

19. $\frac{wv_{rel}}{W+w}$.

21. 220 bullets/min.

25. 5100 lb, 5600 hp.

Chapter 10

1. 2.5 meters/sec.

3. 8.8 meters/sec.

5. $2mv/t$.

7. Slows down to 3.0 meters/sec.

9. $m_1/3$.

11. 310 meters/sec.

13. Block, 4.0 ft/sec; ball, 8.0 ft/sec.

15. (a) 4.1 ft/sec; 2.4×10^3 joules.

$$(b) v_{12} = 3.3 \text{ ft/sec}; v_{24} = 5.3 \text{ ft/sec}.$$

17. (a) The left mass comes to rest; the center mass moves to the right with a speed $v_0(m - M)/(m + M)$; the right mass moves to the right with a speed $2v_0m/(m + M)$.

(b) The left mass moves to the left with a speed $v_0(M - m)/(m + M)$; the center mass comes to rest; the right mass moves to the right with a speed $2v_0m/(m + M)$.

19. 0.25 meter.

21. 12 lb.

23. $v_0 = \left(2E \frac{M+m}{Mm} \right)^{\frac{1}{2}}$.

25. (a) $\frac{1}{2}m_1v_{1i}^2$.

$$(b) \frac{1}{2} \frac{m_1^2v_{1i}^2}{(m_1 + m_2)}$$
.

$$(c) m_2/(m_1 + m_2).$$

$$(d) \frac{1}{2}m_1(v_{1i}^2 + v_{1i}v_{cm} + v_{cm}^2) + \frac{1}{2}m_2v_{cm}^2; \text{ zero; } 100\%; \text{ no.}$$

27. 1.9 meters/sec, 30° to initial direction; no.

29. 117° from final direction of B ; no.

31. (a) 5.0×10^8 cm/sec.

(b) 6.9×10^8 cm/sec, at 14° to the original direction opposite to the neutron and in the plane defined by the helium nucleus and the neutron.

33. $\pi(r_1 + r_2)^2$

35. 4.2×10^{11} transmutations/meter² sec.

37. 8.12 Mev.

Chapter 11

1. (a) 1200 in./min.

(b) 600 in./min.

3. (a) 3.8×10^6 radians/sec.

(b) 190 meters/sec.

5. (a) -0.27 radians/sec².

(b) 20 rev.

7. $0.80\omega_0$.

9. (a) $x^2 + y^2 = R^2$; a circle of radius R ; ω is the angular speed of the body.

(b) $v_x = \omega y$; $v_y = -\omega x$; $v = \omega R$; v is tangential to the circle.

(c) $a_x = -\omega^2 x$; $a_y = -\omega^2 y$;
 $a = \omega^2 R$;
a points radially inward.

11. (a) 2.0×10^{-7} radians/sec;
 3.0×10^4 meters/sec.

(b) 6.0 meters/sec².

13. 5.7.

15. (a) 70 radians/sec.

(b) -13 radians/sec².

(c) 240 ft.

17. 0.12 radian/sec.

Chapter 12

1. (a) $i(yF_z - zF_y) + j(zF_x - xF_z) + k(xF_y - yF_x)$.

13. 6.75×10^{12} radians/sec.

15. (a) 2.6×10^{29} joules.

(b) 2.4×10^9 years.

17. 292 ft-lb.

19. (a) -7.66 radians/sec².

(b) -11.7 newton meter.

(c) 4.58×10^4 joules.

(d) 624 revs.

(e) (c) can be computed and has the same value.

21. 19 ft/sec²; 7.3 lb; 6.7 lb.

23. $g/2$.

25. (a) 11 ft.

(b) 1.4 sec.

27. (a) $K = mg(R - r)$; $\frac{2}{7}$ translational;
 $\frac{5}{7}$ rotational.

(b) $\frac{17}{7}mg$.

29. $50mg$

33. (a) Mg .

(b) $MR^2\omega^2/4$.

(c) $R^2\omega^2/4g$.

37. Axis is $5\sqrt{3}$ ft above ground and 5 ft from the wall.

39. 5.4 meters/sec.

Chapter 13

1. 2.0 rad/sec; clockwise as seen from above.
9. $\frac{1}{2}ab\omega M$; L precesses around the axis of rotation, making an angle $\theta = \tan^{-1} a/b - \tan^{-1} b/a$ with it.
11. (a) $L_{\text{spin}}/L_{\text{orbital}} = \frac{2}{3}(R_m/R_{e-m})^2$ in which R_m is the lunar radius and R_{e-m} is the earth-moon distance.
- (b) Increase or decrease by one-half of present value.
13. 0.77 rad/sec.
15. (a) Linear momentum, angular momentum, and mechanical energy.
- (b) $\frac{Ml^2}{12d^2 + l^2}$, where l is the length of the stick.
19. 250 rev/min.

$$21. \frac{v_1}{1 + (I/MR^2)}$$

$$25. \sqrt{2gr/\cos \theta_0}$$

27. (a) They rotate about the center of mass (the center of the pole) with $\omega = 6.7$ rad/sec.
- (b) As above, but $\omega = 60$ rad/sec.
- (c) $K_a = 5.0 \times 10^3$ joules;
 $K_b = 45 \times 10^3$ joules;
 difference represents work done by the skaters.

Chapter 14

$$1. \frac{W[h(2r - k)]^{\frac{1}{2}}}{r - h}$$

$$3. 74.4 \text{ gm.}$$

5. Along a line extending from the center of the hole through the center of the disk, beyond the latter point by a distance $Rr^2/2(R^2 - r^2)$.

$$9. 7.2 \text{ ft.}$$

$$13. \text{Back: } 880 \text{ lb; front: } 630 \text{ lb.}$$

$$15. F_A = 120 \text{ lb}; F_B = 72 \text{ lb}; T = 47 \text{ lb.}$$

$$17. F_h = 5.0 \text{ lb}; F_v = 30 \text{ lb}; d = 1.0 \text{ ft.}$$

19. (a) $W/2 \sin \theta$, tangent to the chain.
 (b) $\frac{1}{2}W \cot \theta$.

Chapter 15

1. 0.28 sec.
3. (a) 99 nt.
 (b) 99 nt/meter.
5. (a) 4.0 sec.
 (b) $\pi/2$ radians/sec.
 (c) 0.37 cm.
 (d) $0.37 \cos(\pi t/2)$, in cm.

$$(c) -0.58 \sin(\pi t/2), \text{ in cm/sec.}$$

$$(f) 0.58 \text{ cm/sec.}$$

$$(g) 0.91 \text{ cm/sec}^2.$$

$$(h) \text{Zero.}$$

$$(i) 0.58 \text{ cm/sec.}$$

$$7. 3.1 \text{ cm.}$$

$$9. k_1 = k(1 + n)/n.$$

$$k_2 = k(1 + n).$$

$$13. (a) 1.6 \times 10^4 \text{ meters/sec}^2,$$

$$2.5 \text{ meters/sec.}$$

$$(b) 2.2 \text{ meters/sec.}$$

$$7.9 \times 10^3 \text{ meters/sec}^2.$$

$$15. (a) 6.2 \text{ in.}$$

- (b) 1.2%; most of the original energy appears as internal energy in the block.

$$17. 19 \text{ lb.}$$

$$19. \frac{3}{4}, \frac{1}{4}; A/\sqrt{2}.$$

$$23. 9.5 \text{ meters/sec}^2.$$

$$27. \frac{1}{2\pi} \sqrt{\frac{(g^2 + r^4/R^2)^{\frac{1}{2}}}{l}}$$

31. (a) 0.45 cycles/sec. (Does it matter whether the nail is very smooth or rusty?)

$$(b) 4.0 \text{ ft.}$$

$$33. (a) 39 \text{ radians/sec.}$$

$$(b) 34 \text{ radians/sec.}$$

$$(c) 120 \text{ radians/sec}^2.$$

37. (a) O₂: 8.0 amu; HCl: 0.97 amu; CO: 6.8 amu.

$$(b) 500 \text{ nt/meter.}$$

$$41. K_{\text{trans}} = 6.3 \times 10^{-2} \text{ joule.}$$

$$K_{\text{rot}} = 3.1 \times 10^{-2} \text{ joule.}$$

$$43. (b) \sqrt{k/m}.$$

Chapter 16

$$1. 1600 \text{ miles.}$$

$$3. 4.8 \text{ sec.}$$

$$7. (b) 3.2 \text{ meters.}$$

$$9. (b) 84.2 \text{ min.}$$

$$(c) \text{No,}$$

$$11. (a) G(M_1 + M_2)m/a^2.$$

$$(b) GM_1m/b^2$$

$$(c) \text{Zero.}$$

$$13. (a) 2.6 \times 10^4 \text{ ft/sec.}$$

$$(b) 87 \text{ min.}$$

$$15. 2.5 \times 10^4 \text{ km.}$$

$$17. (a) \frac{1}{4}.$$

$$(b) \frac{1}{2}.$$

$$(c) B, \text{ by } 8.5 \times 10^7 \text{ ft-lb.}$$

$$19. (a) -GmM_e/r.$$

$$(b) -2GmM_e/r.$$

$$(c) \text{Falls directly down.}$$

$$21. 1.88 \text{ years.}$$

23. (b) 1.0×10^4 meters/sec.
 (c) Moon: 1.9×10^3 meters/sec.
 Sun: 6.2×10^6 meters/sec.
25. (a) $2\pi \sqrt{d^3/3mG}$.
 (b) 2.
 (c) 2.
27. $\sqrt{GM/L}$.
29. $a/3$.
31. (a) 2.2×10^{-6} nt/kg, \perp to line joining centers.
 (b) -5.3×10^{-7} joules/kg.
33. (a) $-Gm \left(\frac{M_e}{R} + \frac{M_\mu}{r} \right)$.
 (b) Nine-tenths of the way to the moon and at infinity.
 (c) Earth: $U = 6.3 \times 10^7$ joules.
 $g = 9.8$ nt/kg.
 Moon: $U = 3.9 \times 10^6$ joules
 $g = 1.6$ nt/kg

Chapter 17

1. (a) 240 lb/in.²
 (b) 2.3 lb/in.²
3. (b) 6000 lb.
5. (a) $\frac{1}{2}\rho g D^2 W$, $\frac{1}{6}\rho g D^3 W$.
 (b) $D/3$ up from bottom.
9. $\frac{1}{2}A(h_2 - h_1)^2$.
11. (a) fA/a .
 (b) 20 lb.
13. 0.20 ft³.
15. 0.67 gm/cm³, 0.74 gm/cm³.
17. 0.19, no.
19. $0.12 \left(\frac{1}{\rho} - \frac{1}{8} \right)$, ρ in gm/cm³.
21. (b) $p = \rho gh$ where h is the vertical depth below the surface.

Chapter 18

1. 29 ft/sec.
3. 1.1×10^5 ft-lb.
7. $v = 4.1$ meters/sec.
 $v' = 21$ meters/sec.
 $Av = 8.1 \times 10^{-3}$ meter³/sec.
11. (a) $2 \sqrt{h(H-h)}$.
 (b) Yes, a distance h above the bottom.
15. 790 lb; 250 lb (up).
17. 410 meters/sec.

Chapter 19

3. (a) 10 cm, 1.0 vib/sec; 200 cm/sec.
 200 cm.
 (b) 63 cm/sec.

5. (a) 12 cm.
 (b) 180° .
7. 130 meters/sec.
9. v_0 .
11. $1/4\pi$ watts/meter².
13. Intensity proportional to r^{-1} ; amplitude proportional to $r^{-1/2}$.
15. $\lambda = 2 \sqrt{\frac{4(H+h)^2 + d^2}{2 \sqrt{4H^2 + d^2}}}$.
19. (b) Even though the displacement of the string is zero at this instant the transverse velocities are not so that energy is present as kinetic energy.

21. $y = 6 \cos \frac{\pi}{2} (0.005x + 8.00t - 0.57)$.

Chapter 20

1. 17 meters, 0.017 meters.
3. 1.0×10^5 vib/sec.
7. (a) $\frac{l(V-v)}{Vv}$.
 (b) 1600 ft.
11. 3.6×10^{-8} meter.
13. (a) 5.0×10^8 vib/sec.
 (b) $SBD/SAD = \frac{1}{4}$.
15. 31 and 94 vib/sec.
21. 1130, 1500, and 1880 cycles/sec.
23. $L(r-1)r$; 0.13 meter; 0.27 meter.
25. (a) 323 vib/sec.
 (b) 6.
27. 387 vib/sec.
29. No.
31. (a) 970 vib/sec.
 (b) 1030 vib/sec.
 (c) Zero.
33. (a) 0.90 ft.
 (b) 1440 vib/sec.
 (c) 1080 ft/sec.
 (d) 0.75 ft.
35. (a) 42° .
 (b) 20 sec.
37. 990 meters/sec.

Chapter 21

1. 373.15/273.16.
3. Materials, shape, absolute temperature, air currents; dimensions are T^{-1} .
5. -40° ; 575° .
7. 10,000 ohms, $4.124 \times 10^{-3}/^\circ\text{C}$,
 $-1.780 \times 10^{-6}/^\circ\text{C}^2$.
9. 1.002 in.
11. The clock will run about 9 sec slow.

13. 46.4 cm.
 19. $+28.9 \text{ cm}^3$.
 23. (a) $1.4 \times 10^{-2} \text{ kg meter}^2/\text{sec}$;
 0.41 joule.
 (b) $\Delta\omega/\omega = \Delta K/K = -0.32\%$;
 $\Delta L/L = 0$.
 25. $7.44 \times 10^{-4}/\text{C}^\circ$.

Chapter 22

1. 1.17°C .
 3. 190 watts.
 5. $0.13 \text{ Btu/lb F}^\circ$.
 9. $0.59 \text{ cal/gm C}^\circ$.
 11. Mean specific heat exceeds that at
 the midpoint by $B \frac{t^2}{12}$.
 13. (a) 34 Btu.
 (b) 270 F° .
 15. (a) $500 \text{ C}^\circ/\text{meter}$.
 (b) 4.6 cal/sec.
 (c) 75°C .
 17. 8.6°C (Cu-Al) and 57°C (Al-brass).
 19. $1.4 \times 10^{-5} \text{ kcal/meter}^2 \text{ sec}$;
 $6.2 \times 10^{14} \text{ kcal}$.
 21. $0.42 \text{ cal/meter sec C}^\circ$.
 23. (a) 6 cal.
 (b) -43 cal.
 (c) 40 cal.
 (d) 18 cal, 18 cal.
 25. 8000 cal.

Chapter 23

1. 76.5% by mass.
 3. 100 cm^3 .
 5. 653 joules.
 7. 27 lb/in.^2
 9. The mercury drops 41.7 cm .
 11. (a) $565 \times 10^{-23} \text{ joule}$,
 $772 \times 10^{-23} \text{ joule}$.
 (b) 3390 joule, 4630 joule.
 13. (a) $1.01 \times 10^{16} \text{ K}$, $16.2 \times 10^{16} \text{ K}$;
 (b) 450°K , 7200°K .
 15. He: 1400 meters/sec;
 A: 440 meters/sec.

17. 6.6×10^4 .
 19. (a) Eleven times larger.
 (b) Same size.
 25. (a) $6.6 \times 10^{-23} \text{ gm}$.
 (b) 40.
 27. 4.13 joules/cal.
 29. 1910 cal.
 35. 1.41.
 37. Monatomic.
 39. (a) 8.0 atm.
 (b) 600° K .
 41. (a) 2.5 atm, 336° K .
 (b) 0.41 V_i .

Chapter 24

1. $3.2 \times 10^{-8} \text{ cm}$.
 3. (a) $3.5 \times 10^{10} \text{ molecules/cm}^3$.
 (b) 160 meters.
 9. (a) $7.1 \times 10^3 \text{ meters/sec}$.
 (b) $2 \times 10^{-3} \text{ cm}$.
 (c) $5 \times 10^{10} \text{ collisions/sec}$.
 11. $\bar{v}, v_{\text{rms}}, v_p$.
 13. 1.5 cm/sec.
 17. $RT \ln \frac{V_f - b}{V_i - b} + a \left(\frac{1}{V_f} - \frac{1}{V_i} \right)$.
 19. (a) $3.2 \times 10^6 \text{ nt/meter}^2$.
 (b) $4.1 \times 10^6 \text{ nt/meter}^2$.

Chapter 25

1. $5.0 \times 10^4 \text{ joules}$.
 7. (a) 2090 joules.
 (b) 380 cal.
 (c) 1570 joules.
 9. 6.5.
 11. $1.1 \times 10^6 \text{ joules}$.
 15. 10^{-3} N , 10^{-6} N , 0°N , -10^{-6} N ,
 -10^{-3} N , for example.
 17. $+0.20 \text{ cal/}^\circ\text{K}$.
 19. $+0.3 \text{ cal/gm }^\circ\text{K}$.
 21. $+0.1 \text{ cal/}^\circ\text{K}$; no.
 23. (a) 500 meters/sec.
 (b) N_2 .
 (c) Positive.

Index

Absolute thermodynamic temperature
scale, 631
Absolute zero, 532, 632
Acceleration, 39, 57, 80
 angular, 244
 average, 39
 centripetal, 66, 69, 70
 constant, 41, 44, 58
 in free fall, 48
 of gravity, 91, 393, 396
 instantaneous, 40
 radial, 66, 69
 and relative velocity, 71
 tangential, 69
 in uniform circular motion, 65
 variable, 41
Action force, 87
Adhesion, surface, 113
Adiabatic processes, 564
 entropy changes in, 639
Aerodynamic lift, 451
Amontons, G., 112
Amplitude, 346, 469, 501
 of simple harmonic motion, 346, 350
Angular acceleration, 244
Angular frequency, 350
Angular momentum, 299
 conservation of, 305, 310

Angular velocity, 299
Annihilation, 173
Antinode, 504
Archimedes' principle, 430, 450
Aristotle, 48, 52
Astronomical unit, 13
Atmosphere, 433
 of moon, 604
 pressure variation in, 598
 scale height of, 437
Atomic clock, 8
Atomic mass unit, 231
Atwood's machine, 101
Audible range, 497
Austern, N., 80
Avogadro's law, 597
Avogadro's number, 550, 580, 609
Ballistic pendulum, 219
Banked curves, 120
Barometer, 432
Barrell, H., 5
Bascom, W., 463
Beams, J. W., 114
Beats, 510
Benfield, A. E., 299
Bernoulli, Daniel, 447, 572

- Bernoulli's equation, 445
 applications of, Pitot tube, 449, 460
 Venturi meter, 448
- Binding energy, 174
- Blackham, E. D., 506
- Bohr model, 77, 129
- Bohr, N., 314
- Boltzmann, L., 572, 608
- Boltzmann's constant, 641
- Born, Max, 124
- Bose-Einstein distribution, 608
- Bowden, F. P., 112
- Boyle, Robert, 433, 572
- Brahe, T., 385
- British engineering system, 90
- British Thermal Unit, 547, 555
- Brown, R., 608
- Brownian motion, 608
- Brush, S. G., 572
- Bubble chamber, 32, 70, 227
- Bulk modulus, 500
- Buoyancy, 431
- Bureau of Standards, 388
- Calculus, 36, 81
- Calorie, 547, 555
- Carnot cycle, 622, 634
- Carnot, Sadi, 547, 622, 629
- Carnot's theorem, 629
- Cavendish balance, 389
- Cavendish, H., 388
- Celsius degree, 536
- Celsius scale, 533
- Center of gravity, 190, 323
- Center of mass, 183, 323
 motion of, 189
- Center of mass frame, collisions in, 220
- Centrifugal force, 121
- Centripetal acceleration, 66, 69, 70
- Centripetal force, 117
- Cerenkov radiation, 518
- Cesium clock, 8
- Cgs system, 90
- Cgs units, 11
- Chadwick, J., 239
- Chizanowski, P., 388
- Circular motion, radial acceleration in.
 69
 tangential acceleration in, 69
- Classical mechanics, 79, 122
- Clausius, R., 572, 589, 628
- Clausius statement of second law, 639
- Clock, cesium, 8
 quartz crystal, 8
- Cloud chamber, 225, 226
- Colding, L. A., 547
- Collisions, 210
 in center of mass, 220
 in an ideal gas, 575
 inelastic, 218
 momentum conserved in, 213
 in one dimension, 215
 in two dimensions, 223
- Compressibility, 551
- Conservation of angular momentum,
 305, 310
 energy, 150, 170
 linear momentum, 173, 193
 mechanical energy, 168
- Conservative force, 150
 in one dimension, 158
 and path independence, 153
 in two dimensions, 166
- Continuity, equation of, 443
- Conveyer belt problem, 203
- Copernican, system, 48
- Copernican theory, 385
- Copernicus, N., 384
- Coulomb, C. A., 112
- Coulomb's law, 92
- Covolume, 612
- Critical pressure, 423, 614
- Critical temperature, 613
- Critical volume, 614
- Cross product, 24
- Cross section, 227
- Cycle, Carnot, 622
- D'Alambert, J., 222
- Dalton's law, 597
- Debye, P., 550
- Debye temperature, 550
- Degree Celsius, 536
- Densities, some measured, 425
- Density, 424
 weight, 427
- Derivative, 35
- Derjaguim, B. V., 583
- Descartes, René, 222, 433, 460
- Dicke, R. H., 393
- Dimensions, 45
- Dipole flow, 455
- Dipole moment, magnetic, 92
- Dirac, P. A. M., 125
- Disorder, 640
- Displacement vector, 34, 57
- Doppler, C. J., 512
- Doppler effect, 512
- Dot product, 23
- Drumhead, vibrations of, 509

INDEX

- Dry ice puck, 82, 114
 "Dry" water, 442
 Dulong, 544, 550
 Dynamic lift, 449
 Dynamics, fluid, 440
 of particles, 79, 109
 rotational, 295
 of variable-mass systems, 202
 Dyne, 90
 Dyson, F., 640
- Earth, as reference frame, 413
 weight of, 390
 Edgerton, H. E., 211
 Efficiency, of engine, 625, 629
 Eigenfrequency, 508
 Eigenfunction, 509
 Einstein, A., 123, 171, 451, 609
 Eisenbud, L., 80
 Elastic limit, 347
 Electromagnetic force, 120
 Elementary particles, 32
 Emmerich, 532
 Energy, 131
 binding, 174
 conservation of, 150, 170
 equipartition of, 589
 gravitational potential, 406, 410
 and heat, 545
 interatomic, 537, 583
 interval, 560
 mechanical, 156, 168
 of oscillations, 344
 potential, 154
 of simple harmonic motion, 353
 Engine, efficiency of, 629
 heat, 625, 627
 Entropy, 619, 635
 and disorder, 640
 and irreversible processes, 636
 and reversible processes, 633
 and the second law, 638
 Environment, 79, 524
 Eötvös, 393
 Ephemeris time, 8
 Equation of state of ideal gas, 573
 Equilibrium, examples of, 326
 neutral, 164, 333
 of rigid bodies, 320
 saddle point, 334
 stable, 164, 333
 thermal, 525
 thermodynamic, 556
 unstable, 164, 333
 Equipartition of energy, 589
 Equivalence, Eratosthenes, 395
 principle of, 414
 Erg, 134
 Escape velocity, 409
 Essen, L., 7, 10
 Euler, L., 440
 Evaporation, 604
 Expansion, 535
 coefficient of linear, 536, 537
 coefficient of volume, 538
 Fahrenheit scale, 533
 Fermi-Dirac distribution, 608
 Feynman, Richard, 442
 Field, of flow, 453
 Field, gravitational, 404, 453
 Field concept, 405
 First law of thermodynamics, 560
 Fizeau, H. L., 5
 Flettner ship, 451
 Flow, compressible, 441
 dipole, 455
 field of, 453
 incompressible, 441
 irrotational, 441
 nonsteady, 441
 nonviscous, 441
 potential, 453
 rotational, 441, 453
 steady, 441
 streamline, 453
 tube of, 442
 viscous, 441
 vortex, 453
 Fluid dynamics, 440
 Fluid mechanics, conservation of
 momentum in, 452
 Fluid statics, 423
 Force, action, 87
 centrifugal, 121
 centripetal, 117
 classification of, 120
 conservative, 150, 158, 166
 dynamic procedure for measuring, 95
 electromagnetic, 120
 of friction, 109
 gravitational, 120
 inertial, 121
 laws of, 92
 nonconservative, 150, 167
 nuclear, 120
 and potential energy, 157
 pseudo-, 120
 reaction, 87
 static procedure for measuring, 95

- Force, "true" measure of, 222
 Force constant, 138, 346
 Foucault pendulum, 422
 Fps units, 11
 Free-body diagram, 96
 Free expansion, 565, 636, 641
 Free fall, 48
 equations of, 49
 Freitag, E. H., 109
 Frequency, 343
 angular, 659
 Friction, 109
 kinetic, 111
 as nonconservative force, 152
 static, 110
 Fundamental frequency, 505

 g, 48
 Galileo, 48, 51, 52, 59, 75, 81, 82, 384, 385
 Gardner, Martin, 338
 Gas constant, universal, 573
 Gas, ideal, 572
 Gas thermometer, 529
 Gases, kinetic theory of, 571, 599
 Gauge pressure, 432
 Gaussian system, 90
 Gaussian units, 11
 Geiger, H., 230
 Gerjuoy, E., 194, 310
 Gibbs, J. W., 572
 Glaser, D., 225
 Gradient, 167
 Gram, 90
 Gravitation, 382
 constant of, 92, 388
 law of, 92, 387
 Gravitational, field, 404, 453
 force, 120
 mass, 391
 potential energy, 406, 410
 Gravity, acceleration due to, 91, 393, 396
 center of, 190
 screen for, 388
 of a sphere, 397

 Halley, E., 384
 Harmonic motion, 342
 combinations of, 366
 damped, 370
 Harmonic oscillator, 160
 Harmonics, 505
 organ pipe, 507
 Havens, 607
 Hay, G. E., 16

 Heat capacity, 547
 Heat, 169, 545, 640
 conduction of, 551, 637, 641
 mechanical equivalent of, 554
 path dependence of, 559
 of vaporization, 563
 and work, 555
 Heat engine, 625
 Heikes, 532
 Heisenberg, 124
 Henri, V., 610
 Hertz, 343
 Hertz, H., 343
 Heyl, P. R., 388
 Hooke, R., 348, 386
 Hooke's law, 138, 151, 348, 358
 Horsepower, 145
Hydrodynamica, 447
 Hydrostatic paradox, 435

 Iceberg, 431
 Ideal gas, adiabatic process in, 587
 equation of state of, 573
 internal energy of, 584
 macroscopic definition of, 572
 microscopic definition of, 574
 specific heats of, 584, 590
 Ideal gas, diatomic, 590
 monatomic, 590
 polyatomic, 590
 Ideal gas temperature scale, 530
 Impulse, 212
 Inelastic collision, 218
 Inertia, law of, 82
 moment of, 268
 Inertial force, 121
 Inertial frame, 82
 Inertial mass, 391
 Integral, 138
 Interference, 478, 511
 Interferometer, 5
 Intermolecular forces, 164, 582
 Internal energy, 560
 of ideal gas, 584
 International Bureau of Weights and Measures, 83
 Invar, 537
 Invariance, 26, 31
 Irreversible process, 619
 entropy change in, 636
 Isobaric process, 562
 Isotherms, 613
 Isotope separation, 581
 Isotropic, 538

- Jelley, J. V., 518
 joule, 134
 Joule experiment, 567
 Joule, J., 547, 554, 572
- Kelvin degree, 527
 Kelvin, Lord, 1, 628, 629
 Kelvin-Planck statement of second law, 639
 Kelvin temperature scale, 632
 Kepler, J., 385, 401, 460
 Kepler's laws, 460
 Kilocalorie, 547, 555
 thermochemical, 555
 Kilogram, standard, 83, 90
 Kilowatt-hour, 145
 Kinematics, 32
 Kinetic energy, 142, 148, 158
 Kinetic theory of gases, 571, 599
 Kinetic theory of pressure, 575
 Kinetic theory of temperature, 579
 King-Hele, 396
 Kirkpatrick, P., 76
 Koppler, 610
 Kronig, A., 572
 Krypton-86 meter, 6
 Kundt's method, 521
 Kusch, P., 606
- Lagrange, J. L., 440
 Laplace, 589
 Lawrence Radiation Laboratory, 70
 Lee, T. D., 27
 Length, atomic standard of, 6
 Length, standard of, 5
 Lengths, some measured, 4
 Leonardo da Vinci, 112
 Lift, dynamic, 449
 static, 450
 Light year, 13
 Limiting process, 37, 40
 Lindsay, R. B., 171, 444
 Linear momentum, conservation of, 193
 of a particle, 191
 of a system, 192
 Linear motion with constant acceleration, 41, 44
 Lissajous figures, 379
 Locke, J., 525
 Longitudinal waves, 464
- McNish, A. G., 11
 Mach, E., 608
- Macroscopic description, 524
 Magdeburg hemisphères, 436
 Magnetic dipole moment, 92
 Marsden, 230
 Mass, 80, 85
 center of, 183
 and energy, 171
 gravitational, 391
 inertial, 391
 reduced, 368
 relativistic, 171, 192
 rest, 171
 variable, 198
 and weight, 93
- Mass-energy equivalence, 173, 232
 Masses, some measured, 86
 Maxwell, C., 572, 590, 602, 605
 Maxwellian distribution, 602, 605
 Mean free path, 599
 Measurement, 1
 Mechanical energy, 156, 168
 Mechanical equivalent of heat, 554
 Mechanics, 32
 classical, 79, 122
 quantum, 122
 relativistic, 122
 relativistic quantum, 125
 Mercury, 404
 Merriam, J. L., 202
 Meter, standard, 5
 Microscopic description, 524
 Miller, F. H., 371
 Mks system, 90
 Mks units, 11
 Molar heat capacity, 550
 Mole, 550
 Molecular weight, 550
 Molecules, diameter of, 600
 Moment of inertia, 268
 Moment of mass, 187
 Momentum, 212
 angular, 299, 305, 310
 conservation of, 195, 213, 452
 linear, 183, 191, 192
 Moon, atmosphere of, 604
 Moore, A. D., 455, 456
 Morse, P. M., 572
 Motion, in one dimension, 32, 36, 41
 in a plane, 57
 of projectiles, 59
 rectilinear, 36
 of translation, 33
 uniform circular, 64
 Mott, N. F., 594
 Multiples and supmultiples-metric, 12

- National Bureau of Standards, 83
 Negative temperature, 532
 Neutron, discovery of, 239
 newton, 84, 90
 Newton, I., 36, 80, 81, 82, 112, 191, 382,
 385, 460, 589
 Newton's first law, 81, 121
 Newton's law of cooling, 541
 Newton's law of gravitation, 92
 Newton's laws of motion, applications
 of, 96
 Newton's second law, 85
 Newton's third law, 87, 194
 Node, 504
 Nonconservative force, 150, 167
 Noninertial frame, 121
 Nuclear force, 120
 Nucleus, discovery of, 230
- Oliver, Jack, 497
 Operational viewpoint, 2
 Organ pipe, harmonics of, 507
 Oscillations, 342
 energy of, 344
 forced, 372
 frequency of, 343
 life-time of, 371
 period of, 343
 simple harmonic, 345
 two-body, 368
 Ostwald, W., 608
 Overtones, 505
 Owens, Jesse, 75
- Pair production, 173
 Panofsky, W., 643
 Parsec, 13
 Particle, 32, 79
 Particle dynamics, 79, 109
 Particle kinematics, 32
 Pascal, Blaise, 430, 433
 Pascal's principle, 430, 577
 Path independence, 153
 Pendulum, ballistic, 219
 conical, 118
 Foucault, 422
 physical, 360
 simple, 140, 167, 357
 torsional, 358
 Period, 343, 459
 Periodic motion, 342
 Perrin, J., 610
 Petit, 500, 544
 Phase change, 487
 Phase constant, 350
 Phase of simple harmonic motion, 350
 Phillips, M., 643
 Piano, waveform of, 506
 Piezoelectric effect, 497
 Pitot tube, 449, 460
 Planck, M., 625
 Planck's constant, 124
 Plane, motion in, 57
 Plane wave, 466
 Planets, 341
 energy of, 412
 Plasma, 423
 Poiseuille, 462
 Pomeranz, K. B., 203
 Potential energy, 154
 intermolecular, 164, 537
 and work, 161
 Potential flow, 453
 Pound, 90
 Power, 145
 in wave motion, 476
 Poynting, 388
 Prefixes, metric, 12
 Pressure amplitude, 502
 Pressure, 424
 of atmosphere, 428
 in fluid at rest, 425
 gauge, 432
 kinetic theory of, 575
Principia, 81, 191, 384
 Principle of equivalence, 414
 Process, adiabatic, 564
 irreversible, 619
 isobaric, 562
 quasi-static, 621
 reversible, 619
 thermodynamic, 556
 Projectile motion, 59
 Pseudo-force, 82, 120
 Ptolemaic system, 48
 Ptolemy, 384
- Quantities, physical, 2
 Quantum mechanics, 122
 relativistic, 125
 Quantum theory, 593
 Quartz crystal clock, 8
 Quasi-static process, 621
- Rabinowicz, Ernest, 110
 Radioactive decay, 196, 212
 Rainwater, 607
 Ramsey, N. F., 625
 Ray, 466
 Rayleigh, Lord, 581

INDEX

- Hayne, J., 532
Reaction, 211
Reaction force, 87
Reactions, nuclear, 231
Rectilinear motion, 36
Reddick, H. W., 371
Reduced mass, 368
Reference frame, 3, 120
 absolute, 4
 earth as, 413
 inertial, 4
Reference frames, and kinetic energy, 145
 and relative velocity, 71
 and work, 145
Refrigerator, 625, 627
 coefficient of performance of, 644
Rekveld, J., 484
Relative velocity and acceleration, 71
Relativistic mass, 171, 192
Relativistic mechanics, 122
Relativistic quantum mechanics, 125
Relativity theory, 3, 171
 general, 404
Resonance, 372, 488
Rest mass, 171
Reversible processes, 619
 entropy for, 633
Rezneck, S., 555
Ricci, M. A., 433
Rigid body, equilibrium of, 320
Rocket, 201
 thrust on, 451
Rogers, W. E., 455, 456
Root-mean-square speed, 577
Rotational dynamics, 295, 811
Rotational flow, 453
Rotational inertia, 268
Rotor, 119
Rowland, H. A., 555
Ruⁿford, Count, 546, 567
Russel, B., 384
Rutherford, Lord, 230
- Saddle point, 334
Satellite, 67, 396, 401
 energy of, 412
Scalar product, 23, 30, 134
Scalars, 16
Schrödinger, 124
Second law of motion, 85
Shock wave, 517
Simple harmonic motion, 345, 348
 amplitude of, 346, 350
 angular, 359
Simple harmonic motion, applications of, 357
 energy of, 353
 equation of motion of, 347, 348
 frequency of, 350
 period of, 350
 phase constant of, 350
 phase of, 350
 two-body, 368
 uniform circular motion and, 363
Simple harmonic oscillator, 345
Sink, 455
Slater, J. C., 560
Slug, 91
Smyth, H. D., 581
Sommerfeld, A., 303
Sound, sources of, 505
 speed of, 500, 509, 521, 587
Sound waves, 497
Source, 455
Specific heat, 547
 at constant pressure, 549, 586
 at constant volume, 549, 586
 of hydrogen, 542
 of an ideal gas, 584
Specific heats, ratio of, 501, 590
Speed, average, 603
 molecular, 602
 most probable, 603
 root-mean-square, 577, 603
 of sound, 521
 of waves, 472
Spherical wave, 466
Standard, of time, 7
Standards, 2
Standing wave, 483, 486, 489
Static friction, 110
Statics, fluid, 423
Statistical mechanics, 525, 571
 at negative temperatures, 628
Stern, O., 605
Streamlines, 442, 454
Stukeley, 382
Superconductivity, 552
Superposition principle, 470
Surface adhesion, 113
System, 524
- Tabor, D., 112
Tacoma Narrows Bridge, 374
Temperature, 524, 526, 532, 561
 absolute zero of, 632
 Celsius scale of, 533
 Dobye, 550
 Fahrenheit scale of, 533

- Temperature, fixed points of, 535
 ideal gas scale of, 530
 International Practical Scale of, 534
 Kelvin scale of, 632
 and kinetic theory, 579
 negative, 532, 628
 thermodynamic scale of, 631
- Temperature expansion, 535
- Temperature gradient, 552
- Temperatures, some measured, 533
- Terminal velocity, 107
- Thermal conductivity, 552
- Thermal equilibrium, 525
- Thermal expansion, coefficient of, 551
- Thermodynamic process, 556
- Thermodynamic temperature scale, 631
- Thermodynamics, 524, 556, 571
 first law of, 560, 561, 627, 633
 at negative temperatures, 628
 second law of, 619, 626, 633, 638, 639
 third law of, 633
 zeroth law of, 525, 633
- Thermometer, constant volume gas, 529
- Third law of motion, 87
- Thompson, B., 545
- Thomsen, J. S., 526, 621
- Thomson, W., 1, 629
- Thrust, 200
- Tides, 376
- Time, ephemeris, 8
 standard of, 7
 universal, 8
- Time intervals, some measured, 7
- Tolman, R. C., 525
- Top, 296
- Torricelli, E., 432
- Torricelli's law, 459
- Translational motion, 33
- Transverse waves, 464
- Triple point, 527, 533
- Tube of flow, 442
- Uncertainty relation, 124
- Uniform circular motion, 64, 116, 363
- Units, 2, 11, 45
 British engineering, 90
 cgs, 90
 Gaussian, 90
 mechanical, 90
 mks, 90
- Unit vectors, 19, 36
 in polar coordinates, 67
- U.S. Standard Atmosphere, 40
- Universal gas constant, 573
- Universal time, 8
- Vectors, 15
 addition of, 16, 18
 components of, 18
 and laws of physics, 26
 multiplication of, 23
 resolution of, 18
 subtraction of, 17
- Vector product, 24, 30
- Velocity, 57
 angular, 299
 average, 33
 escape, 409
 instantaneous, 35
 relative, 71
 terminal, 107
 variable, 36
- Venturi meter, 448
- Venturi tube, 458
- Vibrating systems, 505
- Vibration, of string, 506
- Violin, waveform of, 506
- von Guericke, Otto, 433, 436
- von Helmholtz, H., 547
- von Mayer, J., 547
- von Neuman, John, 442
- Vortex flow, 453
- van der Waals' equation, 611
- van der Waals, J. D., 611
- WWV, 11
- Water, density of, 539
 specific heat of, 547
- Waterston, J. J., 572
- Watt, 145
- Watt, J., 145
- Wave, intensity of, 503
 plane, 466
 spherical, 466
 standing, 486
 traveling, 467
- Wave speed, 427
- Wavefront, 466
- Wavelength, 459
- Waves, audible, 497
 complex, 481
 elastic, 463
 infrasonic, 497
 interference of, 478
 longitudinal, 464, 498, 501
 mechanical, 463
 power in, 476
 sound, 497
 standing, 483, 489, 504
 transverse, 464
 ultrasonic, 497

- Weight and mass, 93
Weight density, 427
Weinstock, R., 80
Wilson, C. T. R., 225
Work, 131
 as an area on a p - V diagram, 138, 559
 by a constant force, 132
 and heat, 555
 path dependence of, 559
 and reference frame, 148
- Work, as a scalar product, 134
 by a variable force, 136, 139
Work-energy theorem, 141 144, 150
 in fluid flow, 445
- Yang, C. N., 27
Young's modulus, 501
- Zeroth law of thermodynamics, 525

SELECTED NUMERICAL CONSTANTS

$$\pi = 3.14$$

$$\pi^2 = 9.87$$

$$e = 2.72$$

$$e^{-1} = 1/e = 0.368$$

$$\ln 2 = 0.693$$

$$\log e = 0.434$$

$$\sqrt{2} = 1.41$$

$$\sqrt{3} = 1.73$$

$$\sin 30^\circ = \cos 60^\circ = \frac{1}{2} = 0.500$$

$$\cos 30^\circ = \sin 60^\circ = \sqrt{3}/2 = 0.866$$

$$\tan 30^\circ = \cot 60^\circ = \sqrt{3}/3 = 0.577$$

$$\tan 60^\circ = \cot 30^\circ = \sqrt{3} = 1.732$$

$$\sin 45^\circ = \cos 45^\circ = \sqrt{2}/2 = 0.707$$

$$\tan 45^\circ = \cot 45^\circ = 1.00$$

SELECTED CONVERSION FACTORS

(See Appendix H for a more complete list)

$$180^\circ = \pi \text{ rad}$$

$$1 \text{ radian} = 57.3^\circ = 0.159 \text{ rev}$$

$$1 \text{ slug} = 32.2 \text{ lb (mass)} = 14.6 \text{ kg}$$

$$1 \text{ kilogram} = 2.21 \text{ lb (mass)}$$

$$1 \text{ pound (mass)} = 0.454 \text{ kg}$$

$$1 \text{ atomic mass unit} = 1.66 \times 10^{-27} \text{ kg}$$

$$1 \text{ meter} = 39.4 \text{ in.} = 3.28 \text{ ft; } 1 \text{ inch} = 2.54 \text{ cm}$$

$$1 \text{ mile} = 5280 \text{ ft} = 1.61 \text{ km}$$

$$1 \text{ angstrom unit} = 10^{-10} \text{ meter} = 0.1 \text{ m}\mu$$

$$1 \text{ millimicron} = 10^{-9} \text{ meter}$$

$$1 \text{ liter} = 61.0 \text{ in.}^3$$

$$1 \text{ ft}^3 = 28.3 \text{ li}$$

$$1 \text{ day} = 86,400 \text{ sec}$$

$$1 \text{ year} = 3.16 \times 10^7 \text{ sec} = 365 \text{ days}$$

$$1 \text{ mile/hr} = 1.47 \text{ ft/sec} = 0.447 \text{ meter/sec}$$

$$1 \text{ pound} = 4.45 \text{ nt; } 1 \text{ newton} = 0.225 \text{ lb}$$

$$1 \text{ atmosphere} = 29.9 \text{ in.-Hg} = 76.0 \text{ cm-Hg} = 1.01 \times 10^5 \text{ nt/meter}^2$$

$$1 \text{ Btu} = 778 \text{ ft-lb} = 252 \text{ cal} = 1060 \text{ joules}$$

$$1 \text{ calorie} = 4.19 \text{ joules; } 1 \text{ joule} = 0.239 \text{ cal} = 2.78 \times 10^{-7} \text{ kw-hr}$$

$$1 \text{ electron volt} = 1.60 \times 10^{-19} \text{ joule}$$

$$1 \text{ horsepower} = 550 \text{ ft-lb/sec} = 746 \text{ watts}$$

$$1 \text{ weber/meter}^2 = 1 \text{ tesla} = 10^4 \text{ gauss}$$